

wwPDB X-ray Structure Validation Summary Report (i)

Sep 12, 2023 – 04:11 AM EDT

PDB ID : 4LY4

Title : Crystal structure of peptidoglycan deacetylase (HP0310) with Zinc from Heli-

cobacter pylori

Authors: Shaik, M.M.; Zanotti, G.

Deposited on : 2013-07-30

Resolution : 2.20 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at

https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity : 4.02b-467 Xtriage (Phenix) : 1.13

EDS : 2.35.1

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

Refmac : 5.8.0158

CCP4 : 7.0.044 (Gargrove)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.35.1

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive	Similar resolution
Metric	$(\# ext{Entries})$	$(\# ext{Entries}, ext{ resolution range}(ext{Å}))$
R_{free}	130704	4898 (2.20-2.20)
Clashscore	141614	5594 (2.20-2.20)
Ramachandran outliers	138981	5503 (2.20-2.20)
Sidechain outliers	138945	5504 (2.20-2.20)
RSRZ outliers	127900	4800 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain					
1	A	326	74% 12%	•	11%			
1	В	326	71% 16%	•	11%			
1	С	326	75% 13%	•	11%			
1	D	326	74%	•	11%			



2 Entry composition (i)

There are 3 unique types of molecules in this entry. The entry contains 9758 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called peptidoglycan deacetylase.

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace		
1	Λ	290	Total	С	N	О	S	0	0	0
1	A	290	2357	1530	388	428	11	U	0	0
1	В	290	Total	С	N	О	S	0	0	0
1	Б	290	2357	1530	388	428	11	U	0	U
1	С	290	Total	С	N	О	S	0	0	0
1		290	2357	1530	388	428	11	U	0	0
1	D	290	Total	С	N	О	S	0	0	0
1	ש	290	2357	1530	388	428	11	U	U	U

There are 132 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-32	MET	-	expression tag	UNP B5ZA76
A	-31	HIS	-	expression tag	UNP B5ZA76
A	-30	HIS	-	expression tag	UNP B5ZA76
A	-29	HIS	-	expression tag	UNP B5ZA76
A	-28	HIS	-	expression tag	UNP B5ZA76
A	-27	HIS	-	expression tag	UNP B5ZA76
A	-26	HIS	-	expression tag	UNP B5ZA76
A	-25	GLY	-	expression tag	UNP B5ZA76
A	-24	LYS	-	expression tag	UNP B5ZA76
A	-23	PRO	-	expression tag	UNP B5ZA76
A	-22	ILE	-	expression tag	UNP B5ZA76
A	-21	PRO	_	expression tag	UNP B5ZA76
A	-20	ASN	-	expression tag	UNP B5ZA76
A	-19	PRO	_	expression tag	UNP B5ZA76
A	-18	LEU	-	expression tag	UNP B5ZA76
A	-17	LEU	-	expression tag	UNP B5ZA76
A	-16	GLY	-	expression tag	UNP B5ZA76
A	-15	LEU	-	expression tag	UNP B5ZA76
A	-14	ASP		expression tag	UNP B5ZA76
A	-13	SER	-	expression tag	UNP B5ZA76
A	-12	THR	-	expression tag	UNP B5ZA76



 $Continued\ from\ previous\ page...$

Chain	Residue	Modelled	Actual	Comment	Reference
A	-11	GLU	-	expression tag	UNP B5ZA76
A	-10	ASN	-	expression tag	UNP B5ZA76
A	-9	LEU	-	expression tag	UNP B5ZA76
A	-8	TYR	-	expression tag	UNP B5ZA76
A	-7	PHE	-	expression tag	UNP B5ZA76
A	-6	GLN	-	expression tag	UNP B5ZA76
A	-5	GLY	-	expression tag	UNP B5ZA76
A	-4	ILE	-	expression tag	UNP B5ZA76
A	-3	ASP	-	expression tag	UNP B5ZA76
A	-2	PRO	-	expression tag	UNP B5ZA76
A	-1	PHE	-	expression tag	UNP B5ZA76
A	0	THR	-	expression tag	UNP B5ZA76
В	-32	MET	-	expression tag	UNP B5ZA76
В	-31	HIS	-	expression tag	UNP B5ZA76
В	-30	HIS	-	expression tag	UNP B5ZA76
В	-29	HIS	-	expression tag	UNP B5ZA76
В	-28	HIS	-	expression tag	UNP B5ZA76
В	-27	HIS	-	expression tag	UNP B5ZA76
В	-26	HIS	-	expression tag	UNP B5ZA76
В	-25	GLY	-	expression tag	UNP B5ZA76
В	-24	LYS	-	expression tag	UNP B5ZA76
В	-23	PRO	_	expression tag	UNP B5ZA76
В	-22	ILE	-	expression tag	UNP B5ZA76
В	-21	PRO	-	expression tag	UNP B5ZA76
В	-20	ASN	-	expression tag	UNP B5ZA76
В	-19	PRO	-	expression tag	UNP B5ZA76
В	-18	LEU	-	expression tag	UNP B5ZA76
В	-17	LEU	-	expression tag	UNP B5ZA76
В	-16	GLY	-	expression tag	UNP B5ZA76
В	-15	LEU	-	expression tag	UNP B5ZA76
В	-14	ASP	-	expression tag	UNP B5ZA76
В	-13	SER	-	expression tag	UNP B5ZA76
В	-12	THR	-	expression tag	UNP B5ZA76
В	-11	GLU	-	expression tag	UNP B5ZA76
В	-10	ASN	-	expression tag	UNP B5ZA76
В	-9	LEU	-	expression tag	UNP B5ZA76
В	-8	TYR	-	expression tag	UNP B5ZA76
В	-7	PHE	-	expression tag	UNP B5ZA76
В	-6	GLN	-	expression tag	UNP B5ZA76
В	-5	GLY	-	expression tag	UNP B5ZA76
В	-4	ILE	-	expression tag	UNP B5ZA76
В	-3	ASP	-	expression tag	UNP B5ZA76



 $Continued\ from\ previous\ page...$

Chain	Residue	Modelled Modelled	Actual	Comment	Reference
В	-2	PRO	-	expression tag	UNP B5ZA76
В	-1	PHE	-	expression tag	UNP B5ZA76
В	0	THR	-	expression tag	UNP B5ZA76
С	-32	MET	-	expression tag	UNP B5ZA76
С	-31	HIS	-	expression tag	UNP B5ZA76
С	-30	HIS	_	expression tag	UNP B5ZA76
С	-29	HIS	-	expression tag	UNP B5ZA76
С	-28	HIS	-	expression tag	UNP B5ZA76
С	-27	HIS	-	expression tag	UNP B5ZA76
С	-26	HIS	-	expression tag	UNP B5ZA76
С	-25	GLY	-	expression tag	UNP B5ZA76
С	-24	LYS	-	expression tag	UNP B5ZA76
С	-23	PRO	-	expression tag	UNP B5ZA76
С	-22	ILE	-	expression tag	UNP B5ZA76
С	-21	PRO	-	expression tag	UNP B5ZA76
С	-20	ASN	-	expression tag	UNP B5ZA76
С	-19	PRO	-	expression tag	UNP B5ZA76
С	-18	LEU	_	expression tag	UNP B5ZA76
С	-17	LEU	-	expression tag	UNP B5ZA76
С	-16	GLY	-	expression tag	UNP B5ZA76
С	-15	LEU	-	expression tag	UNP B5ZA76
С	-14	ASP	-	expression tag	UNP B5ZA76
С	-13	SER	-	expression tag	UNP B5ZA76
С	-12	THR	-	expression tag	UNP B5ZA76
С	-11	GLU	-	expression tag	UNP B5ZA76
С	-10	ASN	-	expression tag	UNP B5ZA76
С	-9	LEU	-	expression tag	UNP B5ZA76
С	-8	TYR	-	expression tag	UNP B5ZA76
С	-7	PHE	-	expression tag	UNP B5ZA76
С	-6	GLN	_	expression tag	UNP B5ZA76
С	-5	GLY	-	expression tag	UNP B5ZA76
С	-4	ILE	-	expression tag	UNP B5ZA76
С	-3	ASP	-	expression tag	UNP B5ZA76
С	-2	PRO	-	expression tag	UNP B5ZA76
С	-1	PHE	-	expression tag	UNP B5ZA76
С	0	THR	_	expression tag	UNP B5ZA76
D	-32	MET		expression tag	UNP B5ZA76
D	-31	HIS	-	expression tag	UNP B5ZA76
D	-30	HIS	-	expression tag	UNP B5ZA76
D	-29	HIS	-	expression tag	UNP B5ZA76
D	-28	HIS	-	expression tag	UNP B5ZA76
D	-27	HIS	-	expression tag	UNP B5ZA76



 $Continued\ from\ previous\ page...$

Chain	Residue	Modelled	Actual	Comment	Reference
D	-26	HIS	-	expression tag	UNP B5ZA76
D	-25	GLY	-	expression tag	UNP B5ZA76
D	-24	LYS	-	expression tag	UNP B5ZA76
D	-23	PRO	-	expression tag	UNP B5ZA76
D	-22	ILE	-	expression tag	UNP B5ZA76
D	-21	PRO	-	expression tag	UNP B5ZA76
D	-20	ASN	-	expression tag	UNP B5ZA76
D	-19	PRO	-	expression tag	UNP B5ZA76
D	-18	LEU	-	expression tag	UNP B5ZA76
D	-17	LEU	-	expression tag	UNP B5ZA76
D	-16	GLY	-	expression tag	UNP B5ZA76
D	-15	LEU	-	expression tag	UNP B5ZA76
D	-14	ASP	-	expression tag	UNP B5ZA76
D	-13	SER	-	expression tag	UNP B5ZA76
D	-12	THR	-	expression tag	UNP B5ZA76
D	-11	GLU	-	expression tag	UNP B5ZA76
D	-10	ASN	-	expression tag	UNP B5ZA76
D	-9	LEU	-	expression tag	UNP B5ZA76
D	-8	TYR	-	expression tag	UNP B5ZA76
D	-7	PHE	-	expression tag	UNP B5ZA76
D	-6	GLN	-	expression tag	UNP B5ZA76
D	-5	GLY	-	expression tag	UNP B5ZA76
D	-4	ILE		expression tag	UNP B5ZA76
D	-3	ASP		expression tag	UNP B5ZA76
D	-2	PRO		expression tag	UNP B5ZA76
D	-1	PHE	-	expression tag	UNP B5ZA76
D	0	THR	-	expression tag	UNP B5ZA76

• Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total Zn 1 1	0	0
2	В	1	Total Zn 1 1	0	0
2	С	1	Total Zn 1 1	0	0
2	D	1	Total Zn 1 1	0	0

• Molecule 3 is water.




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	67	Total O 67 67	0	0
3	В	88	Total O 88 88	0	0
3	С	79	Total O 79 79	0	0
3	D	92	Total O 92 92	0	0




3 Residue-property plots (i)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: peptidoglycan deacetylase






• Molecule 1: peptidoglycan deacetylase



• Molecule 1: peptidoglycan deacetylase







4 Data and refinement statistics (i)

Property	Value	Source
Space group	I 2 2 2	Depositor
Cell constants	113.64Å 153.93Å 153.88Å	Depositor
a, b, c, α , β , γ	90.00° 90.00° 90.00°	Depositor
Resolution (Å)	30.48 - 2.20	Depositor
Resolution (A)	30.58 - 2.20	EDS
% Data completeness	99.9 (30.48-2.20)	Depositor
(in resolution range)	98.9 (30.58-2.20)	EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	2.28 (at 2.20Å)	Xtriage
Refinement program	PHENIX (phenix.refine: dev_1352)	Depositor
D.D.	0.169 , 0.207	Depositor
R, R_{free}	0.168 , 0.202	DCC
R_{free} test set	3431 reflections $(5.05%)$	wwPDB-VP
Wilson B-factor (Å ²)	28.1	Xtriage
Anisotropy	0.394	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$, $B_{sol}(Å^2)$	0.33 , 13.5	EDS
L-test for twinning ²	$< L > = 0.36, < L^2> = 0.19$	Xtriage
Estimated twinning fraction	0.457 for -h,-l,-k	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	9758	wwPDB-VP
Average B, all atoms (Å ²)	25.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 8.69% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of <|L|>, $<L^2>$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Mol Chain		lengths	Bond angles	
IVIOI			RMSZ $ \# Z > 5$		# Z > 5
1	A	0.45	0/2432	0.58	0/3305
1	В	0.44	0/2432	0.57	0/3305
1	С	0.43	0/2432	0.58	0/3305
1	D	0.44	0/2432	0.58	0/3305
All	All	0.44	0/9728	0.58	0/13220

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2357	0	2284	33	0
1	В	2357	0	2284	48	0
1	С	2357	0	2284	29	0
1	D	2357	0	2284	35	0
2	A	1	0	0	0	0
2	В	1	0	0	0	0
2	С	1	0	0	0	0
2	D	1	0	0	0	0
3	A	67	0	0	3	0



Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	В	88	0	0	2	0
3	С	79	0	0	3	0
3	D	92	0	0	3	0
All	All	9758	0	9136	133	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

The worst 5 of 133 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	$\begin{array}{c} {\rm Interatomic} \\ {\rm distance} \ ({\rm \AA}) \end{array}$	Clash overlap (Å)
1:A:212:ASN:ND2	1:B:210:SER:O	1.98	0.96
1:A:210:SER:O	1:C:212:ASN:ND2	2.04	0.90
1:A:141:HIS:O	1:A:143:PHE:N	2.10	0.84
1:B:212:ASN:ND2	1:D:210:SER:O	2.13	0.82
1:B:141:HIS:O	1:B:143:PHE:N	2.16	0.78

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
1	A	288/326 (88%)	269 (93%)	15 (5%)	4 (1%)	11	8
1	В	288/326 (88%)	272 (94%)	14 (5%)	2 (1%)	22	22
1	С	288/326 (88%)	272 (94%)	13 (4%)	3 (1%)	15	14
1	D	288/326 (88%)	270 (94%)	16 (6%)	2 (1%)	22	22
All	All	1152/1304 (88%)	1083 (94%)	58 (5%)	11 (1%)	15	14

5 of 11 Ramachandran outliers are listed below:



Mol	Chain	Res	Type
1	A	142	GLY
1	В	142	GLY
1	С	142	GLY
1	D	142	GLY
1	A	131	SER

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric Outliers		Percentiles	
1	A	254/287 (88%)	247 (97%)	7 (3%)	43 56	
1	В	254/287 (88%)	247 (97%)	7 (3%)	43 56	
1	С	254/287 (88%)	249 (98%)	5 (2%)	55 69	
1	D	254/287 (88%)	249 (98%)	5 (2%)	55 69	
All	All	1016/1148 (88%)	992 (98%)	24 (2%)	49 62	

5 of 24 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	С	127	TRP
1	С	213	SER
1	С	190	VAL
1	С	256	VAL
1	A	256	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.



5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

Of 4 ligands modelled in this entry, 4 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ $>$	# RSRZ > 2	$OWAB(Å^2)$	Q<0.9
1	A	290/326 (88%)	-0.58	2 (0%) 87 86	16, 24, 39, 61	0
1	В	290/326 (88%)	-0.61	1 (0%) 94 93	15, 24, 37, 62	0
1	С	290/326~(88%)	-0.62	2 (0%) 87 86	14, 23, 39, 61	0
1	D	290/326 (88%)	-0.61	2 (0%) 87 86	16, 23, 38, 53	0
All	All	1160/1304 (88%)	-0.61	7 (0%) 89 88	14, 24, 39, 62	0

The worst 5 of 7 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	141	HIS	4.5
1	В	141	HIS	3.6
1	С	141	HIS	3.4
1	С	209	LYS	3.3
1	D	142	GLY	2.3

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.



Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-}\mathbf{factors}(\mathbf{\mathring{A}}^2)$	Q<0.9
2	ZN	В	301	1/1	0.99	0.07	36,36,36,36	0
2	ZN	С	301	1/1	0.99	0.04	38,38,38,38	0
2	ZN	A	301	1/1	1.00	0.06	34,34,34,34	0
2	ZN	D	301	1/1	1.00	0.06	33,33,33,33	0

6.5 Other polymers (i)

There are no such residues in this entry.

