



wwPDB X-ray Structure Validation Summary Report ⓘ

Nov 9, 2024 – 07:20 AM EST

PDB ID : 4ZPT
Title : Structure of MERS-Coronavirus Spike Receptor-binding Domain (England1 Strain) in Complex with Vaccine-Elicited Murine Neutralizing Antibody D12 (Crystal Form 1)
Authors : Joyce, M.G.; Mascola, J.R.; Graham, B.S.; Kwong, P.D.
Deposited on : 2015-05-08
Resolution : 2.59 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 1.20.1
EDS : 3.0
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4 : 9.0.003 (Gargrove)
Density-Fitness : 1.0.11
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.39


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.59 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	3775 (2.60-2.60)
Clashscore	180529	4181 (2.60-2.60)
Ramachandran outliers	177936	4129 (2.60-2.60)
Sidechain outliers	177891	4129 (2.60-2.60)
RSRZ outliers	164620	3775 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	216	
1	H	216	
2	B	214	
2	L	214	
3	R	208	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	S	208	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	NAG	R	601	X	-	-	-
4	NAG	S	601	X	-	-	-

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 10054 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called D12 Fab Heavy chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	213	Total 1602	C 1008	N 263	O 322	S 9	0	0	0
1	H	214	Total 1605	C 1010	N 263	O 323	S 9	0	0	0

- Molecule 2 is a protein called D12 Fab Light chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	212	Total 1651	C 1023	N 282	O 340	S 6	0	0	0
2	L	212	Total 1651	C 1023	N 282	O 340	S 6	0	0	0

- Molecule 3 is a protein called Spike glycoprotein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	R	208	Total 1611	C 1029	N 256	O 315	S 11	0	0	0
3	S	208	Total 1611	C 1029	N 256	O 315	S 11	0	0	0

- Molecule 4 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: C₈H₁₅NO₆).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	R	1	Total	C	N	O	0	0
			14	8	1	5		
4	S	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 5 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	75	Total	O	0	0
			75	75		
5	B	55	Total	O	0	0
			55	55		
5	R	29	Total	O	0	0
			29	29		
5	H	57	Total	O	0	0
			57	57		
5	S	19	Total	O	0	0
			19	19		
5	L	60	Total	O	0	0
			60	60		



- Molecule 3: Spike glycoprotein



- Molecule 3: Spike glycoprotein



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	74.45Å 128.79Å 170.87Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	37.05 – 2.59 37.05 – 2.59	Depositor EDS
% Data completeness (in resolution range)	98.7 (37.05-2.59) 98.7 (37.05-2.59)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.21 (at 2.58Å)	Xtrriage
Refinement program	PHENIX 1.9_1692	Depositor
R, R_{free}	0.204 , 0.244 0.211 , 0.242	Depositor DCC
R_{free} test set	2529 reflections (4.94%)	wwPDB-VP
Wilson B-factor (Å ²)	40.9	Xtrriage
Anisotropy	0.211	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 43.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	10054	wwPDB-VP
Average B, all atoms (Å ²)	41.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.11% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section:
NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.68	1/1640 (0.1%)	0.64	2/2236 (0.1%)
1	H	0.72	1/1643 (0.1%)	0.68	1/2239 (0.0%)
2	B	0.66	1/1685 (0.1%)	0.59	1/2290 (0.0%)
2	L	0.74	1/1685 (0.1%)	0.63	2/2290 (0.1%)
3	R	0.58	0/1651	0.54	0/2254
3	S	0.55	0/1651	0.59	0/2254
All	All	0.66	4/9955 (0.0%)	0.61	6/13563 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	H	0	1
2	L	0	1
All	All	0	2

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	101	ASP	CB-CG	-6.48	1.38	1.51
1	H	103	TRP	NE1-CE2	-5.82	1.29	1.37
2	L	50	TYR	CE1-CZ	-5.32	1.31	1.38
2	B	35	TRP	NE1-CE2	-5.28	1.30	1.37

The worst 5 of 6 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	H	148	GLU	C-N-CD	-10.23	98.09	120.60

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	101	ASP	CB-CG-OD1	-9.23	109.99	118.30
1	A	101	ASP	CB-CA-C	-6.83	96.75	110.40
2	L	48	ILE	CG1-CB-CG2	-5.42	99.48	111.40
2	L	18	ARG	NE-CZ-NH2	5.33	122.97	120.30

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	H	148	GLU	Peptide
2	L	211	ARG	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1602	0	1564	15	0
1	H	1605	0	1563	31	0
2	B	1651	0	1584	12	0
2	L	1651	0	1584	29	0
3	R	1611	0	1572	22	0
3	S	1611	0	1570	51	0
4	R	14	0	13	4	0
4	S	14	0	13	2	0
5	A	75	0	0	0	0
5	B	55	0	0	0	0
5	H	57	0	0	1	0
5	L	60	0	0	0	0
5	R	29	0	0	1	0
5	S	19	0	0	0	0
All	All	10054	0	9463	154	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

The worst 5 of 154 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:R:587:LYS:HE3	4:R:601:NAG:O6	1.10	1.22
3:R:587:LYS:CE	4:R:601:NAG:O6	1.89	1.21
3:S:412:THR:O	3:S:416:SER:OG	1.62	1.17
1:H:213:ARG:CG	1:H:213:ARG:HH21	1.58	1.12
3:R:383:CYS:SG	3:R:407:CYS:CB	2.38	1.11

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	209/216 (97%)	205 (98%)	4 (2%)	0	100	100
1	H	210/216 (97%)	206 (98%)	2 (1%)	2 (1%)	13	29
2	B	210/214 (98%)	205 (98%)	5 (2%)	0	100	100
2	L	210/214 (98%)	205 (98%)	4 (2%)	1 (0%)	25	47
3	R	206/208 (99%)	196 (95%)	10 (5%)	0	100	100
3	S	206/208 (99%)	199 (97%)	7 (3%)	0	100	100
All	All	1251/1276 (98%)	1216 (97%)	32 (3%)	3 (0%)	44	66

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	H	101	ASP
1	H	149	PRO
2	L	199	LYS

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar

resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	184/186 (99%)	178 (97%)	6 (3%)	33 59
1	H	183/186 (98%)	167 (91%)	16 (9%)	8 17
2	B	190/192 (99%)	183 (96%)	7 (4%)	29 55
2	L	190/192 (99%)	180 (95%)	10 (5%)	19 40
3	R	190/190 (100%)	176 (93%)	14 (7%)	11 24
3	S	190/190 (100%)	173 (91%)	17 (9%)	8 17
All	All	1127/1136 (99%)	1057 (94%)	70 (6%)	15 33

5 of 70 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
3	S	581	THR
3	S	587	LYS
2	L	105	GLU
3	R	565	GLU
3	R	543	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
3	S	486	HIS
3	S	566	GLN
3	S	582	ASN
2	L	37	GLN
2	L	89	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	NAG	R	601	3	14,14,15	0.28	0	17,19,21	0.57	0
4	NAG	S	601	3	14,14,15	0.29	0	17,19,21	0.57	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	R	601	3	1/1/5/7	4/6/23/26	0/1/1/1
4	NAG	S	601	3	1/1/5/7	4/6/23/26	0/1/1/1

There are no bond length outliers.

There are no bond angle outliers.

All (2) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
4	R	601	NAG	C1
4	S	601	NAG	C1

5 of 8 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	S	601	NAG	C8-C7-N2-C2
4	S	601	NAG	O7-C7-N2-C2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
4	S	601	NAG	O5-C5-C6-O6
4	S	601	NAG	C4-C5-C6-O6
4	R	601	NAG	C4-C5-C6-O6

There are no ring outliers.

2 monomers are involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	R	601	NAG	4	0
4	S	601	NAG	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	213/216 (98%)	-0.19	2 (0%) 81 77	26, 31, 45, 62	0
1	H	214/216 (99%)	0.11	6 (2%) 55 49	29, 39, 56, 80	0
2	B	212/214 (99%)	-0.01	2 (0%) 81 77	24, 32, 53, 77	0
2	L	212/214 (99%)	-0.06	2 (0%) 81 77	24, 33, 46, 81	0
3	R	208/208 (100%)	0.49	15 (7%) 23 18	28, 42, 87, 142	0
3	S	208/208 (100%)	0.73	17 (8%) 19 16	39, 55, 93, 119	0
All	All	1267/1276 (99%)	0.17	44 (3%) 47 41	24, 37, 73, 142	0

The worst 5 of 44 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	R	578	GLY	5.9
3	R	458	VAL	5.3
3	R	381	VAL	5.1
3	S	458	VAL	5.0
3	R	580	ASP	4.8

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	NAG	S	601	14/15	0.39	0.26	127,133,136,138	0
4	NAG	R	601	14/15	0.40	0.24	124,129,132,132	0

6.5 Other polymers [i](#)

There are no such residues in this entry.