



# Full wwPDB X-ray Structure Validation Report ⓘ

Sep 19, 2023 – 12:49 PM EDT

PDB ID : 5E29  
Title : Crystal Structure of Deoxygenated Hemoglobin in Complex with an Allosteric Effector and Nitric Oxide  
Authors : Safo, M.K.; Deshpande, T.M.  
Deposited on : 2015-09-30  
Resolution : 1.85 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtrriage (Phenix) : 1.13  
EDS : 2.35.1  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.35.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*





The reported resolution of this entry is 1.85 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	2469 (1.86-1.86)
Clashscore	141614	2625 (1.86-1.86)
Ramachandran outliers	138981	2592 (1.86-1.86)
Sidechain outliers	138945	2592 (1.86-1.86)
RSRZ outliers	127900	2436 (1.86-1.86)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	141	 90% 10%
1	C	141	 86% 14%
2	B	145	 75% 24% .
2	D	145	 86% 14%

## 2 Entry composition [i](#)

There are 9 unique types of molecules in this entry. The entry contains 5613 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Hemoglobin subunit alpha.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	141	1069	685	187	194	3	0	0	0
1	C	141	1069	685	187	194	3	0	0	0

- Molecule 2 is a protein called Hemoglobin subunit beta.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	145	1116	719	194	200	3	0	0	0
2	D	145	1116	719	194	200	3	0	0	0

- Molecule 3 is NITRIC OXIDE (three-letter code: NO) (formula: NO).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	N	O	0	0
			2	1	1		
3	C	1	Total	N	O	0	0
			2	1	1		

- Molecule 4 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (formula:  $C_{34}H_{32}FeN_4O_4$ ).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
4	A	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		
4	B	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		
4	C	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		
4	D	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		

- Molecule 5 is 2-(nitrooxy)ethyl 2-(4-{2-[(3,5-dimethylphenyl)amino]-2-oxoethyl}phenoxy)-2-methylpropanoate (three-letter code: 5JN) (formula:  $C_{22}H_{26}N_2O_7$ ).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
5	A	1	31	22	2	7	0	1

- Molecule 6 is 2-{4-[(3,5-DIMETHYLANILINO)-CARBONYL-METHYL]-PHENOXY}-2-METHYLPROPIONIC ACID (three-letter code: RQ3) (formula: C<sub>20</sub>H<sub>23</sub>NO<sub>4</sub>).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
6	A	1	25	20	1	4	0	0
6	B	1	25	20	1	4	0	1

- Molecule 7 is NITRATE ION (three-letter code: NO3) (formula: NO<sub>3</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	B	1	Total	N	O	0	0
			4	1	3		
7	D	1	Total	N	O	0	0
			4	1	3		

- Molecule 8 is CACODYLATE ION (three-letter code: CAC) (formula: C<sub>2</sub>H<sub>6</sub>AsO<sub>2</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
8	D	1	Total	As	C	O	0	0
			5	1	2	2		

- Molecule 9 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
9	A	221	Total O 221 221	0	0
9	B	220	Total O 220 220	0	0
9	C	262	Total O 262 262	0	0
9	D	270	Total O 270 270	0	0

### 3 Residue-property plots


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Hemoglobin subunit alpha

Chain A:  90% 10%




- Molecule 1: Hemoglobin subunit alpha

Chain C:  86% 14%





- Molecule 2: Hemoglobin subunit beta

Chain B:  % 75% 24%



- Molecule 2: Hemoglobin subunit beta

Chain D:  % 86% 14%





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	95.88Å 98.42Å 65.44Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	29.54 – 1.85 29.54 – 1.85	Depositor EDS
% Data completeness (in resolution range)	98.1 (29.54-1.85) 98.2 (29.54-1.85)	Depositor EDS
$R_{merge}$	0.08	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.96 (at 1.85Å)	Xtrriage
Refinement program	CNS	Depositor
R, $R_{free}$	0.198 , 0.253 0.194 , 0.250	Depositor DCC
$R_{free}$ test set	2669 reflections (5.07%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	18.5	Xtrriage
Anisotropy	0.150	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.30 , 60.4	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.38$ , $\langle L^2 \rangle = 0.21$	Xtrriage
Estimated twinning fraction	0.045 for k,h,-l	Xtrriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	5613	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	25.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.51% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: RQ3, NO3, HEM, NO, 5JN, CAC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.39	0/1097	0.56	0/1491
1	C	0.42	0/1097	0.59	0/1491
2	B	0.40	0/1146	0.57	0/1556
2	D	0.44	0/1146	0.59	0/1556
All	All	0.41	0/4486	0.58	0/6094

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1069	0	1073	11	0
1	C	1069	0	1073	18	0
2	B	1116	0	1106	36	0
2	D	1116	0	1106	18	0
3	A	2	0	0	0	0
3	C	2	0	0	0	0
4	A	43	0	30	0	0
4	B	43	0	30	3	0
4	C	43	0	30	0	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	D	43	0	30	2	0
5	A	31	0	19	2	0
6	A	25	0	22	0	0
6	B	25	0	17	1	0
7	B	4	0	0	1	0
7	D	4	0	0	0	0
8	D	5	0	0	1	0
9	A	221	0	0	2	0
9	B	220	0	0	11	0
9	C	262	0	0	8	0
9	D	270	0	0	8	0
All	All	5613	0	4536	82	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (82) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:13:ALA:HB3	9:B:301:HOH:O	1.58	1.03
1:C:56:LYS:HE2	9:C:476:HOH:O	1.82	0.79
2:B:143:HIS:HD2	2:B:144:LYS:NZ	1.82	0.76
2:B:43:GLU:HG3	9:B:305:HOH:O	1.88	0.74
2:B:97:HIS:NE2	7:B:203:NO3:O2	2.21	0.74
1:C:7:LYS:O	1:C:11:LYS:HG3	1.90	0.72
1:C:113:LEU:HB3	1:C:116:GLU:HG2	1.73	0.71
2:D:66:LYS:HD3	4:D:201:HEM:HAA2	1.73	0.69
1:C:16:LYS:HD3	9:C:523:HOH:O	1.94	0.67
2:D:65:LYS:HE2	9:D:478:HOH:O	1.96	0.65
2:B:37:TRP:HB2	1:C:92:ARG:HB3	1.80	0.63
1:C:96:VAL:HB	9:C:420:HOH:O	1.98	0.62
2:D:2:HIS:CD2	2:D:3:LEU:H	2.16	0.62
1:C:6:ASP:O	1:C:10:VAL:HG13	2.01	0.61
2:B:10:ALA:HA	9:B:301:HOH:O	2.01	0.61
1:C:90:LYS:HG2	9:C:504:HOH:O	2.01	0.60
2:D:143:HIS:HD2	2:D:144:LYS:NZ	2.03	0.57
1:A:74:ASP:HB2	9:A:401:HOH:O	2.04	0.57
2:B:6:GLU:H	2:B:6:GLU:CD	2.06	0.57
2:B:19:ASN:C	2:B:19:ASN:HD22	2.08	0.57
2:D:2:HIS:HD2	2:D:3:LEU:H	1.52	0.56
2:B:143:HIS:HD2	2:B:144:LYS:HZ2	1.52	0.55

Continued on next page...

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:D:63:HIS:CE1	4:D:201:HEM:HBD2	2.42	0.55
2:B:6:GLU:HG3	9:B:422:HOH:O	2.07	0.55
2:B:80:ASN:ND2	2:B:83:GLY:H	2.05	0.54
2:B:75:LEU:HD21	2:B:133:VAL:HG11	1.88	0.54
2:B:53:ALA:O	2:B:57:ASN:HB2	2.08	0.53
2:D:123:THR:OG1	2:D:126:VAL:HG23	2.08	0.53
1:A:16:LYS:HE3	1:A:116:GLU:OE2	2.08	0.53
2:B:143:HIS:HD2	2:B:144:LYS:HZ1	1.55	0.53
2:D:80:ASN:HB2	9:D:433:HOH:O	2.08	0.53
2:D:3:LEU:HD22	2:D:7:GLU:HB3	1.91	0.53
1:C:42:TYR:C	1:C:44:PRO:HD3	2.28	0.52
2:B:117:HIS:HE1	9:B:342:HOH:O	1.93	0.52
2:B:120:LYS:HG3	9:B:451:HOH:O	2.13	0.49
2:B:143:HIS:CD2	2:B:144:LYS:NZ	2.71	0.49
2:B:143:HIS:CD2	2:B:144:LYS:HZ2	2.29	0.49
1:C:45:HIS:HB2	9:C:501:HOH:O	2.13	0.49
1:C:51:GLY:O	1:C:56:LYS:HE3	2.12	0.49
8:D:202:CAC:AS	9:D:310:HOH:O	2.90	0.49
1:A:131:SER:HA	5:A:203[A]:5JN:O30	2.13	0.48
1:C:88:ALA:HB2	9:C:389:HOH:O	2.12	0.48
2:D:82:LYS:HG2	9:D:424:HOH:O	2.13	0.48
2:B:66:LYS:HD3	4:B:201:HEM:HAA2	1.95	0.48
1:A:16:LYS:HG2	1:A:116:GLU:OE2	2.13	0.48
1:A:116:GLU:OE1	1:A:116:GLU:N	2.44	0.48
1:C:43:PHE:N	1:C:44:PRO:HD3	2.27	0.48
1:A:99:LYS:NZ	9:A:307:HOH:O	2.46	0.48
2:D:2:HIS:CD2	2:D:3:LEU:N	2.81	0.47
2:B:82:LYS:HE3	9:B:337:HOH:O	2.13	0.47
2:D:124:PRO:HB2	2:D:125:PRO:HD3	1.96	0.47
2:B:92:HIS:HA	2:B:96:LEU:HB2	1.97	0.47
1:A:21:ALA:HB1	1:A:63:ALA:HB1	1.97	0.46
2:B:101:GLU:HG2	1:C:94:ASP:OD2	2.15	0.46
1:A:76:MET:N	1:A:77:PRO:CD	2.79	0.46
2:B:43:GLU:HG2	9:B:365:HOH:O	2.15	0.46
1:A:94:ASP:OD2	1:A:96:VAL:HG22	2.17	0.45
1:A:51:GLY:O	1:A:56:LYS:HE3	2.17	0.45
1:C:94:ASP:OD2	1:C:96:VAL:HG13	2.16	0.44
2:B:80:ASN:HD21	2:B:83:GLY:H	1.66	0.43
2:B:104:ARG:HH22	2:B:139:ASN:ND2	2.16	0.43
2:B:15:TRP:CD2	2:B:75:LEU:HD12	2.54	0.42
2:B:77:HIS:CD2	2:B:77:HIS:N	2.86	0.42

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:143:HIS:CD2	2:B:144:LYS:HZ1	2.37	0.42
1:C:68:ASN:HB3	9:C:334:HOH:O	2.19	0.42
5:A:203[A]:5JN:H24	9:B:424:HOH:O	2.20	0.42
9:B:396:HOH:O	1:C:45:HIS:HD2	2.03	0.41
2:D:20:VAL:HG12	9:D:379:HOH:O	2.19	0.41
2:B:5:PRO:HD2	2:B:6:GLU:OE2	2.20	0.41
1:A:92:ARG:HB3	2:D:37:TRP:HB2	2.03	0.41
1:C:76:MET:HE3	9:C:407:HOH:O	2.21	0.41
2:B:124:PRO:HB2	2:B:125:PRO:HD3	2.02	0.41
2:D:21:ASP:HB3	2:D:65:LYS:HE3	2.02	0.41
2:B:37:TRP:CZ3	6:B:202[B]:RQ3:H61	2.55	0.41
2:B:25:GLY:HA3	9:B:378:HOH:O	2.20	0.41
2:D:105:LEU:HA	9:D:303:HOH:O	2.21	0.41
2:B:99:ASP:HA	2:B:100:PRO:HD3	1.94	0.41
2:D:108:ASN:ND2	9:D:303:HOH:O	2.37	0.40
2:B:45:PHE:HA	2:B:59:LYS:HD2	2.03	0.40
2:D:43:GLU:HB2	9:D:418:HOH:O	2.20	0.40
2:B:67:VAL:HG13	4:B:201:HEM:C2B	2.55	0.40
2:B:67:VAL:HG13	4:B:201:HEM:C1B	2.56	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	139/141 (99%)	134 (96%)	5 (4%)	0	100	100
1	C	139/141 (99%)	135 (97%)	4 (3%)	0	100	100
2	B	143/145 (99%)	138 (96%)	5 (4%)	0	100	100
2	D	143/145 (99%)	140 (98%)	3 (2%)	0	100	100
All	All	564/572 (99%)	547 (97%)	17 (3%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	113/113 (100%)	113 (100%)	0	100	100
1	C	113/113 (100%)	113 (100%)	0	100	100
2	B	117/117 (100%)	115 (98%)	2 (2%)	60	47
2	D	117/117 (100%)	117 (100%)	0	100	100
All	All	460/460 (100%)	458 (100%)	2 (0%)	91	89

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	B	2	HIS
2	B	19	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (14) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	9	ASN
1	A	87	HIS
2	B	19	ASN
2	B	63	HIS
2	B	77	HIS
2	B	80	ASN
2	B	117	HIS
2	B	139	ASN
2	B	143	HIS
1	C	20	HIS
2	D	2	HIS
2	D	77	HIS
2	D	80	ASN
2	D	143	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

12 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	HEM	A	202	3	41,50,50	1.43	7 (17%)	45,82,82	0.98	3 (6%)
6	RQ3	B	202[B]	-	26,26,26	2.28	12 (46%)	37,37,37	1.59	3 (8%)
8	CAC	D	202	-	0,4,4	-	-	0,6,6	-	-
4	HEM	D	201	2	41,50,50	1.52	8 (19%)	45,82,82	1.27	5 (11%)
4	HEM	B	201	2	41,50,50	1.21	5 (12%)	45,82,82	1.20	4 (8%)
6	RQ3	A	204	-	26,26,26	2.96	10 (38%)	37,37,37	1.59	6 (16%)
3	NO	C	201	4	0,1,1	-	-	-	-	-
5	5JN	A	203[A]	-	30,32,32	1.99	13 (43%)	39,44,44	1.35	3 (7%)
7	NO3	B	203	-	1,3,3	3.61	1 (100%)	0,3,3	-	-
4	HEM	C	202	3	41,50,50	1.34	7 (17%)	45,82,82	1.15	3 (6%)
3	NO	A	201	4	0,1,1	-	-	-	-	-
7	NO3	D	203	-	1,3,3	3.61	1 (100%)	0,3,3	-	-

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the

Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	HEM	A	202	3	-	4/12/54/54	-
6	RQ3	B	202[B]	-	-	4/19/19/19	0/2/2/2
4	HEM	D	201	2	-	5/12/54/54	-
4	HEM	B	201	2	-	6/12/54/54	-
6	RQ3	A	204	-	-	0/19/19/19	0/2/2/2
5	5JN	A	203[A]	-	-	11/24/26/26	0/2/2/2
4	HEM	C	202	3	-	3/12/54/54	-

All (64) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	A	204	RQ3	C13-C11	8.41	1.57	1.38
6	B	202[B]	RQ3	C10-C9	5.14	1.48	1.38
6	A	204	RQ3	C9-C8	4.98	1.48	1.38
6	A	204	RQ3	C22-C18	4.86	1.47	1.39
6	A	204	RQ3	C14-C15	-4.48	1.42	1.51
4	A	202	HEM	C3C-CAC	-4.41	1.38	1.47
5	A	203[A]	5JN	C15-C14	4.36	1.46	1.38
6	A	204	RQ3	C14-C11	-4.00	1.45	1.51
5	A	203[A]	5JN	C12-C10	-3.98	1.43	1.51
6	B	202[B]	RQ3	C14-C15	-3.71	1.43	1.51
6	A	204	RQ3	C10-C9	3.65	1.45	1.38
4	D	201	HEM	CHB-C1B	3.63	1.44	1.35
7	B	203	NO3	O1-N	3.61	1.40	1.24
7	D	203	NO3	O1-N	3.61	1.40	1.24
6	A	204	RQ3	C21-C20	3.53	1.44	1.39
4	D	201	HEM	C3C-CAC	-3.28	1.41	1.47
4	C	202	HEM	C3C-CAC	-3.17	1.41	1.47
6	B	202[B]	RQ3	C22-C18	3.09	1.44	1.39
6	B	202[B]	RQ3	C19-C20	3.09	1.44	1.39
6	A	204	RQ3	O7-C4	3.03	1.50	1.45
4	A	202	HEM	CBB-CAB	3.00	1.45	1.30
4	B	201	HEM	C3C-CAC	-2.94	1.41	1.47
4	D	201	HEM	C4A-NA	2.92	1.42	1.36
6	A	204	RQ3	C21-C23	2.86	1.43	1.39
4	C	202	HEM	CBB-CAB	2.84	1.44	1.30
6	B	202[B]	RQ3	C9-C8	2.80	1.44	1.38
4	A	202	HEM	C3C-C2C	-2.78	1.36	1.40
5	A	203[A]	5JN	C15-C16	2.75	1.44	1.38

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	A	203[A]	5JN	C8-C3	2.72	1.43	1.39
4	A	202	HEM	CAB-C3B	-2.65	1.40	1.47
6	B	202[B]	RQ3	C14-C11	2.63	1.55	1.51
6	B	202[B]	RQ3	O7-C4	2.62	1.49	1.45
6	A	204	RQ3	C6-C4	2.54	1.57	1.52
4	D	201	HEM	CHA-C4D	2.54	1.41	1.35
4	B	201	HEM	CBB-CAB	2.53	1.42	1.30
4	C	202	HEM	C3C-C2C	-2.52	1.36	1.40
6	B	202[B]	RQ3	C13-C12	2.52	1.43	1.38
4	D	201	HEM	C1B-C2B	2.51	1.49	1.44
6	B	202[B]	RQ3	C22-C23	2.48	1.43	1.39
5	A	203[A]	5JN	C10-N9	-2.48	1.30	1.35
5	A	203[A]	5JN	C4-C5	2.45	1.43	1.39
4	D	201	HEM	CBB-CAB	2.45	1.42	1.30
4	B	201	HEM	CAB-C3B	-2.44	1.40	1.47
4	D	201	HEM	CAB-C3B	-2.39	1.40	1.47
5	A	203[A]	5JN	C4-C3	2.38	1.43	1.39
5	A	203[A]	5JN	C17-C16	2.36	1.43	1.38
4	B	201	HEM	CBC-CAC	2.34	1.44	1.29
4	A	202	HEM	CHB-C1B	2.34	1.41	1.35
5	A	203[A]	5JN	C8-C7	2.31	1.42	1.39
4	C	202	HEM	C2C-C1C	2.28	1.47	1.42
4	D	201	HEM	CBC-CAC	2.26	1.44	1.29
4	C	202	HEM	C1D-C2D	2.25	1.48	1.44
4	A	202	HEM	CBC-CAC	2.24	1.44	1.29
5	A	203[A]	5JN	C6-C5	2.24	1.42	1.39
6	B	202[B]	RQ3	C12-C8	2.22	1.43	1.38
6	B	202[B]	RQ3	C19-C18	2.22	1.43	1.39
4	C	202	HEM	CAB-C3B	-2.18	1.41	1.47
5	A	203[A]	5JN	C17-C18	2.16	1.42	1.38
4	A	202	HEM	CMA-C3A	2.14	1.56	1.51
4	B	201	HEM	CHA-C4D	2.13	1.40	1.35
5	A	203[A]	5JN	C6-C7	2.12	1.42	1.39
6	B	202[B]	RQ3	C21-C23	2.03	1.42	1.39
4	C	202	HEM	CBC-CAC	2.01	1.42	1.29
5	A	203[A]	5JN	C18-C13	2.01	1.43	1.38

All (27) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	A	204	RQ3	C4-O7-C8	6.16	130.84	120.95
6	B	202[B]	RQ3	C4-O7-C8	5.62	129.97	120.95

*Continued on next page...*

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	B	202[B]	RQ3	O7-C4-C1	5.33	117.26	109.99
5	A	203[A]	5JN	C20-O19-C16	4.94	128.87	120.95
4	C	202	HEM	C4B-CHC-C1C	3.81	127.59	122.56
4	B	201	HEM	CBA-CAA-C2A	-3.35	106.90	112.62
5	A	203[A]	5JN	O25-C23-C20	3.29	117.11	111.64
4	D	201	HEM	CMB-C2B-C1B	3.20	129.92	125.04
4	A	202	HEM	C4C-CHD-C1D	2.77	126.21	122.56
5	A	203[A]	5JN	O19-C20-C23	2.75	116.55	110.67
4	B	201	HEM	C4B-CHC-C1C	2.66	126.07	122.56
4	C	202	HEM	C4C-CHD-C1D	2.60	125.99	122.56
4	D	201	HEM	CMA-C3A-C4A	-2.54	124.55	128.46
4	B	201	HEM	C4A-C3A-C2A	-2.51	105.25	107.00
6	A	204	RQ3	C14-C11-C13	-2.41	117.44	120.89
4	D	201	HEM	CMA-C3A-C2A	2.39	129.45	124.94
4	B	201	HEM	C4C-CHD-C1D	2.38	125.70	122.56
4	A	202	HEM	C4B-CHC-C1C	2.36	125.68	122.56
6	A	204	RQ3	O2-C1-C4	2.31	120.96	113.99
6	A	204	RQ3	C21-C20-C19	2.23	120.75	118.09
6	A	204	RQ3	C18-N17-C15	2.21	131.37	127.50
4	C	202	HEM	C4A-C3A-C2A	-2.17	105.49	107.00
6	A	204	RQ3	C9-C10-C11	-2.11	118.12	121.03
4	D	201	HEM	CAA-C2A-C3A	2.09	133.25	127.25
4	D	201	HEM	C2C-C3C-C4C	-2.06	105.46	106.90
6	B	202[B]	RQ3	C21-C20-C19	2.05	120.53	118.09
4	A	202	HEM	CMC-C2C-C3C	2.03	128.48	124.68

There are no chirality outliers.

All (33) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	B	201	HEM	C2B-C3B-CAB-CBB
5	A	203[A]	5JN	C21-C20-C23-O24
5	A	203[A]	5JN	C21-C20-C23-O25
5	A	203[A]	5JN	O19-C20-C23-O24
5	A	203[A]	5JN	O19-C20-C23-O25
5	A	203[A]	5JN	C20-C23-O25-C26
6	B	202[B]	RQ3	O3-C1-C4-O7
5	A	203[A]	5JN	O24-C23-O25-C26
4	B	201	HEM	C4B-C3B-CAB-CBB
6	B	202[B]	RQ3	O3-C1-C4-C6
4	D	201	HEM	C2B-C3B-CAB-CBB
4	D	201	HEM	C2A-CAA-CBA-CGA

Continued on next page...

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms
6	B	202[B]	RQ3	O2-C1-C4-C6
6	B	202[B]	RQ3	O2-C1-C4-O7
4	D	201	HEM	C4B-C3B-CAB-CBB
5	A	203[A]	5JN	C26-C27-O28-N29
4	C	202	HEM	CAD-CBD-CGD-O2D
4	C	202	HEM	CAD-CBD-CGD-O1D
4	C	202	HEM	C2D-C3D-CAD-CBD
5	A	203[A]	5JN	C4-C3-N9-C10
4	D	201	HEM	CAD-CBD-CGD-O1D
4	D	201	HEM	CAD-CBD-CGD-O2D
5	A	203[A]	5JN	C22-C20-C23-O25
4	A	202	HEM	CAD-CBD-CGD-O2D
4	B	201	HEM	CAA-CBA-CGA-O2A
4	A	202	HEM	CAA-CBA-CGA-O2A
5	A	203[A]	5JN	C22-C20-C23-O24
4	B	201	HEM	C2D-C3D-CAD-CBD
4	A	202	HEM	CAD-CBD-CGD-O1D
4	B	201	HEM	CAA-CBA-CGA-O1A
4	A	202	HEM	CAA-CBA-CGA-O1A
5	A	203[A]	5JN	C8-C3-N9-C10
4	B	201	HEM	CAD-CBD-CGD-O2D


There are no ring outliers.


6 monomers are involved in 10 short contacts:



Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	B	202[B]	RQ3	1	0
8	D	202	CAC	1	0
4	D	201	HEM	2	0
4	B	201	HEM	3	0
5	A	203[A]	5JN	2	0
7	B	203	NO3	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and

any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.









## 5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.



## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	141/141 (100%)	-0.52	0 <a href="#">100</a> <a href="#">100</a>	13, 21, 32, 40	0
1	C	141/141 (100%)	-0.64	0 <a href="#">100</a> <a href="#">100</a>	12, 19, 28, 39	0
2	B	145/145 (100%)	-0.33	1 (0%) <a href="#">87</a> <a href="#">88</a>	15, 25, 41, 60	0
2	D	145/145 (100%)	-0.50	1 (0%) <a href="#">87</a> <a href="#">88</a>	12, 19, 34, 60	0
All	All	572/572 (100%)	-0.50	2 (0%) <a href="#">94</a> <a href="#">93</a>	12, 21, 35, 60	0

All (2) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	D	2	HIS	3.9
2	B	2	HIS	2.2

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)


There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
5	5JN	A	203[A]	31/31	0.75	0.27	31,34,41,48	31
6	RQ3	B	202[B]	25/25	0.75	0.27	29,33,40,47	25
6	RQ3	A	204	25/25	0.80	0.24	31,40,47,50	25
8	CAC	D	202	5/5	0.86	0.14	93,93,95,100	0
7	NO3	B	203	4/4	0.89	0.19	45,46,53,59	0
7	NO3	D	203	4/4	0.90	0.15	25,41,49,49	0
4	HEM	B	201	43/43	0.97	0.08	15,25,42,44	0
4	HEM	A	202	43/43	0.98	0.07	7,18,29,39	0
4	HEM	C	202	43/43	0.98	0.07	10,16,37,44	0
4	HEM	D	201	43/43	0.98	0.08	10,15,22,25	0
3	NO	A	201	2/2	0.99	0.09	17,17,17,19	0
3	NO	C	201	2/2	1.00	0.04	17,17,17,19	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.






**Electron density around RQ3 B 202 (B):**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)




**Electron density around RQ3 A 204:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around HEM B 201:**


$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)




**Electron density around HEM A 202:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)







## 6.5 Other polymers [i](#)

There are no such residues in this entry.