



# wwPDB X-ray Structure Validation Summary Report i

Sep 20, 2023 – 09:07 PM EDT

PDB ID : 5J3D  
Title : Crystal structure of human Fab 14N4 in complex with post-fusion RSV F  
Authors : Mousa, J.J.; Crowe, J.E.  
Deposited on : 2016-03-30  
Resolution : 4.08 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>  
with specific help available everywhere you see the i symbol.

The types of validation reports are described at  
<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references](#) ①) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.13
EDS	:	2.35.1
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.35.1


# 1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*


The reported resolution of this entry is 4.08 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1133 (4.46-3.70)
Clashscore	141614	1013 (4.42-3.74)
Ramachandran outliers	138981	1151 (4.46-3.70)
Sidechain outliers	138945	1139 (4.46-3.70)
RSRZ outliers	127900	1012 (4.48-3.68)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.



*Continued on next page...*

Continued from previous page...



## 2 Entry composition [\(i\)](#)

There are 4 unique types of molecules in this entry. The entry contains 19902 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 14N4 heavy chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	220	Total	C	N	O	S	0	0	0
			1624	1020	275	322	7			
1	C	220	Total	C	N	O	S	0	0	0
			1624	1020	275	322	7			
1	H	220	Total	C	N	O	S	0	0	0
			1624	1020	275	322	7			

- Molecule 2 is a protein called 14N4 light chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	215	Total	C	N	O	S	0	0	0
			1665	1044	276	339	6			
2	D	215	Total	C	N	O	S	0	0	0
			1665	1044	276	339	6			
2	L	215	Total	C	N	O	S	0	0	0
			1665	1044	276	339	6			

- Molecule 3 is a protein called Fusion glycoprotein F0.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	E	73	Total	C	N	O	S	0	0	0
			580	366	95	116	3			
3	G	73	Total	C	N	O	S	0	0	0
			580	366	95	116	3			
3	J	73	Total	C	N	O	S	0	0	0
			580	366	95	116	3			

- Molecule 4 is a protein called Fusion glycoprotein F0.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	F	358	Total	C	N	O	S	0	0	0
			2765	1744	458	545	18			

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	I	358	Total	C	N	O	S	0	0	0
			2765	1744	458	545	18			
4	K	358	Total	C	N	O	S	0	0	0
			2765	1744	458	545	18			

There are 87 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
F	379	VAL	ILE	conflict	UNP P03420
F	447	VAL	MET	conflict	UNP P03420
F	514	GLY	-	expression tag	UNP P03420
F	515	LEU	-	expression tag	UNP P03420
F	516	GLU	-	expression tag	UNP P03420
F	517	VAL	-	expression tag	UNP P03420
F	518	LEU	-	expression tag	UNP P03420
F	519	PHE	-	expression tag	UNP P03420
F	520	GLN	-	expression tag	UNP P03420
F	521	GLY	-	expression tag	UNP P03420
F	522	PRO	-	expression tag	UNP P03420
F	523	HIS	-	expression tag	UNP P03420
F	524	HIS	-	expression tag	UNP P03420
F	525	HIS	-	expression tag	UNP P03420
F	526	HIS	-	expression tag	UNP P03420
F	527	HIS	-	expression tag	UNP P03420
F	528	HIS	-	expression tag	UNP P03420
F	529	HIS	-	expression tag	UNP P03420
F	530	HIS	-	expression tag	UNP P03420
F	531	SER	-	expression tag	UNP P03420
F	532	ALA	-	expression tag	UNP P03420
F	533	TRP	-	expression tag	UNP P03420
F	534	SER	-	expression tag	UNP P03420
F	535	HIS	-	expression tag	UNP P03420
F	536	PRO	-	expression tag	UNP P03420
F	537	GLN	-	expression tag	UNP P03420
F	538	PHE	-	expression tag	UNP P03420
F	539	GLU	-	expression tag	UNP P03420
F	540	LYS	-	expression tag	UNP P03420
I	379	VAL	ILE	conflict	UNP P03420
I	447	VAL	MET	conflict	UNP P03420
I	514	GLY	-	expression tag	UNP P03420
I	515	LEU	-	expression tag	UNP P03420
I	516	GLU	-	expression tag	UNP P03420

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
I	517	VAL	-	expression tag	UNP P03420
I	518	LEU	-	expression tag	UNP P03420
I	519	PHE	-	expression tag	UNP P03420
I	520	GLN	-	expression tag	UNP P03420
I	521	GLY	-	expression tag	UNP P03420
I	522	PRO	-	expression tag	UNP P03420
I	523	HIS	-	expression tag	UNP P03420
I	524	HIS	-	expression tag	UNP P03420
I	525	HIS	-	expression tag	UNP P03420
I	526	HIS	-	expression tag	UNP P03420
I	527	HIS	-	expression tag	UNP P03420
I	528	HIS	-	expression tag	UNP P03420
I	529	HIS	-	expression tag	UNP P03420
I	530	HIS	-	expression tag	UNP P03420
I	531	SER	-	expression tag	UNP P03420
I	532	ALA	-	expression tag	UNP P03420
I	533	TRP	-	expression tag	UNP P03420
I	534	SER	-	expression tag	UNP P03420
I	535	HIS	-	expression tag	UNP P03420
I	536	PRO	-	expression tag	UNP P03420
I	537	GLN	-	expression tag	UNP P03420
I	538	PHE	-	expression tag	UNP P03420
I	539	GLU	-	expression tag	UNP P03420
I	540	LYS	-	expression tag	UNP P03420
K	379	VAL	ILE	conflict	UNP P03420
K	447	VAL	MET	conflict	UNP P03420
K	514	GLY	-	expression tag	UNP P03420
K	515	LEU	-	expression tag	UNP P03420
K	516	GLU	-	expression tag	UNP P03420
K	517	VAL	-	expression tag	UNP P03420
K	518	LEU	-	expression tag	UNP P03420
K	519	PHE	-	expression tag	UNP P03420
K	520	GLN	-	expression tag	UNP P03420
K	521	GLY	-	expression tag	UNP P03420
K	522	PRO	-	expression tag	UNP P03420
K	523	HIS	-	expression tag	UNP P03420
K	524	HIS	-	expression tag	UNP P03420
K	525	HIS	-	expression tag	UNP P03420
K	526	HIS	-	expression tag	UNP P03420
K	527	HIS	-	expression tag	UNP P03420
K	528	HIS	-	expression tag	UNP P03420
K	529	HIS	-	expression tag	UNP P03420

*Continued on next page...*


*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
K	530	HIS	-	expression tag	UNP P03420
K	531	SER	-	expression tag	UNP P03420
K	532	ALA	-	expression tag	UNP P03420
K	533	TRP	-	expression tag	UNP P03420
K	534	SER	-	expression tag	UNP P03420
K	535	HIS	-	expression tag	UNP P03420
K	536	PRO	-	expression tag	UNP P03420
K	537	GLN	-	expression tag	UNP P03420
K	538	PHE	-	expression tag	UNP P03420
K	539	GLU	-	expression tag	UNP P03420
K	540	LYS	-	expression tag	UNP P03420


### 3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: 14N4 heavy chain




- Molecule 1: 14N4 heavy chain



- Molecule 1: 14N4 heavy chain




- Molecule 2: 14N4 light chain




- Molecule 2: 14N4 light chain



- Molecule 2: 14N4 light chain



- Molecule 3: Fusion glycoprotein F0




- Molecule 3: Fusion glycoprotein F0





- Molecule 3: Fusion glycoprotein F0



- Molecule 4: Fusion glycoprotein F0




- Molecule 4: Fusion glycoprotein F0



- Molecule 4: Fusion glycoprotein F0





## 4 Data and refinement statistics i

Property	Value	Source
Space group	P 42 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	235.13Å    235.13Å    220.14Å 90.00°    90.00°    90.00°	Depositor
Resolution (Å)	49.50 – 4.08 49.50 – 4.08	Depositor EDS
% Data completeness (in resolution range)	97.3 (49.50-4.08) 84.8 (49.50-4.08)	Depositor EDS
$R_{merge}$	0.30	Depositor
$R_{sym}$	(Not available)	Depositor
$< I/\sigma(I) >$ <sup>1</sup>	1.06 (at 4.14Å)	Xtriage
Refinement program	PHENIX (1.10.1_2155: ???)	Depositor
$R$ , $R_{free}$	0.257 , 0.282 0.257 , 0.282	Depositor DCC
$R_{free}$ test set	1999 reflections (4.13%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	104.5	Xtriage
Anisotropy	0.118	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.32 , 105.9	EDS
L-test for twinning <sup>2</sup>	$<  L  > = 0.45$ , $< L^2 > = 0.28$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.85	EDS
Total number of atoms	19902	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	153.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.33% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $< |L| >$ ,  $< L^2 >$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality i

### 5.1 Standard geometry i

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.34	0/1660	0.56	0/2262
1	C	0.45	0/1660	0.63	0/2262
1	H	0.27	0/1660	0.51	0/2262
2	B	0.34	0/1704	0.55	0/2317
2	D	0.42	0/1704	0.59	0/2317
2	L	0.29	0/1704	0.50	0/2317
3	E	0.50	0/586	0.75	0/789
3	G	0.64	0/586	0.85	1/789 (0.1%)
3	J	0.50	0/586	0.73	0/789
4	F	0.50	0/2805	0.74	2/3803 (0.1%)
4	I	0.54	0/2805	0.81	4/3803 (0.1%)
4	K	0.56	1/2805 (0.0%)	0.80	5/3803 (0.1%)
All	All	0.46	1/20265 (0.0%)	0.68	12/27513 (0.0%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	K	439	CYS	CB-SG	-6.17	1.71	1.82

The worst 5 of 12 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	I	212	CYS	CA-CB-SG	12.46	136.43	114.00
4	K	172	LEU	CA-CB-CG	-6.41	100.56	115.30
4	I	231	LEU	CA-CB-CG	-6.26	100.91	115.30
4	I	373	LEU	CA-CB-CG	-6.26	100.90	115.30
4	I	305	LEU	CB-CG-CD1	-5.41	101.80	111.00

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts [\(i\)](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1624	0	1599	43	0
1	C	1624	0	1599	45	1
1	H	1624	0	1599	36	0
2	B	1665	0	1600	48	0
2	D	1665	0	1600	65	0
2	L	1665	0	1600	33	0
3	E	580	0	589	25	0
3	G	580	0	589	13	0
3	J	580	0	589	26	0
4	F	2765	0	2798	112	0
4	I	2765	0	2798	122	0
4	K	2765	0	2798	109	0
All	All	19902	0	19758	514	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 13.

The worst 5 of 514 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:D:38:TRP:CE2	4:I:268:ASN:HB3	1.81	1.15
2:D:38:TRP:HH2	4:I:269:ASP:HA	1.35	0.89
1:A:35:SER:O	4:I:465:LYS:NZ	2.09	0.86
2:D:109:THR:HG23	4:I:267:THR:HA	1.55	0.85
4:F:345:ASN:HD22	4:I:455:THR:HG21	1.42	0.84

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:14:GLN:OE1	1:C:20:ARG:NH1[8_554]	2.13	0.07

## 5.3 Torsion angles [\(i\)](#)

### 5.3.1 Protein backbone [\(i\)](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	218/224 (97%)	198 (91%)	20 (9%)	0	100 100
1	C	218/224 (97%)	199 (91%)	19 (9%)	0	100 100
1	H	218/224 (97%)	199 (91%)	19 (9%)	0	100 100
2	B	213/218 (98%)	198 (93%)	15 (7%)	0	100 100
2	D	213/218 (98%)	198 (93%)	13 (6%)	2 (1%)	17 55
2	L	213/218 (98%)	198 (93%)	15 (7%)	0	100 100
3	E	71/73 (97%)	65 (92%)	5 (7%)	1 (1%)	11 46
3	G	71/73 (97%)	65 (92%)	5 (7%)	1 (1%)	11 46
3	J	71/73 (97%)	65 (92%)	5 (7%)	1 (1%)	11 46
4	F	354/394 (90%)	333 (94%)	21 (6%)	0	100 100
4	I	354/394 (90%)	334 (94%)	20 (6%)	0	100 100
4	K	354/394 (90%)	332 (94%)	21 (6%)	1 (0%)	41 75
All	All	2568/2727 (94%)	2384 (93%)	178 (7%)	6 (0%)	47 80

5 of 6 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
3	G	69	CYS
3	J	69	CYS
3	E	69	CYS
4	K	153	ALA
2	D	233	GLU

### 5.3.2 Protein sidechains [\(i\)](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	183/187 (98%)	178 (97%)	5 (3%)	44 66
1	C	183/187 (98%)	174 (95%)	9 (5%)	25 52
1	H	183/187 (98%)	177 (97%)	6 (3%)	38 62
2	B	190/193 (98%)	187 (98%)	3 (2%)	62 79
2	D	190/193 (98%)	183 (96%)	7 (4%)	34 60
2	L	190/193 (98%)	187 (98%)	3 (2%)	62 79
3	E	66/66 (100%)	63 (96%)	3 (4%)	27 54
3	G	66/66 (100%)	63 (96%)	3 (4%)	27 54
3	J	66/66 (100%)	63 (96%)	3 (4%)	27 54
4	F	329/362 (91%)	309 (94%)	20 (6%)	18 47
4	I	329/362 (91%)	306 (93%)	23 (7%)	15 42
4	K	329/362 (91%)	310 (94%)	19 (6%)	20 48
All	All	2304/2424 (95%)	2200 (96%)	104 (4%)	27 54

5 of 104 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
4	I	204	LEU
4	I	416	CYS
4	K	477	PHE
4	I	244	THR
4	I	345	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 12 such sidechains are listed below:

Mol	Chain	Res	Type
3	G	81	GLN
4	I	225	GLN
2	L	6	GLN
3	J	26	GLN
3	E	81	GLN

### 5.3.3 RNA [\(i\)](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [\(i\)](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [\(i\)](#)

There are no ligands in this entry.

## 5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [\(i\)](#)

### 6.1 Protein, DNA and RNA chains [\(i\)](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	220/224 (98%)	0.32	15 (6%) 17 14	125, 169, 259, 296	0
1	C	220/224 (98%)	-0.04	5 (2%) 60 51	71, 108, 207, 239	0
1	H	220/224 (98%)	1.56	63 (28%) 0 0	162, 259, 353, 380	0
2	B	215/218 (98%)	0.52	11 (5%) 28 24	130, 208, 264, 282	0
2	D	215/218 (98%)	0.04	5 (2%) 60 51	86, 143, 209, 223	0
2	L	215/218 (98%)	1.80	85 (39%) 0 0	167, 277, 375, 387	0
3	E	73/73 (100%)	-0.18	0 100 100	82, 108, 139, 163	0
3	G	73/73 (100%)	-0.22	0 100 100	79, 102, 128, 162	0
3	J	73/73 (100%)	-0.29	0 100 100	86, 110, 143, 169	0
4	F	358/394 (90%)	-0.06	5 (1%) 75 66	74, 109, 145, 185	0
4	I	358/394 (90%)	-0.13	4 (1%) 80 72	72, 97, 137, 195	0
4	K	358/394 (90%)	-0.17	0 100 100	76, 99, 133, 170	0
All	All	2598/2727 (95%)	0.28	193 (7%) 14 12	71, 125, 322, 387	0

The worst 5 of 193 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
4	F	147	ALA	7.2
4	I	149	ALA	6.5
1	H	228	PRO	6.5
2	L	216	VAL	5.7
2	L	142	ASP	5.7

### 6.2 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [\(i\)](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [\(i\)](#)

There are no ligands in this entry.

### 6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.