



# wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 23, 2022 – 06:10 PM EDT

PDB ID : 5SCJ  
Title : Structure of liver pyruvate kinase in complex with anthraquinone derivative  
106  
Authors : Lulla, A.; Foller, A.; Nain-Perez, A.; Grotli, M.; Brear, P.; Hyvonen, M.  
Deposited on : 2021-12-01  
Resolution : 2.35 Å (reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.27  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.27


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.35 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.






Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1164 (2.36-2.36)
Clashscore	141614	1232 (2.36-2.36)
Ramachandran outliers	138981	1211 (2.36-2.36)
Sidechain outliers	138945	1212 (2.36-2.36)
RSRZ outliers	127900	1150 (2.36-2.36)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	447	 13% 82% 12% • 6%
1	B	447	 6% 85% 12% ••
1	C	447	 2% 83% 11% • 5%
1	D	447	 % 81% 13% 5%
1	E	447	 6% 81% 12% • 6%

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
1	F	447	
1	G	447	
1	H	447	

## 2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 27221 atoms, of which 87 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Pyruvate kinase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	422	Total 3236	C 2034	N 585	O 597	S 20	0	6	0
1	B	436	Total 3329	C 2090	N 604	O 615	S 20	0	4	0
1	C	425	Total 3247	C 2040	N 585	O 603	S 19	0	4	0
1	D	425	Total 3252	C 2042	N 590	O 601	S 19	0	6	0
1	E	419	Total 3210	C 2018	N 579	O 593	S 20	0	5	0
1	F	432	Total 3321	C 2090	N 597	O 614	S 20	0	7	0
1	G	421	Total 3231	C 2031	N 581	O 600	S 19	0	6	0
1	H	425	Total 3251	C 2040	N 594	O 598	S 19	0	4	0

There are 48 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-1	GLY	-	expression tag	UNP Q16716
A	0	SER	-	expression tag	UNP Q16716
A	12	ASP	SER	conflict	UNP Q16716
A	130	GLY	-	linker	UNP Q16716
A	131	SER	-	linker	UNP Q16716
A	132	GLY	-	linker	UNP Q16716
B	-1	GLY	-	expression tag	UNP Q16716
B	0	SER	-	expression tag	UNP Q16716
B	12	ASP	SER	conflict	UNP Q16716
B	130	GLY	-	linker	UNP Q16716
B	131	SER	-	linker	UNP Q16716
B	132	GLY	-	linker	UNP Q16716
C	-1	GLY	-	expression tag	UNP Q16716

*Continued on next page...*

*Continued from previous page...*


Chain	Residue	Modelled	Actual	Comment	Reference
C	0	SER	-	expression tag	UNP Q16716
C	12	ASP	SER	conflict	UNP Q16716
C	130	GLY	-	linker	UNP Q16716
C	131	SER	-	linker	UNP Q16716
C	132	GLY	-	linker	UNP Q16716
D	-1	GLY	-	expression tag	UNP Q16716
D	0	SER	-	expression tag	UNP Q16716
D	12	ASP	SER	conflict	UNP Q16716
D	130	GLY	-	linker	UNP Q16716
D	131	SER	-	linker	UNP Q16716
D	132	GLY	-	linker	UNP Q16716
E	-1	GLY	-	expression tag	UNP Q16716
E	0	SER	-	expression tag	UNP Q16716
E	12	ASP	SER	conflict	UNP Q16716
E	228	GLY	-	linker	UNP Q16716
E	229	SER	-	linker	UNP Q16716
E	230	GLY	-	linker	UNP Q16716
F	-1	GLY	-	expression tag	UNP Q16716
F	0	SER	-	expression tag	UNP Q16716
F	12	ASP	SER	conflict	UNP Q16716
F	228	GLY	-	linker	UNP Q16716
F	229	SER	-	linker	UNP Q16716
F	230	GLY	-	linker	UNP Q16716
G	-1	GLY	-	expression tag	UNP Q16716
G	0	SER	-	expression tag	UNP Q16716
G	12	ASP	SER	conflict	UNP Q16716
G	130	GLY	-	linker	UNP Q16716
G	131	SER	-	linker	UNP Q16716
G	132	GLY	-	linker	UNP Q16716
H	-1	GLY	-	expression tag	UNP Q16716
H	0	SER	-	expression tag	UNP Q16716
H	12	ASP	SER	conflict	UNP Q16716
H	130	GLY	-	linker	UNP Q16716
H	131	SER	-	linker	UNP Q16716
H	132	GLY	-	linker	UNP Q16716

- Molecule 2 is 1,6-di-O-phosphono-beta-D-fructofuranose (three-letter code: FBP) (formula:  $C_6H_{14}O_{12}P_2$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	O	P		
2	A	1	20	6	12	2	0	0
2	B	1	20	6	12	2	0	0
2	C	1	20	6	12	2	0	0
2	D	1	20	6	12	2	0	0
2	E	1	20	6	12	2	0	0
2	F	1	20	6	12	2	0	0
2	G	1	20	6	12	2	0	0
2	H	1	20	6	12	2	0	0

- Molecule 3 is OXALATE ION (three-letter code: OXL) (formula:  $C_2O_4$ ).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C O 6 2 4	0	0
3	B	1	Total C O 6 2 4	0	0
3	C	1	Total C O 6 2 4	0	0
3	D	1	Total C O 6 2 4	0	0
3	E	1	Total C O 6 2 4	0	0
3	F	1	Total C O 6 2 4	0	0
3	G	1	Total C O 6 2 4	0	0
3	H	1	Total C O 6 2 4	0	0

- Molecule 4 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total Mg 1 1	0	0
4	B	1	Total Mg 1 1	0	0
4	C	1	Total Mg 1 1	0	0
4	D	1	Total Mg 1 1	0	0

Continued on next page...

*Continued from previous page...*


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	E	1	Total Mg 1 1	0	0
4	F	1	Total Mg 1 1	0	0
4	G	1	Total Mg 1 1	0	0
4	H	1	Total Mg 1 1	0	0

- Molecule 5 is POTASSIUM ION (three-letter code: K) (formula: K).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total K 1 1	0	0
5	B	1	Total K 1 1	0	0
5	C	1	Total K 1 1	0	0
5	D	1	Total K 1 1	0	0
5	E	1	Total K 1 1	0	0
5	F	1	Total K 1 1	0	0
5	G	1	Total K 1 1	0	0
5	H	1	Total K 1 1	0	0

- Molecule 6 is (2R)-2-hydroxy-2-{2-[4-(3-hydroxy-9,10-dioxo-9,10-dihydroanthracene-2-sulfonyl)piperazin-1-yl]-2-oxoethyl}butanedioic acid (three-letter code: I9K) (formula: C<sub>24</sub>H<sub>22</sub>N<sub>2</sub>O<sub>11</sub>S) (labeled as "Ligand of Interest" by depositor).





Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	
			Total	C	H	N	O			S
6	A	1	Total	C	H	N	O	S	22	0
			60	24	22	2	11	1		
6	C	1	Total	C	H	N	O	S	21	0
			59	24	21	2	11	1		
6	F	1	Total	C	H	N	O	S	22	0
			60	24	22	2	11	1		
6	G	1	Total	C	H	N	O	S	22	0
			60	24	22	2	11	1		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	30	Total	O	0	0
			30	30		
7	B	44	Total	O	0	0
			44	44		
7	C	92	Total	O	0	0
			92	92		
7	D	113	Total	O	0	0
			113	113		
7	E	47	Total	O	0	0
			47	47		
7	F	94	Total	O	0	0
			94	94		
7	G	109	Total	O	0	0
			109	109		
7	H	152	Total	O	0	0
			152	152		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Pyruvate kinase




- Molecule 1: Pyruvate kinase





- Molecule 1: Pyruvate kinase









- Molecule 1: Pyruvate kinase




- Molecule 1: Pyruvate kinase





- Molecule 1: Pyruvate kinase







● Molecule 1: Pyruvate kinase



● Molecule 1: Pyruvate kinase



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	208.14Å 113.06Å 188.96Å 90.00° 91.54° 90.00°	Depositor
Resolution (Å)	188.89 – 2.35 188.89 – 2.35	Depositor EDS
% Data completeness (in resolution range)	56.7 (188.89-2.35) 56.7 (188.89-2.35)	Depositor EDS
$R_{merge}$	0.15	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.61 (at 2.34Å)	Xtrriage
Refinement program	BUSTER 2.10.4 (16-JUL-2021)	Depositor
R, $R_{free}$	0.218 , 0.253 0.206 , 0.231	Depositor DCC
$R_{free}$ test set	4891 reflections (4.76%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	47.7	Xtrriage
Anisotropy	0.032	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.32 , 40.4	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.47$ , $\langle L^2 \rangle = 0.30$	Xtrriage
Estimated twinning fraction	0.010 for -h,-k,l	Xtrriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	27221	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	58.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 56.13 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 2.8637e-05. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, I9K, FBP, K, OXL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.37	0/3307	0.54	0/4469
1	B	0.41	0/3396	0.55	0/4592
1	C	0.44	0/3313	0.58	0/4479
1	D	0.47	0/3326	0.58	0/4497
1	E	0.38	0/3278	0.54	0/4430
1	F	0.44	0/3396	0.56	0/4591
1	G	0.45	0/3303	0.57	0/4465
1	H	0.46	0/3316	0.58	0/4483
All	All	0.43	0/26635	0.56	0/36006

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3236	0	3299	31	0
1	B	3329	0	3394	29	1
1	C	3247	0	3299	33	0
1	D	3252	0	3311	34	0
1	E	3210	0	3270	31	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	F	3321	0	3393	28	0
1	G	3231	0	3284	44	0
1	H	3251	0	3306	35	0
2	A	20	0	10	2	0
2	B	20	0	10	0	0
2	C	20	0	10	0	0
2	D	20	0	10	0	0
2	E	20	0	10	0	0
2	F	20	0	10	0	0
2	G	20	0	10	0	0
2	H	20	0	10	0	0
3	A	6	0	0	0	0
3	B	6	0	0	0	0
3	C	6	0	0	0	0
3	D	6	0	0	0	0
3	E	6	0	0	0	0
3	F	6	0	0	0	0
3	G	6	0	0	0	0
3	H	6	0	0	0	0
4	A	1	0	0	0	0
4	B	1	0	0	0	0
4	C	1	0	0	0	0
4	D	1	0	0	0	0
4	E	1	0	0	0	0
4	F	1	0	0	0	0
4	G	1	0	0	0	0
4	H	1	0	0	0	0
5	A	1	0	0	0	0
5	B	1	0	0	0	0
5	C	1	0	0	0	0
5	D	1	0	0	0	0
5	E	1	0	0	0	0
5	F	1	0	0	0	0
5	G	1	0	0	0	0
5	H	1	0	0	0	0
6	A	38	22	0	0	0
6	C	38	21	0	1	0
6	F	38	22	0	0	0
6	G	38	22	0	2	0
7	A	30	0	0	0	0
7	B	44	0	0	0	1
7	C	92	0	0	0	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
7	D	113	0	0	1	0
7	E	47	0	0	1	0
7	F	94	0	0	1	0
7	G	109	0	0	1	0
7	H	152	0	0	2	0
All	All	27134	87	26636	235	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 235 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:422[A]:GLU:HG3	1:C:452:LEU:HD13	1.56	0.85
1:G:422[A]:GLU:HG3	1:G:452:LEU:HD13	1.60	0.84
1:G:538:ARG:HG2	1:H:536:ILE:HG12	1.66	0.77
1:A:536:ILE:HG12	1:B:538:ARG:HG2	1.70	0.72
1:E:408:GLU:OE2	1:F:411:ARG:NH2	2.23	0.71

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
7:B:744:HOH:O	7:B:744:HOH:O[2_555]	1.86	0.34
1:B:511:LEU:O	1:B:511:LEU:O[2_555]	2.09	0.11

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	424/447 (95%)	418 (99%)	5 (1%)	1 (0%)	47 56

Continued on next page...



Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	438/447 (98%)	430 (98%)	4 (1%)	4 (1%)	17	17
1	C	425/447 (95%)	419 (99%)	3 (1%)	3 (1%)	22	23
1	D	429/447 (96%)	420 (98%)	6 (1%)	3 (1%)	22	23
1	E	420/447 (94%)	416 (99%)	3 (1%)	1 (0%)	47	56
1	F	435/447 (97%)	429 (99%)	5 (1%)	1 (0%)	47	56
1	G	423/447 (95%)	415 (98%)	5 (1%)	3 (1%)	22	23
1	H	427/447 (96%)	420 (98%)	4 (1%)	3 (1%)	22	23
All	All	3421/3576 (96%)	3367 (98%)	35 (1%)	19 (1%)	25	27

5 of 19 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	271	GLY
1	B	129	PRO
1	B	231	PRO
1	C	231	PRO
1	G	271	GLY

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	340/352 (97%)	329 (97%)	11 (3%)	39	47
1	B	349/352 (99%)	336 (96%)	13 (4%)	34	42
1	C	341/352 (97%)	332 (97%)	9 (3%)	46	56
1	D	342/352 (97%)	333 (97%)	9 (3%)	46	56
1	E	338/352 (96%)	327 (97%)	11 (3%)	38	46
1	F	350/352 (99%)	337 (96%)	13 (4%)	34	42
1	G	340/352 (97%)	332 (98%)	8 (2%)	49	59
1	H	340/352 (97%)	333 (98%)	7 (2%)	53	65
All	All	2740/2816 (97%)	2659 (97%)	81 (3%)	44	50

5 of 81 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	F	116	SER
1	G	358	SER
1	F	331	LEU
1	F	537[B]	MET
1	H	331	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	285	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 36 ligands modelled in this entry, 16 are monoatomic - leaving 20 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	FBP	B	601	-	18,20,20	0.53	0	23,32,32	1.18	3 (13%)
2	FBP	F	601	-	18,20,20	0.64	1 (5%)	23,32,32	0.84	1 (4%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
6	I9K	G	605	-	35,41,41	0.36	0	51,63,63	0.54	0
3	OXL	A	602	4	0,5,5	-	-	0,6,6	-	-
3	OXL	B	602	4	0,5,5	-	-	0,6,6	-	-
2	FBP	C	601	-	18,20,20	0.56	0	23,32,32	1.15	3 (13%)
3	OXL	D	602	4	0,5,5	-	-	0,6,6	-	-
6	I9K	C	605	-	35,41,41	0.34	0	51,63,63	0.68	0
3	OXL	C	602	4	0,5,5	-	-	0,6,6	-	-
2	FBP	E	601	-	18,20,20	0.63	0	23,32,32	0.94	2 (8%)
2	FBP	A	601	-	18,20,20	0.67	0	23,32,32	1.11	1 (4%)
2	FBP	D	601	-	18,20,20	0.65	1 (5%)	23,32,32	1.11	2 (8%)
6	I9K	A	605	-	35,41,41	0.20	0	51,63,63	0.65	1 (1%)
3	OXL	E	602	4	0,5,5	-	-	0,6,6	-	-
3	OXL	G	602	4	0,5,5	-	-	0,6,6	-	-
2	FBP	H	601	-	18,20,20	0.83	0	23,32,32	0.90	1 (4%)
6	I9K	F	605	-	35,41,41	0.38	0	51,63,63	0.82	2 (3%)
3	OXL	F	602	4	0,5,5	-	-	0,6,6	-	-
3	OXL	H	602	4	0,5,5	-	-	0,6,6	-	-
2	FBP	G	601	-	18,20,20	0.61	0	23,32,32	0.85	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	FBP	B	601	-	-	4/13/32/32	0/1/1/1
2	FBP	F	601	-	-	2/13/32/32	0/1/1/1
6	I9K	G	605	-	-	8/24/58/58	0/4/4/4
3	OXL	A	602	4	-	0/0/4/4	-
3	OXL	B	602	4	-	0/0/4/4	-
2	FBP	C	601	-	-	2/13/32/32	0/1/1/1
3	OXL	D	602	4	-	0/0/4/4	-
6	I9K	C	605	-	-	9/24/58/58	0/4/4/4
3	OXL	C	602	4	-	0/0/4/4	-
2	FBP	E	601	-	-	4/13/32/32	0/1/1/1
2	FBP	A	601	-	-	5/13/32/32	0/1/1/1
2	FBP	D	601	-	-	2/13/32/32	0/1/1/1
6	I9K	A	605	-	-	4/24/58/58	0/4/4/4
3	OXL	E	602	4	-	0/0/4/4	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	OXL	G	602	4	-	0/0/4/4	-
2	FBP	H	601	-	-	2/13/32/32	0/1/1/1
6	I9K	F	605	-	-	11/24/58/58	0/4/4/4
3	OXL	F	602	4	-	0/0/4/4	-
3	OXL	H	602	4	-	0/0/4/4	-
2	FBP	G	601	-	-	2/13/32/32	0/1/1/1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	F	601	FBP	P2-O4P	2.28	1.57	1.50
2	D	601	FBP	P2-O5P	-2.11	1.46	1.54

The worst 5 of 16 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	601	FBP	P1-O1-C1	3.87	128.96	118.30
6	F	605	I9K	C12-N-S	3.33	123.09	117.05
2	B	601	FBP	P1-O1-C1	3.27	127.31	118.30
2	D	601	FBP	O5P-P2-O6	3.16	115.15	106.73
6	F	605	I9K	C21-N-S	3.06	122.61	117.05

There are no chirality outliers.

5 of 55 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	601	FBP	O5-C5-C6-O6
2	A	601	FBP	C6-O6-P2-O4P
2	A	601	FBP	C6-O6-P2-O5P
2	A	601	FBP	C6-O6-P2-O6P
2	B	601	FBP	C1-O1-P1-O2P


There are no ring outliers.


3 monomers are involved in 5 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	G	605	I9K	2	0
6	C	605	I9K	1	0
2	A	601	FBP	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



































## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.



## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	422/447 (94%)	0.77	58 (13%) 3 4	49, 75, 107, 120	0
1	B	436/447 (97%)	0.45	29 (6%) 17 26	30, 64, 98, 108	0
1	C	425/447 (95%)	0.20	11 (2%) 56 65	32, 52, 81, 133	0
1	D	425/447 (95%)	0.14	4 (0%) 84 90	29, 41, 73, 122	0
1	E	419/447 (93%)	0.47	28 (6%) 17 26	44, 67, 97, 117	0
1	F	432/447 (96%)	0.21	15 (3%) 44 56	32, 53, 84, 100	0
1	G	421/447 (94%)	0.13	7 (1%) 70 78	30, 46, 73, 106	1 (0%)
1	H	425/447 (95%)	0.10	4 (0%) 84 90	27, 39, 67, 116	0
All	All	3405/3576 (95%)	0.31	156 (4%) 32 45	27, 55, 94, 133	1 (0%)

The worst 5 of 156 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	543	SER	9.8
1	D	24	PHE	8.6
1	D	25	PHE	7.0
1	B	540[A]	LEU	6.5
1	H	24	PHE	6.2

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)


There are no monosaccharides in this entry.

## 6.4 Ligands i

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
5	K	E	604	1/1	0.67	0.25	113,113,113,113	0
5	K	A	604	1/1	0.77	0.14	101,101,101,101	0
6	I9K	A	605	38/38	0.79	0.32	129,130,131,131	22
3	OXL	E	602	6/6	0.87	0.16	85,86,86,86	0
6	I9K	F	605	38/38	0.87	0.25	76,79,89,89	22
5	K	H	604	1/1	0.89	0.08	73,73,73,73	0
6	I9K	C	605	38/38	0.91	0.15	93,94,98,99	21
4	MG	B	603	1/1	0.91	0.11	45,45,45,45	0
6	I9K	G	605	38/38	0.91	0.15	79,86,94,94	22
2	FBP	A	601	20/20	0.92	0.14	81,82,84,84	0
4	MG	D	603	1/1	0.93	0.17	57,57,57,57	0
3	OXL	G	602	6/6	0.93	0.12	47,48,49,49	0
3	OXL	C	602	6/6	0.94	0.12	51,52,53,54	0
3	OXL	A	602	6/6	0.94	0.29	76,77,77,77	0
5	K	F	604	1/1	0.95	0.13	79,79,79,79	0
5	K	C	604	1/1	0.95	0.08	77,77,77,77	0
4	MG	E	603	1/1	0.95	0.05	70,70,70,70	0
5	K	B	604	1/1	0.96	0.10	80,80,80,80	0
2	FBP	E	601	20/20	0.96	0.14	69,70,73,74	0
3	OXL	H	602	6/6	0.96	0.11	48,49,49,49	0
2	FBP	B	601	20/20	0.96	0.13	64,65,67,67	0
5	K	G	604	1/1	0.96	0.08	65,65,65,65	0
2	FBP	F	601	20/20	0.97	0.14	55,60,65,65	0
3	OXL	D	602	6/6	0.97	0.12	43,44,45,46	0
4	MG	G	603	1/1	0.97	0.18	30,30,30,30	0
4	MG	H	603	1/1	0.97	0.18	53,53,53,53	0
4	MG	C	603	1/1	0.97	0.17	59,59,59,59	0
4	MG	A	603	1/1	0.98	0.06	73,73,73,73	0
3	OXL	B	602	6/6	0.98	0.09	48,49,49,49	0
3	OXL	F	602	6/6	0.98	0.15	59,60,61,62	0
2	FBP	H	601	20/20	0.98	0.17	32,35,37,39	0
2	FBP	D	601	20/20	0.98	0.16	35,37,40,40	0
5	K	D	604	1/1	0.98	0.08	67,67,67,67	0
4	MG	F	603	1/1	0.98	0.12	46,46,46,46	0
2	FBP	G	601	20/20	0.99	0.13	36,40,45,45	0
2	FBP	C	601	20/20	0.99	0.14	35,37,42,42	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.






**Electron density around I9K F 605:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around I9K C 605:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)




**Electron density around I9K G 605:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around FBP A 601:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)




**Electron density around FBP E 601:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around FBP B 601:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)




**Electron density around FBP F 601:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around FBP H 601:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

**Electron density around FBP D 601:**




$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)








**Electron density around FBP G 601:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

**Electron density around FBP C 601:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



## 6.5 Other polymers [i](#)

There are no such residues in this entry.