



Full wwPDB X-ray Structure Validation Report ⓘ

Sep 25, 2023 – 04:18 PM EDT

PDB ID : 6BHQ
Title : Mouse Immunoglobulin G 2c Fc fragment with complex-type glycan
Authors : Falconer, D.; Barb, A.
Deposited on : 2017-10-31
Resolution : 2.05 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtrriage (Phenix) : 1.13
EDS : 2.35.1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.35.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.05 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1692 (2.04-2.04)
Clashscore	141614	1773 (2.04-2.04)
Ramachandran outliers	138981	1752 (2.04-2.04)
Sidechain outliers	138945	1752 (2.04-2.04)
RSRZ outliers	127900	1672 (2.04-2.04)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	224	 4% 83% 10% 7%
1	B	224	 14% 72% 14% 11%
2	C	8	 100%
2	D	8	 100%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	NAG	D	1	-	-	-	X
3	GOL	A	509	-	-	X	-

2 Entry composition i


There are 4 unique types of molecules in this entry. The entry contains 3551 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Igh protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	208	Total 1618	C 1028	N 270	O 310	S 10	0	0	0
1	B	199	Total 1542	C 986	N 255	O 291	S 10	0	0	0

- Molecule 2 is an oligosaccharide called beta-D-galactopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-2)-alpha-D-mannopyranose-(1-6)-[alpha-D-mannopyranose-(1-3)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[beta-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
2	C	8	Total 96	C 54	N 3	O 39	0	0	0
2	D	8	Total 96	C 54	N 3	O 39	0	0	0

- Molecule 3 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C O 6 3 3	0	0

- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	100	Total O 100 100	0	0
4	B	93	Total O 93 93	0	0

MAG1
MAG2
BMA3
MAN4
MAG5
GAL6
MAN7
FUL8

4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	67.73Å 73.40Å 118.41Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	46.08 – 2.05 46.08 – 2.05	Depositor EDS
% Data completeness (in resolution range)	99.0 (46.08-2.05) 99.0 (46.08-2.05)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.68 (at 2.05Å)	Xtrriage
Refinement program	PHENIX (1.10.1_2155: ???)	Depositor
R, R_{free}	0.237 , 0.258 0.237 , 0.259	Depositor DCC
R_{free} test set	3698 reflections (9.90%)	wwPDB-VP
Wilson B-factor (Å ²)	40.6	Xtrriage
Anisotropy	0.201	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.29 , 40.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	3551	wwPDB-VP
Average B, all atoms (Å ²)	55.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.12% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, MAN, GAL, NAG, BMA, FUL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.95	0/1659	0.92	0/2264
1	B	1.00	2/1580 (0.1%)	0.99	5/2152 (0.2%)
All	All	0.97	2/3239 (0.1%)	0.95	5/4416 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	317	LYS	C-N	9.38	1.55	1.34
1	B	318	GLU	C-N	-5.69	1.21	1.34

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	318	GLU	CA-C-N	-9.03	97.33	117.20
1	B	317	LYS	C-N-CA	-6.47	105.53	121.70
1	B	370	THR	N-CA-CB	-5.45	99.95	110.30
1	B	321	CYS	CA-CB-SG	5.32	123.58	114.00
1	B	270	ASP	C-N-CD	5.07	139.05	128.40

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	318	GLU	Mainchain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1618	0	1576	19	0
1	B	1542	0	1491	41	0
2	C	96	0	82	0	0
2	D	96	0	82	0	0
3	A	6	0	8	7	0
4	A	100	0	0	1	0
4	B	93	0	0	6	0
All	All	3551	0	3239	59	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (59) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:273:VAL:HG12	1:B:325:ASN:CB	1.41	1.47
1:B:273:VAL:CG1	1:B:325:ASN:HB2	1.50	1.41
1:B:273:VAL:CG1	1:B:325:ASN:CB	2.02	1.37
1:B:273:VAL:HG12	1:B:325:ASN:HB3	1.11	1.08
1:B:273:VAL:HA	1:B:324:ASN:O	1.56	1.04
1:B:273:VAL:HG13	1:B:325:ASN:HB2	1.36	1.03
1:B:416:THR:HG21	4:B:601:HOH:O	1.70	0.91
1:B:273:VAL:HG12	1:B:325:ASN:HB2	1.18	0.88
1:B:273:VAL:HG13	1:B:325:ASN:CB	1.93	0.86
1:B:273:VAL:CA	1:B:324:ASN:O	2.29	0.80
1:A:275:ILE:HG12	1:A:323:VAL:HG12	1.63	0.78
1:B:273:VAL:CG1	1:B:325:ASN:HB3	1.90	0.74
1:B:273:VAL:HG13	1:B:325:ASN:ND2	2.05	0.72
1:A:341:GLY:N	3:A:509:GOL:O3	2.24	0.70
1:A:442:SER:O	1:A:443:ARG:HB3	1.93	0.69
1:B:433:HIS:HD2	4:B:675:HOH:O	1.81	0.64

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:338:LYS:NZ	3:A:509:GOL:O1	2.25	0.62
1:B:273:VAL:HG13	1:B:325:ASN:CG	2.20	0.62
1:A:399:ASP:CG	1:B:411:ARG:HH22	2.04	0.62
1:A:339:PRO:O	3:A:509:GOL:H11	2.01	0.61
1:A:347:GLN:HB3	1:A:370:THR:HG22	1.84	0.58
1:B:416:THR:CG2	4:B:601:HOH:O	2.40	0.58
1:B:290:GLN:NE2	1:B:292:HIS:CE1	2.73	0.57
1:B:273:VAL:HG13	1:B:325:ASN:HD22	1.67	0.57
1:B:413:GLN:O	1:B:416:THR:HG22	2.06	0.56
1:A:340:ARG:HA	3:A:509:GOL:H31	1.88	0.56
1:B:273:VAL:CG1	1:B:325:ASN:CG	2.73	0.56
1:B:333:GLU:O	1:B:334:LYS:HD3	2.05	0.55
1:B:273:VAL:HG21	1:B:302:VAL:HG21	1.91	0.52
1:B:257:PRO:O	1:B:258:MET:HG2	2.11	0.50
1:B:416:THR:O	1:B:421:SER:OG	2.25	0.50
1:B:335:THR:HG23	4:B:677:HOH:O	2.12	0.50
1:A:399:ASP:OD1	1:B:411:ARG:NH2	2.42	0.49
1:B:263:VAL:O	1:B:301:ARG:HA	2.12	0.49
1:A:322:LYS:HG3	1:A:333:GLU:HG2	1.94	0.49
1:B:263:VAL:HG21	1:B:323:VAL:HG21	1.94	0.49
1:B:275:ILE:CD1	1:B:323:VAL:HG23	2.43	0.48
1:B:239:SER:O	1:B:264:VAL:HG22	2.13	0.48
1:A:340:ARG:HA	3:A:509:GOL:C3	2.44	0.48
1:B:268:GLU:O	1:B:269:ASP:C	2.51	0.47
1:B:433:HIS:CD2	4:B:675:HOH:O	2.63	0.47
1:B:292:HIS:HB2	1:B:301:ARG:O	2.15	0.47
1:A:382:THR:HG22	1:A:387:THR:HA	1.97	0.46
1:A:341:GLY:H	3:A:509:GOL:HO3	1.57	0.46
1:B:274:GLN:N	1:B:324:ASN:O	2.48	0.46
3:A:509:GOL:H12	4:A:674:HOH:O	2.16	0.45
1:A:240:VAL:HG21	1:A:323:VAL:HG21	1.99	0.45
1:A:275:ILE:HD12	1:A:304:SER:HB2	1.99	0.44
1:A:414:LYS:HE2	1:A:418:GLU:OE2	2.18	0.43
1:A:275:ILE:HG23	1:A:323:VAL:HG12	2.00	0.43
1:B:290:GLN:O	1:B:290:GLN:HG3	2.19	0.43
1:A:441:ILE:HG12	1:A:442:SER:N	2.34	0.43
1:B:273:VAL:CA	1:B:325:ASN:HB2	2.48	0.42
1:A:440:THR:HG22	1:A:441:ILE:N	2.35	0.42
1:B:428:VAL:HA	1:B:435:HIS:O	2.20	0.42
1:B:273:VAL:HA	1:B:325:ASN:HB2	2.01	0.41
1:B:274:GLN:O	1:B:324:ASN:N	2.49	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:290:GLN:HE21	1:B:292:HIS:CE1	2.37	0.41
1:B:382:THR:HG21	4:B:602:HOH:O	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	206/224 (92%)	204 (99%)	2 (1%)	0	100	100
1	B	191/224 (85%)	186 (97%)	5 (3%)	0	100	100
All	All	397/448 (89%)	390 (98%)	7 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	185/203 (91%)	182 (98%)	3 (2%)	62	59
1	B	174/203 (86%)	166 (95%)	8 (5%)	27	19
All	All	359/406 (88%)	348 (97%)	11 (3%)	40	33

All (11) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	282	VAL
1	A	301	ARG
1	A	370	THR
1	B	253	ILE
1	B	281	ASN
1	B	291	THR
1	B	300	LEU
1	B	323	VAL
1	B	325	ASN
1	B	335	THR
1	B	421	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	290	GLN
1	B	325	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

16 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	NAG	C	1	2,1	14,14,15	0.84	0	17,19,21	2.02	3 (17%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	NAG	C	2	2	14,14,15	0.70	0	17,19,21	1.90	4 (23%)
2	BMA	C	3	2	11,11,12	0.42	0	15,15,17	1.36	1 (6%)
2	MAN	C	4	2	11,11,12	0.34	0	15,15,17	2.23	3 (20%)
2	NAG	C	5	2	14,14,15	0.84	1 (7%)	17,19,21	1.26	1 (5%)
2	GAL	C	6	2	11,11,12	1.37	3 (27%)	15,15,17	1.68	3 (20%)
2	MAN	C	7	2	11,11,12	0.81	0	15,15,17	1.28	3 (20%)
2	FUL	C	8	2	10,10,11	1.02	1 (10%)	14,14,16	2.02	6 (42%)
2	NAG	D	1	2	14,14,15	0.68	0	17,19,21	1.55	5 (29%)
2	NAG	D	2	2	14,14,15	0.40	0	17,19,21	1.63	3 (17%)
2	BMA	D	3	2	11,11,12	0.66	0	15,15,17	2.51	8 (53%)
2	MAN	D	4	2	11,11,12	0.60	0	15,15,17	1.75	2 (13%)
2	NAG	D	5	2	14,14,15	0.68	0	17,19,21	1.72	2 (11%)
2	GAL	D	6	2	11,11,12	1.19	1 (9%)	15,15,17	2.29	5 (33%)
2	MAN	D	7	2	11,11,12	1.11	1 (9%)	15,15,17	2.51	4 (26%)
2	FUL	D	8	2	10,10,11	0.58	0	14,14,16	1.82	3 (21%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '2' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	C	1	2,1	-	2/6/23/26	0/1/1/1
2	NAG	C	2	2	-	1/6/23/26	0/1/1/1
2	BMA	C	3	2	-	0/2/19/22	0/1/1/1
2	MAN	C	4	2	-	0/2/19/22	0/1/1/1
2	NAG	C	5	2	-	0/6/23/26	0/1/1/1
2	GAL	C	6	2	-	1/2/19/22	0/1/1/1
2	MAN	C	7	2	-	2/2/19/22	0/1/1/1
2	FUL	C	8	2	-	-	0/1/1/1
2	NAG	D	1	2	-	2/6/23/26	0/1/1/1
2	NAG	D	2	2	-	1/6/23/26	0/1/1/1
2	BMA	D	3	2	-	0/2/19/22	0/1/1/1
2	MAN	D	4	2	-	0/2/19/22	0/1/1/1
2	NAG	D	5	2	-	2/6/23/26	0/1/1/1
2	GAL	D	6	2	-	0/2/19/22	0/1/1/1
2	MAN	D	7	2	-	0/2/19/22	0/1/1/1
2	FUL	D	8	2	-	-	0/1/1/1

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	6	GAL	O2-C2	2.76	1.49	1.43
2	C	8	FUL	C2-C3	-2.63	1.48	1.52
2	D	7	MAN	C2-C3	2.50	1.56	1.52
2	C	5	NAG	C2-N2	2.28	1.50	1.46
2	D	6	GAL	C1-C2	2.24	1.57	1.52
2	C	6	GAL	C2-C3	2.16	1.55	1.52
2	C	6	GAL	C1-C2	2.02	1.56	1.52

All (56) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	2	NAG	C1-O5-C5	5.81	120.06	112.19
2	D	5	NAG	C1-O5-C5	5.15	119.18	112.19
2	C	1	NAG	C1-O5-C5	4.85	118.76	112.19
2	C	4	MAN	C1-O5-C5	4.77	118.65	112.19
2	D	7	MAN	C1-C2-C3	4.74	115.49	109.67
2	D	7	MAN	C1-O5-C5	4.62	118.46	112.19
2	D	6	GAL	O3-C3-C2	-4.61	101.16	109.99
2	D	4	MAN	C1-C2-C3	4.60	115.32	109.67
2	C	4	MAN	C1-C2-C3	4.52	115.22	109.67
2	D	7	MAN	C2-C3-C4	4.48	118.65	110.89
2	C	1	NAG	C1-C2-N2	-4.32	103.11	110.49
2	D	3	BMA	C1-C2-C3	4.28	114.93	109.67
2	D	3	BMA	O5-C5-C6	4.20	113.80	107.20
2	C	1	NAG	O5-C1-C2	-4.16	104.71	111.29
2	C	3	BMA	C1-O5-C5	4.09	117.74	112.19
2	C	4	MAN	O2-C2-C3	-3.95	102.23	110.14
2	C	6	GAL	O5-C5-C6	3.94	113.38	107.20
2	D	7	MAN	C3-C4-C5	3.92	117.24	110.24
2	D	6	GAL	O2-C2-C1	3.89	117.12	109.15
2	D	6	GAL	C1-O5-C5	3.80	117.34	112.19
2	D	8	FUL	C1-C2-C3	3.71	114.23	109.67
2	D	2	NAG	C4-C3-C2	-3.71	105.58	111.02
2	D	3	BMA	O5-C5-C4	-3.69	101.86	110.83
2	C	6	GAL	O2-C2-C1	3.27	115.84	109.15
2	D	1	NAG	O5-C1-C2	-3.26	106.14	111.29
2	D	4	MAN	O5-C5-C6	3.25	112.30	107.20
2	C	8	FUL	C1-O5-C5	3.21	120.05	112.78
2	D	5	NAG	O4-C4-C3	-3.20	102.96	110.35
2	D	8	FUL	C3-C4-C5	3.19	114.74	109.77
2	D	3	BMA	O5-C1-C2	3.19	115.69	110.77
2	D	3	BMA	O3-C3-C2	-3.13	104.00	109.99

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	2	NAG	C1-O5-C5	3.04	116.31	112.19
2	C	5	NAG	O3-C3-C2	-2.92	103.42	109.47
2	D	1	NAG	C1-O5-C5	2.87	116.08	112.19
2	C	8	FUL	C2-C3-C4	-2.81	106.03	110.89
2	C	8	FUL	O5-C5-C4	2.81	114.56	109.52
2	D	6	GAL	O2-C2-C3	-2.80	104.52	110.14
2	D	2	NAG	C1-C2-N2	2.77	115.22	110.49
2	C	8	FUL	C3-C4-C5	2.70	113.97	109.77
2	D	8	FUL	O5-C1-C2	2.68	114.91	110.77
2	D	6	GAL	C1-C2-C3	2.67	112.95	109.67
2	D	3	BMA	C3-C4-C5	2.61	114.90	110.24
2	C	2	NAG	O5-C5-C4	2.61	117.18	110.83
2	D	3	BMA	C2-C3-C4	2.48	115.19	110.89
2	C	7	MAN	C3-C4-C5	2.43	114.57	110.24
2	C	7	MAN	O4-C4-C3	-2.38	104.84	110.35
2	C	6	GAL	O5-C5-C4	-2.34	105.12	110.83
2	C	2	NAG	O4-C4-C5	-2.32	103.54	109.30
2	D	1	NAG	O5-C5-C6	2.27	110.76	107.20
2	C	8	FUL	O5-C1-C2	2.25	114.24	110.77
2	D	3	BMA	O3-C3-C4	2.22	115.49	110.35
2	C	8	FUL	O2-C2-C3	-2.22	105.69	110.14
2	D	1	NAG	C1-C2-N2	2.14	114.14	110.49
2	D	1	NAG	C4-C3-C2	-2.11	107.92	111.02
2	C	7	MAN	O5-C5-C6	2.08	110.47	107.20
2	C	2	NAG	C1-C2-N2	-2.03	107.03	110.49

There are no chirality outliers.


All (11) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	D	1	NAG	O5-C5-C6-O6
2	C	1	NAG	C4-C5-C6-O6
2	D	1	NAG	C4-C5-C6-O6
2	C	7	MAN	O5-C5-C6-O6
2	C	1	NAG	O5-C5-C6-O6
2	C	7	MAN	C4-C5-C6-O6
2	D	5	NAG	C4-C5-C6-O6
2	D	5	NAG	O5-C5-C6-O6
2	D	2	NAG	O5-C5-C6-O6
2	C	2	NAG	C4-C5-C6-O6
2	C	6	GAL	O5-C5-C6-O6

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.





5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	GOL	A	509	-	5,5,5	0.37	0	5,5,5	0.65	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	A	509	-	-	2/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	509	GOL	C1-C2-C3-O3
3	A	509	GOL	O2-C2-C3-O3

There are no ring outliers.

1 monomer is involved in 7 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	509	GOL	7	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	208/224 (92%)	0.43	8 (3%) 40 44	28, 51, 72, 94	0
1	B	199/224 (88%)	1.00	31 (15%) 2 1	27, 49, 113, 143	0
All	All	407/448 (90%)	0.71	39 (9%) 8 8	27, 50, 94, 143	0

All (39) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	272	ASP	7.3
1	B	293	ARG	6.7
1	B	296	TYR	6.1
1	A	236	GLY	6.0
1	B	302	VAL	5.9
1	B	273	VAL	5.8
1	B	291	THR	5.3
1	B	271	PRO	4.6
1	B	332	ILE	4.5
1	B	295	ASP	4.3
1	B	324	ASN	4.0
1	B	292	HIS	3.8
1	B	330	SER	3.7
1	B	323	VAL	3.4
1	B	301	ARG	3.3
1	B	238	PRO	3.3
1	B	263	VAL	3.3
1	A	327	ALA	3.3
1	B	294	GLU	3.2
1	B	270	ASP	3.0
1	B	268	GLU	3.0
1	B	243	PHE	3.0
1	B	241	PHE	2.9
1	B	325	ASN	2.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	237	GLY	2.9
1	B	239	SER	2.8
1	B	265	ASP	2.8
1	B	274	GLN	2.7
1	B	269	ASP	2.6
1	B	262	VAL	2.6
1	B	420	GLY	2.6
1	A	422	LEU	2.4
1	B	359	THR	2.4
1	B	331	PRO	2.4
1	B	300	LEU	2.3
1	A	272	ASP	2.3
1	A	384	ASN	2.3
1	A	419	ARG	2.2
1	A	329	PRO	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	MAN	D	7	11/12	0.49	0.39	96,101,106,107	0
2	NAG	D	1	14/15	0.50	0.41	116,125,141,142	0
2	NAG	D	5	14/15	0.59	0.25	76,82,84,85	0
2	MAN	C	7	11/12	0.70	0.23	92,94,96,98	0
2	NAG	D	2	14/15	0.78	0.26	101,107,114,115	0
2	BMA	D	3	11/12	0.80	0.18	80,84,91,94	0
2	FUL	D	8	10/11	0.80	0.32	98,105,109,110	0
2	NAG	C	5	14/15	0.86	0.17	51,52,53,54	0
2	MAN	D	4	11/12	0.87	0.23	105,112,114,118	0
2	GAL	D	6	11/12	0.87	0.19	63,66,69,71	0
2	BMA	C	3	11/12	0.88	0.10	50,53,54,56	0
2	GAL	C	6	11/12	0.89	0.28	52,52,55,56	0
2	FUL	C	8	10/11	0.89	0.15	67,68,69,70	0


Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	NAG	C	2	14/15	0.93	0.10	51,52,56,57	0
2	NAG	C	1	14/15	0.94	0.12	53,56,58,58	0
2	MAN	C	4	11/12	0.94	0.12	51,53,54,54	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.





6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	GOL	A	509	6/6	0.84	0.39	20,20,20,20	0

6.5 Other polymers [i](#)

There are no such residues in this entry.