



wwPDB EM Validation Summary Report ⓘ

Nov 8, 2022 – 03:06 AM EST

PDB ID : 6D1W
EMDB ID : EMD-7786
Title : human PKD2 F604P mutant
Authors : Zheng, W.; Yang, X.; Bulkley, D.; Chen, X.Z.; Cao, E.
Deposited on : 2018-04-12
Resolution : 3.54 Å (reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:


EMDB validation analysis : 0.0.1.dev43
Mogul : 1.8.5 (274361), CSD as541be (2020)
MolProbity : 4.02b-467
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
MapQ : 1.9.9
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.31.2

1 Overall quality at a glance

The following experimental techniques were used to determine the structure:
ELECTRON MICROSCOPY

The reported resolution of this entry is 3.54 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)
Clashscore	158937	4297
Ramachandran outliers	154571	4023
Sidechain outliers	154315	3826

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	740	
1	B	740	
1	C	740	
1	D	740	

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 14492 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Polycystin-2.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
1	A	451	3581	2381	575	607	18	0	0
1	B	451	3581	2381	575	607	18	0	0
1	C	451	3581	2381	575	607	18	0	0
1	D	451	3581	2381	575	607	18	0	0

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	604	PRO	PHE	engineered mutation	UNP Q13563
B	604	PRO	PHE	engineered mutation	UNP Q13563
C	604	PRO	PHE	engineered mutation	UNP Q13563
D	604	PRO	PHE	engineered mutation	UNP Q13563

- Molecule 2 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: C₈H₁₅NO₆).




Mol	Chain	Residues	Atoms				AltConf
			Total	C	N	O	
2	A	1	Total	C	N	O	0
			42	24	3	15	
2	A	1	Total	C	N	O	0
			42	24	3	15	
2	A	1	Total	C	N	O	0
			42	24	3	15	
2	B	1	Total	C	N	O	0
			42	24	3	15	
2	B	1	Total	C	N	O	0
			42	24	3	15	
2	B	1	Total	C	N	O	0
			42	24	3	15	
2	C	1	Total	C	N	O	0
			42	24	3	15	
2	C	1	Total	C	N	O	0
			42	24	3	15	
2	C	1	Total	C	N	O	0
			42	24	3	15	
2	D	1	Total	C	N	O	0
			42	24	3	15	
2	D	1	Total	C	N	O	0
			42	24	3	15	
2	D	1	Total	C	N	O	0
			42	24	3	15	


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Polycystin-2



• Molecule 1: Polycystin-2



• Molecule 1: Polycystin-2



VAL
LYS
SER
ASP
LEU
THR
ALA
GLN
GLN
LYS
GLU
ALA
MET
GLU
LEU
SER
TYR
ASP
LEU
ILE
ARG
GLY
LYS
GLY
TYR
HIS
HIS
LYS
ALA
LEU
VAL
LYS
LYS
LYS
LEU
LYS
LYS
LYS
ASN
THR
VAL
ASP
ASP
ASP
ILE
ILE
GLU
SER
GLU
SER
LEU
ARG
GLN
GLY
GLY
LYS
LEU
ASN
PHE
ASP
GLU
LEU
ARG
GLN
ASP
LEU
LEU
LYS

GLY
LYS
GLY
HIS
THR
ASP
ALA
GLU
ILE
GLU
ALA
ILE
PHE
THR
LYS
TYR
ASP
GLN
ASP
GLY
ASP
GLN
GLU
GLU
THR
GLU
HIS
HIS
GLN
GLN
MET
ARG
ASP
ASP
LEU
GLU
LYS
GLU
ARG
SER
GLU
ASP
LEU
ASP
LEU
LEU
ASP

4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	387454	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	FEI POLARA 300	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	40	Depositor
Minimum defocus (nm)	Not provided	
Maximum defocus (nm)	Not provided	
Magnification	Not provided	
Image detector	GATAN K2 SUMMIT (4k x 4k)	Depositor
Maximum map value	0.219	Depositor
Minimum map value	-0.125	Depositor
Average map value	0.001	Depositor
Map value standard deviation	0.008	Depositor
Recommended contour level	0.04	Depositor
Map size (Å)	233.3952, 233.3952, 233.3952	wwPDB
Map dimensions	192, 192, 192	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	1.2156, 1.2156, 1.2156	Depositor

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.44	1/3677 (0.0%)	0.47	0/5008
1	B	0.44	1/3677 (0.0%)	0.47	0/5008
1	C	0.44	1/3677 (0.0%)	0.47	0/5008
1	D	0.44	1/3677 (0.0%)	0.47	0/5008
All	All	0.44	4/14708 (0.0%)	0.47	0/20032

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	356	ALA	C-N	-10.18	1.15	1.34
1	A	356	ALA	C-N	-10.14	1.15	1.34
1	D	356	ALA	C-N	-10.14	1.15	1.34
1	B	356	ALA	C-N	-10.13	1.15	1.34

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3581	0	3473	147	0
1	B	3581	0	3473	146	0
1	C	3581	0	3473	147	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	D	3581	0	3473	147	0
2	A	42	0	39	0	0
2	B	42	0	39	0	0
2	C	42	0	39	0	0
2	D	42	0	39	0	0
All	All	14492	0	14048	531	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 19.

The worst 5 of 531 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:319:PRO:HB2	1:D:395:LEU:HD12	1.60	0.83
1:B:319:PRO:HB2	1:B:395:LEU:HD12	1.61	0.83
1:A:319:PRO:HB2	1:A:395:LEU:HD12	1.61	0.82
1:C:319:PRO:HB2	1:C:395:LEU:HD12	1.60	0.82
1:B:505:SER:H	1:B:508:ASN:HB2	1.57	0.70

There are no symmetry-related clashes.

5.3 Torsion angles [\(i\)](#)

5.3.1 Protein backbone [\(i\)](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	445/740 (60%)	415 (93%)	29 (6%)	1 (0%)	47 80
1	B	445/740 (60%)	415 (93%)	29 (6%)	1 (0%)	47 80
1	C	445/740 (60%)	415 (93%)	29 (6%)	1 (0%)	47 80
1	D	445/740 (60%)	415 (93%)	29 (6%)	1 (0%)	47 80
All	All	1780/2960 (60%)	1660 (93%)	116 (6%)	4 (0%)	50 80

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	532	VAL
1	B	532	VAL
1	C	532	VAL
1	D	532	VAL

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	368/647 (57%)	368 (100%)	0	100	100
1	B	368/647 (57%)	368 (100%)	0	100	100
1	C	368/647 (57%)	368 (100%)	0	100	100
1	D	368/647 (57%)	368 (100%)	0	100	100
All	All	1472/2588 (57%)	1472 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 20 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	636	GLN
1	D	543	ASN
1	D	636	GLN
1	D	560	ASN
1	B	458	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

12 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	NAG	B	801	1	14,14,15	0.19	0	17,19,21	0.43	0
2	NAG	A	801	1	14,14,15	0.19	0	17,19,21	0.43	0
2	NAG	C	803	1	14,14,15	0.19	0	17,19,21	0.34	0
2	NAG	A	803	1	14,14,15	0.19	0	17,19,21	0.34	0
2	NAG	A	802	1	14,14,15	0.38	0	17,19,21	0.65	1 (5%)
2	NAG	C	801	1	14,14,15	0.20	0	17,19,21	0.43	0
2	NAG	D	801	1	14,14,15	0.19	0	17,19,21	0.44	0
2	NAG	B	802	1	14,14,15	0.36	0	17,19,21	0.67	1 (5%)
2	NAG	D	803	1	14,14,15	0.20	0	17,19,21	0.33	0
2	NAG	D	802	1	14,14,15	0.37	0	17,19,21	0.66	1 (5%)
2	NAG	C	802	1	14,14,15	0.36	0	17,19,21	0.66	1 (5%)
2	NAG	B	803	1	14,14,15	0.20	0	17,19,21	0.34	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	B	801	1	-	2/6/23/26	0/1/1/1
2	NAG	A	801	1	-	2/6/23/26	0/1/1/1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	C	803	1	-	4/6/23/26	0/1/1/1
2	NAG	A	803	1	-	4/6/23/26	0/1/1/1
2	NAG	A	802	1	-	2/6/23/26	0/1/1/1
2	NAG	C	801	1	-	2/6/23/26	0/1/1/1
2	NAG	D	801	1	-	2/6/23/26	0/1/1/1
2	NAG	B	802	1	-	2/6/23/26	0/1/1/1
2	NAG	D	803	1	-	4/6/23/26	0/1/1/1
2	NAG	D	802	1	-	2/6/23/26	0/1/1/1
2	NAG	C	802	1	-	2/6/23/26	0/1/1/1
2	NAG	B	803	1	-	4/6/23/26	0/1/1/1

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	802	NAG	C1-O5-C5	2.37	115.40	112.19
2	C	802	NAG	C1-O5-C5	2.33	115.35	112.19
2	D	802	NAG	C1-O5-C5	2.33	115.35	112.19
2	A	802	NAG	C1-O5-C5	2.30	115.31	112.19

There are no chirality outliers.

5 of 32 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	801	NAG	C4-C5-C6-O6
2	B	801	NAG	C4-C5-C6-O6
2	C	801	NAG	C4-C5-C6-O6
2	D	801	NAG	C4-C5-C6-O6
2	A	802	NAG	C4-C5-C6-O6

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues

The following chains have linkage breaks:

Mol	Chain	Number of breaks
1	A	1
1	B	1
1	D	1
1	C	1

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	A	356:ALA	C	357:PRO	N	1.15
1	B	356:ALA	C	357:PRO	N	1.15
1	D	356:ALA	C	357:PRO	N	1.15
1	C	356:ALA	C	357:PRO	N	1.14


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-7786. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X




Y




Z


6.1.2 Raw map



X



Y



Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map



X Index: 96




Y Index: 96




Z Index: 96


6.2.2 Raw map



X Index: 96



Y Index: 96




Z Index: 96


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 109




Y Index: 109




Z Index: 78


6.3.2 Raw map



X Index: 96



Y Index: 96




Z Index: 78

The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal surface views [i](#)

6.4.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.04. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

6.4.2 Raw map



These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.


6.5 Mask visualisation [i](#)

This section was not generated. No masks/segmentation were deposited.

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [\(i\)](#)



The volume at the recommended contour level is 103 nm³; this corresponds to an approximate mass of 93 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum i




*Reported resolution corresponds to spatial frequency of 0.282 Å⁻¹

8 Fourier-Shell correlation [\(i\)](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC [\(i\)](#)



*Reported resolution corresponds to spatial frequency of 0.282 \AA^{-1}

8.2 Resolution estimates [i](#)


Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	3.54	-	-
Author-provided FSC curve	-	-	-
Unmasked-calculated*	3.80	4.37	3.94

*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps.

9 Map-model fit [i](#)


This section contains information regarding the fit between EMDB map EMD-7786 and PDB model 6D1W. Per-residue inclusion information can be found in section [3](#) on page [5](#).

9.1 Map-model overlay [i](#)




The images above show the 3D surface view of the map at the recommended contour level 0.04 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.04).


9.4 Atom inclusion [i](#)



At the recommended contour level, 85% of all backbone atoms, 73% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary [i](#)

The table lists the average atom inclusion at the recommended contour level (0.04) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	 0.7267	 0.4050
A	 0.7266	 0.4050
B	 0.7258	 0.4040
C	 0.7275	 0.4050
D	 0.7269	 0.4040

