



Full wwPDB X-ray Structure Validation Report ⓘ

Jun 23, 2024 – 01:24 AM EDT

PDB ID : 6JN2
Title : Crystal structure of the coiled-coil domains of human DOT1L in complex with AF10
Authors : Song, X.; Wang, M.; Yang, N.; Xu, R.M.
Deposited on : 2019-03-13
Resolution : 3.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.20.1
EDS : 2.37.1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.37.1


1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1257 (3.70-3.50)
Clashscore	141614	1353 (3.70-3.50)
Ramachandran outliers	138981	1307 (3.70-3.50)
Sidechain outliers	138945	1307 (3.70-3.50)
RSRZ outliers	127900	1161 (3.70-3.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	98	
2	B	82	

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 1098 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Protein AF-10.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	67	547	342	99	105	1	0	0	0

There are 14 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	685	MET	-	expression tag	UNP P55197
A	686	GLY	-	expression tag	UNP P55197
A	687	SER	-	expression tag	UNP P55197
A	688	SER	-	expression tag	UNP P55197
A	689	HIS	-	expression tag	UNP P55197
A	690	HIS	-	expression tag	UNP P55197
A	691	HIS	-	expression tag	UNP P55197
A	692	HIS	-	expression tag	UNP P55197
A	693	HIS	-	expression tag	UNP P55197
A	694	HIS	-	expression tag	UNP P55197
A	695	SER	-	expression tag	UNP P55197
A	696	GLN	-	expression tag	UNP P55197
A	697	ASP	-	expression tag	UNP P55197
A	698	PRO	-	expression tag	UNP P55197

- Molecule 2 is a protein called Histone-lysine N-methyltransferase, H3 lysine-79 specific.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	67	551	353	96	101	1	0	0	0

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	469	MET	-	expression tag	UNP Q8TEK3

4 Data and refinement statistics

Property	Value	Source
Space group	I 41 2 2	Depositor
Cell constants a, b, c, α , β , γ	107.81Å 107.81Å 114.03Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	50.00 – 3.60 31.69 – 3.60	Depositor EDS
% Data completeness (in resolution range)	98.3 (50.00-3.60) 98.8 (31.69-3.60)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.43 (at 3.56Å)	Xtrriage
Refinement program	REFMAC 5.6.0117	Depositor
R, R_{free}	0.324 , 0.381 0.320 , 0.373	Depositor DCC
R_{free} test set	387 reflections (9.55%)	wwPDB-VP
Wilson B-factor (Å ²)	145.7	Xtrriage
Anisotropy	0.049	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.27 , 122.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	1098	wwPDB-VP
Average B, all atoms (Å ²)	191.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.18% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.42	0/552	0.52	0/741
2	B	0.33	0/558	0.58	0/746
All	All	0.38	0/1110	0.55	0/1487

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	547	0	565	18	0
2	B	551	0	570	38	0
All	All	1098	0	1135	47	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 21.

All (47) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:504:THR:CB	2:B:505:PRO:HD3	1.49	1.31
2:B:504:THR:HB	2:B:505:PRO:CD	1.66	1.24

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:499:LEU:O	2:B:503:LYS:HE3	1.45	1.14
2:B:504:THR:CG2	2:B:505:PRO:HD3	1.94	0.97
2:B:504:THR:HB	2:B:505:PRO:HD3	0.90	0.88
2:B:499:LEU:O	2:B:503:LYS:CE	2.22	0.87
2:B:504:THR:CB	2:B:505:PRO:CD	2.35	0.84
2:B:499:LEU:O	2:B:503:LYS:HG3	1.79	0.83
2:B:495:TYR:O	2:B:499:LEU:HB2	1.86	0.75
2:B:488:LEU:HA	2:B:491:PHE:HD2	1.56	0.70
1:A:770:ARG:HA	1:A:773:LEU:HD12	1.75	0.68
1:A:764:LEU:HD11	2:B:533:LEU:HG	1.77	0.66
2:B:499:LEU:O	2:B:503:LYS:CD	2.46	0.64
2:B:499:LEU:O	2:B:503:LYS:CG	2.45	0.64
2:B:492:LYS:HA	2:B:495:TYR:HD2	1.63	0.64
1:A:747:LEU:HG	2:B:515:LEU:HD12	1.82	0.62
1:A:757:LEU:HD23	2:B:525:LEU:CB	2.29	0.62
1:A:757:LEU:HD23	2:B:525:LEU:HB3	1.80	0.61
2:B:533:LEU:O	2:B:537:GLN:HG2	2.00	0.60
2:B:495:TYR:O	2:B:499:LEU:CB	2.48	0.60
1:A:717:GLU:HA	1:A:720:LEU:HB2	1.84	0.60
2:B:501:TYR:CD2	2:B:502:THR:HG23	2.37	0.59
2:B:504:THR:HG22	2:B:505:PRO:HD3	1.85	0.57
1:A:767:LYS:O	1:A:771:LEU:HB2	2.04	0.57
1:A:771:LEU:HD21	2:B:540:LYS:HG3	1.88	0.54
2:B:492:LYS:HA	2:B:495:TYR:CD2	2.43	0.53
2:B:507:TYR:CD2	2:B:508:LYS:HG3	2.45	0.52
2:B:520:GLU:O	2:B:524:GLN:HG3	2.11	0.51
2:B:499:LEU:HD12	2:B:503:LYS:HE3	1.92	0.50
1:A:747:LEU:HD22	2:B:514:LEU:HD22	1.94	0.50
1:A:716:ILE:HG13	1:A:720:LEU:HD12	1.94	0.49
2:B:504:THR:CG2	2:B:505:PRO:CD	2.81	0.48
2:B:499:LEU:C	2:B:503:LYS:HE3	2.28	0.48
1:A:757:LEU:HD21	2:B:526:LEU:HD23	1.96	0.48
2:B:504:THR:HG22	2:B:505:PRO:CD	2.45	0.47
2:B:521:LYS:O	2:B:524:GLN:HB2	2.15	0.46
2:B:496:LEU:HA	2:B:499:LEU:HB3	1.98	0.45
2:B:501:TYR:CE2	2:B:502:THR:HG23	2.52	0.45
2:B:500:ALA:HA	2:B:503:LYS:HD2	1.99	0.44
1:A:719:LEU:HA	1:A:722:ARG:HD2	1.99	0.43
1:A:756:ARG:NH1	1:A:760:GLN:HB2	2.34	0.42
1:A:755:ARG:O	1:A:758:GLU:HB2	2.20	0.42
1:A:718:GLN:O	1:A:722:ARG:HG3	2.21	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:490:SER:HA	2:B:493:ILE:HD12	2.01	0.41
1:A:757:LEU:HD23	2:B:525:LEU:HB2	1.99	0.41
1:A:750:LEU:HB2	2:B:518:GLU:OE2	2.21	0.41
1:A:774:LEU:O	1:A:777:GLN:HB2	2.22	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	65/98 (66%)	64 (98%)	1 (2%)	0	100	100
2	B	65/82 (79%)	58 (89%)	6 (9%)	1 (2%)	10	47
All	All	130/180 (72%)	122 (94%)	7 (5%)	1 (1%)	19	59

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	505	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	62/89 (70%)	59 (95%)	3 (5%)	25	60

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	B	58/72 (81%)	58 (100%)	0	100	100
All	All	120/161 (74%)	117 (98%)	3 (2%)	47	75

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	716	ILE
1	A	744	LEU
1	A	759	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	67/98 (68%)	-0.37	0 100 100	110, 178, 243, 266	0
2	B	67/82 (81%)	-0.19	0 100 100	110, 189, 258, 273	0
All	All	134/180 (74%)	-0.28	0 100 100	110, 188, 258, 273	0

There are no RSRZ outliers to report.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.