wwPDB X-ray Structure Validation Summary Report (i)

Nov 16, 2023 – 06:20 AM JST

PDB ID	:	6L8Q
Title	:	Complex structure of bat CD26 and MERS-RBD
Authors	:	Yuan, Y.
Deposited on	:	2019-11-07
Resolution	:	3.10 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.36
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.36

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 3.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1094 (3.10-3.10)
Clashscore	141614	1184 (3.10-3.10)
Ramachandran outliers	138981	1141 (3.10-3.10)
Sidechain outliers	138945	1141 (3.10-3.10)
RSRZ outliers	127900	1067 (3.10-3.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	731	80% 18%
1	C	731	77% 21%
1	E	731	83% 15%
1	G	731	80% 18%
2	B	246	73% 11% 15%
2	D	246	70% 15% 15%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	F	246	
2	H	246	
3	I	2	
3	K	2	
3	L	2	
3	M	2	
3	O	2	
3	P	2	
3	Q	2	
3	R	2	
3	T	2	
3	U	2	
3	V	2	
3	X	2	
3	Y	2	
4	J	3	
4	N	3	
4	S	3	
4	W	3	

2 Entry composition (i)

There are 5 unique types of molecules in this entry. The entry contains 30730 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Dipeptidyl peptidase 4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	724	Total	C	N	O	S	0	0	0
			5906	3775	993	1114	24			
1	C	724	Total	C	N	O	S	0	0	0
			5906	3775	993	1114	24			
1	E	724	Total	C	N	O	S	0	0	0
			5906	3775	993	1114	24			
1	G	724	Total	C	N	O	S	0	0	0
			5906	3775	993	1114	24			

There are 24 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	762	HIS	-	expression tag	UNP L5LQ33
A	763	HIS	-	expression tag	UNP L5LQ33
A	764	HIS	-	expression tag	UNP L5LQ33
A	765	HIS	-	expression tag	UNP L5LQ33
A	766	HIS	-	expression tag	UNP L5LQ33
A	767	HIS	-	expression tag	UNP L5LQ33
C	762	HIS	-	expression tag	UNP L5LQ33
C	763	HIS	-	expression tag	UNP L5LQ33
C	764	HIS	-	expression tag	UNP L5LQ33
C	765	HIS	-	expression tag	UNP L5LQ33
C	766	HIS	-	expression tag	UNP L5LQ33
C	767	HIS	-	expression tag	UNP L5LQ33
E	762	HIS	-	expression tag	UNP L5LQ33
E	763	HIS	-	expression tag	UNP L5LQ33
E	764	HIS	-	expression tag	UNP L5LQ33
E	765	HIS	-	expression tag	UNP L5LQ33
E	766	HIS	-	expression tag	UNP L5LQ33
E	767	HIS	-	expression tag	UNP L5LQ33
G	762	HIS	-	expression tag	UNP L5LQ33
G	763	HIS	-	expression tag	UNP L5LQ33
G	764	HIS	-	expression tag	UNP L5LQ33

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
G	765	HIS	-	expression tag	UNP L5LQ33
G	766	HIS	-	expression tag	UNP L5LQ33
G	767	HIS	-	expression tag	UNP L5LQ33

- Molecule 2 is a protein called Spike glycoprotein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	208	Total	C	N	O	S	0	0	0
			1608	1026	256	315	11			
2	D	208	Total	C	N	O	S	0	0	0
			1608	1026	256	315	11			
2	F	208	Total	C	N	O	S	0	0	0
			1608	1026	256	315	11			
2	H	208	Total	C	N	O	S	0	0	0
			1608	1026	256	315	11			

There are 24 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	607	HIS	-	expression tag	UNP A0A0A0Q7F3
B	608	HIS	-	expression tag	UNP A0A0A0Q7F3
B	609	HIS	-	expression tag	UNP A0A0A0Q7F3
B	610	HIS	-	expression tag	UNP A0A0A0Q7F3
B	611	HIS	-	expression tag	UNP A0A0A0Q7F3
B	612	HIS	-	expression tag	UNP A0A0A0Q7F3
D	607	HIS	-	expression tag	UNP A0A0A0Q7F3
D	608	HIS	-	expression tag	UNP A0A0A0Q7F3
D	609	HIS	-	expression tag	UNP A0A0A0Q7F3
D	610	HIS	-	expression tag	UNP A0A0A0Q7F3
D	611	HIS	-	expression tag	UNP A0A0A0Q7F3
D	612	HIS	-	expression tag	UNP A0A0A0Q7F3
F	607	HIS	-	expression tag	UNP A0A0A0Q7F3
F	608	HIS	-	expression tag	UNP A0A0A0Q7F3
F	609	HIS	-	expression tag	UNP A0A0A0Q7F3
F	610	HIS	-	expression tag	UNP A0A0A0Q7F3
F	611	HIS	-	expression tag	UNP A0A0A0Q7F3
F	612	HIS	-	expression tag	UNP A0A0A0Q7F3
H	607	HIS	-	expression tag	UNP A0A0A0Q7F3
H	608	HIS	-	expression tag	UNP A0A0A0Q7F3
H	609	HIS	-	expression tag	UNP A0A0A0Q7F3
H	610	HIS	-	expression tag	UNP A0A0A0Q7F3
H	611	HIS	-	expression tag	UNP A0A0A0Q7F3

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
H	612	HIS	-	expression tag	UNP A0A0A0Q7F3

- Molecule 3 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf	Trace
3	I	2	Total C N O 28 16 2 10	0	0	0
3	K	2	Total C N O 28 16 2 10	0	0	0
3	L	2	Total C N O 28 16 2 10	0	0	0
3	M	2	Total C N O 28 16 2 10	0	0	0
3	O	2	Total C N O 28 16 2 10	0	0	0
3	P	2	Total C N O 28 16 2 10	0	0	0
3	Q	2	Total C N O 28 16 2 10	0	0	0
3	R	2	Total C N O 28 16 2 10	0	0	0
3	T	2	Total C N O 28 16 2 10	0	0	0
3	U	2	Total C N O 28 16 2 10	0	0	0
3	V	2	Total C N O 28 16 2 10	0	0	0
3	X	2	Total C N O 28 16 2 10	0	0	0
3	Y	2	Total C N O 28 16 2 10	0	0	0

- Molecule 4 is an oligosaccharide called beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf	Trace
4	J	3	Total C N O 39 22 2 15	0	0	0
4	N	3	Total C N O 39 22 2 15	0	0	0
4	S	3	Total C N O 39 22 2 15	0	0	0
4	W	3	Total C N O 39 22 2 15	0	0	0

- Molecule 5 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: $C_8H_{15}NO_6$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total C N O 14 8 1 5	0	0
5	A	1	Total C N O 14 8 1 5	0	0
5	B	1	Total C N O 14 8 1 5	0	0
5	C	1	Total C N O 14 8 1 5	0	0
5	C	1	Total C N O 14 8 1 5	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	D	1	Total	C	N	O	0	0
			14	8	1	5		
5	E	1	Total	C	N	O	0	0
			14	8	1	5		
5	F	1	Total	C	N	O	0	0
			14	8	1	5		
5	G	1	Total	C	N	O	0	0
			14	8	1	5		
5	G	1	Total	C	N	O	0	0
			14	8	1	5		
5	H	1	Total	C	N	O	0	0
			14	8	1	5		

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Dipeptidyl peptidase 4

Chain A:



• Molecule 1: Dipeptidyl peptidase 4

Chain C:





• Molecule 1: Dipeptidyl peptidase 4

Chain E:



• Molecule 1: Dipeptidyl peptidase 4

Chain G:



• Molecule 2: Spike glycoprotein

Chain B:





• Molecule 2: Spike glycoprotein

Chain D:



• Molecule 2: Spike glycoprotein

Chain F:



• Molecule 2: Spike glycoprotein

Chain H:



• Molecule 3: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain I:

• Molecule 3: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain K:



• Molecule 3: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain L:



• Molecule 3: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain M:

• Molecule 3: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain O:



• Molecule 3: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain P:



• Molecule 3: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain Q:

• Molecule 3: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain R:

• Molecule 3: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain T:

• Molecule 3: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain U:



• Molecule 3: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain V:

• Molecule 3: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain X:



• Molecule 3: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain Y:



• Molecule 4: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain J:



• Molecule 4: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain N:

• Molecule 4: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain S:

• Molecule 4: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain W:

4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α, β, γ	114.66Å 273.67Å 115.23Å 90.00° 119.68° 90.00°	Depositor
Resolution (Å)	49.81 – 3.10 49.81 – 3.10	Depositor EDS
% Data completeness (in resolution range)	85.3 (49.81-3.10) 84.9 (49.81-3.10)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ [1]	2.83 (at 3.12Å)	Xtriage
Refinement program	PHENIX 1.16_3549	Depositor
R, R_{free}	0.200 , 0.241 0.200 , 0.241	Depositor DCC
R_{free} test set	4577 reflections (4.80%)	wwPDB-VP
Wilson B-factor (Å^2)	64.0	Xtriage
Anisotropy	0.278	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.30 , 12.9	EDS
L-test for twinning[2]	$< \|L\| > = 0.49$, $< L^2 > = 0.32$	Xtriage
Estimated twinning fraction	0.007 for -h-l,k,h 0.007 for l,k,-h-l 0.438 for h,-k,-h-l 0.019 for -h-l,-k,l 0.015 for l,-k,h	Xtriage
F_o,F_c correlation	0.91	EDS
Total number of atoms	30730	wwPDB-VP
Average B, all atoms (Å^2)	63.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.92% of the height of the origin peak. No significant pseudotranslation is detected.*

[1] Intensities estimated from amplitudes.

[2] Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NAG, BMA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#$\|Z\|$ >5	RMSZ	#$\|Z\|$ >5
1	A	0.28	0/6078	0.47	0/8266
1	C	0.27	0/6078	0.46	0/8266
1	E	0.26	0/6078	0.46	0/8266
1	G	0.32	0/6078	0.49	0/8266
2	B	0.25	0/1647	0.45	0/2249
2	D	0.25	0/1647	0.45	0/2249
2	F	0.26	0/1647	0.45	0/2249
2	H	0.25	0/1647	0.45	0/2249
All	All	0.28	0/30900	0.47	0/42060

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5906	0	5607	81	0
1	C	5906	0	5607	97	0
1	E	5906	0	5607	74	0
1	G	5906	0	5607	90	0
2	B	1608	0	1573	14	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	D	1608	0	1573	21	0
2	F	1608	0	1573	18	0
2	H	1608	0	1573	24	0
3	I	28	0	25	1	0
3	K	28	0	25	0	0
3	L	28	0	25	0	0
3	M	28	0	25	1	0
3	O	28	0	25	0	0
3	P	28	0	25	0	0
3	Q	28	0	25	1	0
3	R	28	0	25	1	0
3	T	28	0	25	0	0
3	U	28	0	25	0	0
3	V	28	0	25	1	0
3	X	28	0	25	0	0
3	Y	28	0	25	0	0
4	J	39	0	34	0	0
4	N	39	0	34	0	0
4	S	39	0	34	0	0
4	W	39	0	34	0	0
5	A	28	0	26	1	0
5	B	14	0	13	1	0
5	C	28	0	26	2	0
5	D	14	0	13	0	0
5	E	14	0	13	0	0
5	F	14	0	13	0	0
5	G	28	0	26	0	0
5	H	14	0	13	1	0
All	All	30730	0	29324	413	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

The worst 5 of 413 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:G:472:ILE:HD11	1:G:495:LEU:CD1	1.42	1.46
1:G:472:ILE:CD1	1:G:495:LEU:HD12	1.50	1.41
1:A:80:ASP:OD1	1:A:462:TYR:OH	1.59	1.19
1:E:233:ARG:HH21	1:E:253:PRO:HG3	1.20	1.05
1:E:233:ARG:NH2	1:E:253:PRO:HG3	1.78	0.97

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	722/731 (99%)	682 (94%)	39 (5%)	1 (0%)	51 83
1	C	722/731 (99%)	682 (94%)	40 (6%)	0	100 100
1	E	722/731 (99%)	688 (95%)	34 (5%)	0	100 100
1	G	722/731 (99%)	685 (95%)	37 (5%)	0	100 100
2	B	206/246 (84%)	197 (96%)	9 (4%)	0	100 100
2	D	206/246 (84%)	193 (94%)	13 (6%)	0	100 100
2	F	206/246 (84%)	199 (97%)	7 (3%)	0	100 100
2	H	206/246 (84%)	199 (97%)	7 (3%)	0	100 100
All	All	3712/3908 (95%)	3525 (95%)	186 (5%)	1 (0%)	100 100

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	97	GLY

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	641/649 (99%)	628 (98%)	13 (2%)	55 80

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	C	641/649 (99%)	619 (97%)	22 (3%)	37	69
1	E	641/649 (99%)	624 (97%)	17 (3%)	44	74
1	G	641/649 (99%)	623 (97%)	18 (3%)	43	73
2	B	190/221 (86%)	185 (97%)	5 (3%)	46	74
2	D	190/221 (86%)	186 (98%)	4 (2%)	53	79
2	F	190/221 (86%)	188 (99%)	2 (1%)	73	89
2	H	190/221 (86%)	183 (96%)	7 (4%)	34	66
All	All	3324/3480 (96%)	3236 (97%)	88 (3%)	46	74

5 of 88 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	E	433	ASN
1	G	380	CYS
1	E	467	CYS
1	G	188	ARG
1	G	473	PRO

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 14 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	425	ASN
1	E	72	ASN
2	H	408	ASN
1	G	500	GLN
1	G	726	GLN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

38 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
3	NAG	I	1	3,1	14,14,15	0.35	0	17,19,21	0.64	1 (5%)
3	NAG	I	2	3	14,14,15	0.36	0	17,19,21	0.39	0
4	NAG	J	1	4,1	14,14,15	0.56	1 (7%)	17,19,21	0.61	0
4	NAG	J	2	4	14,14,15	0.55	0	17,19,21	0.65	0
4	BMA	J	3	4	11,11,12	1.22	1 (9%)	15,15,17	1.16	1 (6%)
3	NAG	K	1	3,1	14,14,15	0.39	0	17,19,21	0.46	0
3	NAG	K	2	3	14,14,15	0.23	0	17,19,21	0.48	0
3	NAG	L	1	3,1	14,14,15	0.29	0	17,19,21	0.42	0
3	NAG	L	2	3	14,14,15	0.35	0	17,19,21	0.40	0
3	NAG	M	1	3,1	14,14,15	0.35	0	17,19,21	0.63	1 (5%)
3	NAG	M	2	3	14,14,15	0.34	0	17,19,21	0.36	0
4	NAG	N	1	4,1	14,14,15	0.45	0	17,19,21	0.63	0
4	NAG	N	2	4	14,14,15	0.51	0	17,19,21	0.67	0
4	BMA	N	3	4	11,11,12	1.25	1 (9%)	15,15,17	1.14	2 (13%)
3	NAG	O	1	3,1	14,14,15	0.38	0	17,19,21	0.57	0
3	NAG	O	2	3	14,14,15	0.21	0	17,19,21	0.51	0
3	NAG	P	1	3,1	14,14,15	0.32	0	17,19,21	0.47	0
3	NAG	P	2	3	14,14,15	0.22	0	17,19,21	0.44	0
3	NAG	Q	1	3,1	14,14,15	0.58	0	17,19,21	0.86	0
3	NAG	Q	2	3	14,14,15	0.48	0	17,19,21	0.52	0
3	NAG	R	1	3,1	14,14,15	0.38	0	17,19,21	0.66	1 (5%)
3	NAG	R	2	3	14,14,15	0.17	0	17,19,21	0.40	0
4	NAG	S	1	4,1	14,14,15	0.44	0	17,19,21	0.62	0
4	NAG	S	2	4	14,14,15	0.33	0	17,19,21	0.61	0
4	BMA	S	3	4	11,11,12	1.20	1 (9%)	15,15,17	1.20	2 (13%)
3	NAG	T	1	3,1	14,14,15	0.32	0	17,19,21	0.48	0
3	NAG	T	2	3	14,14,15	0.14	0	17,19,21	0.74	1 (5%)
3	NAG	U	1	3,1	14,14,15	0.48	0	17,19,21	0.51	0
3	NAG	U	2	3	14,14,15	0.16	0	17,19,21	0.51	0
3	NAG	V	1	3,1	14,14,15	0.34	0	17,19,21	0.65	1 (5%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\|>2$	Counts	RMSZ	$\#\|Z\|>2$
3	NAG	V	2	3	14,14,15	0.19	0	17,19,21	0.48	0
4	NAG	W	1	4,1	14,14,15	0.40	0	17,19,21	0.55	0
4	NAG	W	2	4	14,14,15	0.21	0	17,19,21	0.51	0
4	BMA	W	3	4	11,11,12	1.19	1 (9%)	15,15,17	1.27	3 (20%)
3	NAG	X	1	3,1	14,14,15	0.28	0	17,19,21	0.51	0
3	NAG	X	2	3	14,14,15	0.19	0	17,19,21	0.38	0
3	NAG	Y	1	3,1	14,14,15	0.37	0	17,19,21	0.42	0
3	NAG	Y	2	3	14,14,15	0.17	0	17,19,21	0.56	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	I	1	3,1	-	2/6/23/26	0/1/1/1
3	NAG	I	2	3	-	1/6/23/26	0/1/1/1
4	NAG	J	1	4,1	-	2/6/23/26	0/1/1/1
4	NAG	J	2	4	-	3/6/23/26	0/1/1/1
4	BMA	J	3	4	-	1/2/19/22	0/1/1/1
3	NAG	K	1	3,1	-	2/6/23/26	0/1/1/1
3	NAG	K	2	3	-	2/6/23/26	0/1/1/1
3	NAG	L	1	3,1	-	2/6/23/26	0/1/1/1
3	NAG	L	2	3	-	2/6/23/26	0/1/1/1
3	NAG	M	1	3,1	-	2/6/23/26	0/1/1/1
3	NAG	M	2	3	-	0/6/23/26	0/1/1/1
4	NAG	N	1	4,1	-	2/6/23/26	0/1/1/1
4	NAG	N	2	4	-	3/6/23/26	0/1/1/1
4	BMA	N	3	4	-	1/2/19/22	0/1/1/1
3	NAG	O	1	3,1	-	2/6/23/26	0/1/1/1
3	NAG	O	2	3	-	2/6/23/26	0/1/1/1
3	NAG	P	1	3,1	-	2/6/23/26	0/1/1/1
3	NAG	P	2	3	-	2/6/23/26	0/1/1/1
3	NAG	Q	1	3,1	-	4/6/23/26	0/1/1/1
3	NAG	Q	2	3	-	2/6/23/26	0/1/1/1
3	NAG	R	1	3,1	-	2/6/23/26	0/1/1/1
3	NAG	R	2	3	-	2/6/23/26	0/1/1/1
4	NAG	S	1	4,1	-	0/6/23/26	0/1/1/1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	S	2	4	-	2/6/23/26	0/1/1/1
4	BMA	S	3	4	-	1/2/19/22	0/1/1/1
3	NAG	T	1	3,1	-	0/6/23/26	0/1/1/1
3	NAG	T	2	3	-	2/6/23/26	0/1/1/1
3	NAG	U	1	3,1	-	2/6/23/26	0/1/1/1
3	NAG	U	2	3	-	2/6/23/26	0/1/1/1
3	NAG	V	1	3,1	-	2/6/23/26	0/1/1/1
3	NAG	V	2	3	-	2/6/23/26	0/1/1/1
4	NAG	W	1	4,1	-	0/6/23/26	0/1/1/1
4	NAG	W	2	4	-	3/6/23/26	0/1/1/1
4	BMA	W	3	4	-	2/2/19/22	0/1/1/1
3	NAG	X	1	3,1	-	2/6/23/26	0/1/1/1
3	NAG	X	2	3	-	2/6/23/26	0/1/1/1
3	NAG	Y	1	3,1	-	2/6/23/26	0/1/1/1
3	NAG	Y	2	3	-	2/6/23/26	0/1/1/1

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	N	3	BMA	C1-C2	3.57	1.60	1.52
4	J	3	BMA	C1-C2	3.35	1.59	1.52
4	S	3	BMA	C1-C2	2.50	1.57	1.52
4	W	3	BMA	C1-C2	2.47	1.57	1.52
4	J	1	NAG	O5-C1	-2.00	1.40	1.43

The worst 5 of 13 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
4	S	3	BMA	O2-C2-C3	-2.72	104.69	110.14
4	W	3	BMA	O2-C2-C3	-2.65	104.83	110.14
4	W	3	BMA	C1-C2-C3	-2.62	106.45	109.67
3	T	2	NAG	C1-O5-C5	2.56	115.67	112.19
3	R	1	NAG	C1-O5-C5	2.26	115.26	112.19

There are no chirality outliers.

5 of 69 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	K	1	NAG	O5-C5-C6-O6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
3	P	2	NAG	O5-C5-C6-O6
3	Q	1	NAG	C1-C2-N2-C7
3	O	2	NAG	O5-C5-C6-O6
3	X	1	NAG	O5-C5-C6-O6

There are no ring outliers.

6 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	R	1	NAG	1	0
3	M	1	NAG	1	0
3	I	1	NAG	1	0
3	V	1	NAG	1	0
3	I	2	NAG	1	0
3	Q	1	NAG	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.

Oligosaccharide Chain I



Bond lengths



Bond angles



Torsions



Rings

Oligosaccharide Chain K



Bond lengths



Bond angles



Torsions



Rings

Oligosaccharide Chain L



Bond lengths



Bond angles



Torsions



Rings

Oligosaccharide Chain M



Bond lengths



Bond angles



Torsions



Rings

Oligosaccharide Chain O



Bond lengths



Bond angles



Torsions



Rings

Oligosaccharide Chain P



Bond lengths



Bond angles



Torsions



Rings

Oligosaccharide Chain Q



Bond lengths



Bond angles



Torsions



Rings

Oligosaccharide Chain R



Bond lengths



Bond angles



Torsions



Rings

Oligosaccharide Chain T



Bond lengths



Bond angles



Torsions



Rings

Oligosaccharide Chain U



Bond lengths



Bond angles



Torsions



Rings

Oligosaccharide Chain V



Bond lengths



Bond angles



Torsions



Rings

Oligosaccharide Chain X



Bond lengths



Bond angles



Torsions



Rings

Oligosaccharide Chain Y



Bond lengths



Bond angles



Torsions



Rings

Oligosaccharide Chain J



Bond lengths



Bond angles



Torsions



Rings

Oligosaccharide Chain N



Bond lengths



Bond angles



Torsions



Rings

Oligosaccharide Chain S



Bond lengths



Bond angles



Torsions



Rings

Oligosaccharide Chain W



Bond lengths



Bond angles



Torsions



Rings

5.6 Ligand geometry (i)

11 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
5	NAG	D	701	2	14,14,15	0.19	0	17,19,21	0.42	0
5	NAG	H	701	2	14,14,15	0.22	0	17,19,21	0.37	0
5	NAG	B	701	2	14,14,15	0.20	0	17,19,21	0.49	0
5	NAG	G	801	1	14,14,15	0.17	0	17,19,21	1.02	1 (5%)
5	NAG	F	701	2	14,14,15	0.19	0	17,19,21	0.37	0
5	NAG	G	809	1	14,14,15	0.32	0	17,19,21	0.45	0
5	NAG	E	810	1	14,14,15	0.28	0	17,19,21	0.42	0
5	NAG	C	809	1	14,14,15	0.25	0	17,19,21	0.46	0
5	NAG	A	809	1	14,14,15	0.24	0	17,19,21	0.44	0
5	NAG	A	801	1	14,14,15	0.40	0	17,19,21	0.58	0
5	NAG	C	801	1	14,14,15	0.74	1 (7%)	17,19,21	1.38	3 (17%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	NAG	D	701	2	-	2/6/23/26	0/1/1/1
5	NAG	H	701	2	-	1/6/23/26	0/1/1/1
5	NAG	B	701	2	-	2/6/23/26	0/1/1/1
5	NAG	G	801	1	-	3/6/23/26	0/1/1/1
5	NAG	F	701	2	-	2/6/23/26	0/1/1/1
5	NAG	G	809	1	-	0/6/23/26	0/1/1/1
5	NAG	E	810	1	-	2/6/23/26	0/1/1/1
5	NAG	C	809	1	-	2/6/23/26	0/1/1/1
5	NAG	A	809	1	-	2/6/23/26	0/1/1/1
5	NAG	A	801	1	-	2/6/23/26	0/1/1/1
5	NAG	C	801	1	-	4/6/23/26	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	C	801	NAG	C1-C2	2.36	1.55	1.52

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
5	C	801	NAG	C2-N2-C7	4.25	128.96	122.90
5	G	801	NAG	C1-O5-C5	3.83	117.38	112.19
5	C	801	NAG	C1-O5-C5	2.57	115.67	112.19
5	C	801	NAG	C1-C2-N2	2.08	114.04	110.49

There are no chirality outliers.

5 of 22 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	D	701	NAG	C4-C5-C6-O6
5	B	701	NAG	C4-C5-C6-O6
5	A	809	NAG	O5-C5-C6-O6
5	C	809	NAG	O5-C5-C6-O6
5	D	701	NAG	O5-C5-C6-O6

There are no ring outliers.

5 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	H	701	NAG	1	0
5	B	701	NAG	1	0
5	C	809	NAG	1	0
5	A	809	NAG	1	0
5	C	801	NAG	1	0

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å^2)	Q<0.9
1	A	724/731 (99%)	-0.11	2 (0%)	94	88	28, 54, 90, 147	0
1	C	724/731 (99%)	-0.10	2 (0%)	94	88	30, 53, 89, 154	0
1	E	724/731 (99%)	-0.09	5 (0%)	87	75	31, 63, 105, 188	0
1	G	724/731 (99%)	-0.09	5 (0%)	87	75	29, 62, 105, 161	0
2	B	208/246 (84%)	-0.06	1 (0%)	91	81	35, 58, 94, 166	0
2	D	208/246 (84%)	-0.08	0	100	100	37, 58, 98, 173	0
2	F	208/246 (84%)	-0.07	3 (1%)	75	56	37, 60, 102, 219	0
2	H	208/246 (84%)	-0.10	1 (0%)	91	81	38, 62, 104, 196	0
All	All	3728/3908 (95%)	-0.09	19 (0%)	91	81	28, 58, 98, 219	0

The worst 5 of 19 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	F	579	THR	5.4
1	E	277	LEU	5.2
1	E	276	ASP	5.1
1	C	276	ASP	3.2
1	C	277	LEU	3.0

6.2 Non-standard residues in protein, DNA, RNA chains

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,

median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
3	NAG	Q	1	14/15	0.69	0.26	36,70,77,78	0
3	NAG	M	2	14/15	0.71	0.24	87,112,129,131	0
3	NAG	I	2	14/15	0.72	0.19	97,119,133,133	0
3	NAG	K	2	14/15	0.72	0.29	120,135,140,146	0
4	BMA	S	3	11/12	0.72	0.20	78,88,93,94	0
3	NAG	O	2	14/15	0.75	0.28	108,126,133,137	0
3	NAG	T	2	14/15	0.76	0.21	104,122,132,135	0
4	BMA	N	3	11/12	0.78	0.29	71,81,88,92	0
4	BMA	J	3	11/12	0.78	0.25	70,82,89,89	0
3	NAG	R	2	14/15	0.79	0.22	65,93,105,107	0
4	BMA	W	3	11/12	0.79	0.21	70,74,83,83	0
3	NAG	L	2	14/15	0.81	0.19	80,91,95,96	0
3	NAG	U	2	14/15	0.84	0.15	93,100,106,109	0
3	NAG	P	2	14/15	0.86	0.18	75,86,92,94	0
3	NAG	X	2	14/15	0.86	0.19	101,115,126,127	0
3	NAG	V	2	14/15	0.87	0.18	72,99,108,109	0
3	NAG	Y	2	14/15	0.89	0.18	89,97,105,109	0
3	NAG	M	1	14/15	0.92	0.15	70,83,88,99	0
3	NAG	R	1	14/15	0.92	0.20	91,96,105,109	0
3	NAG	Q	2	14/15	0.93	0.21	35,47,65,78	0
3	NAG	V	1	14/15	0.93	0.18	84,96,107,114	0
3	NAG	K	1	14/15	0.93	0.17	71,80,98,109	0
4	NAG	N	2	14/15	0.94	0.19	55,63,75,76	0
3	NAG	I	1	14/15	0.94	0.14	73,84,91,102	0
4	NAG	J	1	14/15	0.94	0.18	43,49,58,60	0
3	NAG	L	1	14/15	0.94	0.15	65,75,83,85	0
4	NAG	S	1	14/15	0.95	0.21	59,63,67,68	0
4	NAG	S	2	14/15	0.95	0.15	51,68,77,80	0
3	NAG	T	1	14/15	0.95	0.11	60,74,85,91	0
4	NAG	W	2	14/15	0.95	0.15	52,70,79,80	0
3	NAG	X	1	14/15	0.95	0.14	59,71,80,85	0
4	NAG	N	1	14/15	0.96	0.17	38,45,54,57	0
3	NAG	P	1	14/15	0.96	0.16	59,72,80,81	0
4	NAG	W	1	14/15	0.96	0.19	53,60,63,65	0
4	NAG	J	2	14/15	0.96	0.15	51,59,69,71	0
3	NAG	O	1	14/15	0.96	0.22	62,76,89,102	0
3	NAG	Y	1	14/15	0.97	0.12	69,76,80,86	0
3	NAG	U	1	14/15	0.98	0.15	67,73,80,89	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around Chain I:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)







Electron density around Chain K:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)







Electron density around Chain L:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)







Electron density around Chain M:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)







Electron density around Chain O:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)







Electron density around Chain P:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)







Electron density around Chain Q:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)



Electron density around Chain R:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)







Electron density around Chain T:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)







Electron density around Chain U:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)







Electron density around Chain V:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)







Electron density around Chain X:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)



Electron density around Chain Y:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)



Electron density around Chain J:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)



Electron density around Chain N:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)



Electron density around Chain S:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)







Electron density around Chain W:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)







6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
5	NAG	G	809	14/15	0.61	0.23	77,113,119,120	0
5	NAG	E	810	14/15	0.71	0.23	114,123,133,137	0
5	NAG	C	809	14/15	0.71	0.26	105,110,113,118	0
5	NAG	G	801	14/15	0.73	0.27	39,69,83,85	0
5	NAG	A	801	14/15	0.75	0.20	81,92,99,99	0
5	NAG	A	809	14/15	0.75	0.19	104,112,117,119	0
5	NAG	D	701	14/15	0.78	0.23	90,110,115,123	0
5	NAG	B	701	14/15	0.81	0.29	93,101,106,113	0
5	NAG	C	801	14/15	0.84	0.25	54,83,97,98	0
5	NAG	H	701	14/15	0.85	0.17	82,89,94,99	0
5	NAG	F	701	14/15	0.86	0.19	69,84,89,90	0

6.5 Other polymers (i)

There are no such residues in this entry.