

wwPDB X-ray Structure Validation Summary Report (i)

Jun 24, 2024 – 06:15 PM EDT

PDB ID	:	6PA5
Title	:	ECAII(T89V,K162T) MUTANT IN COMPLEX WITH L-ASN AT PH 8.3 IN
		SPACE GROUP P2(1)
Authors	:	Lubkowski, J.; Wlodawer, A.
Deposited on	:	2019-06-11
Resolution	:	2.00 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:


MolProbity Mogul	:	4.02b-467 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)		
EDS	:	2.37.1
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.37.1

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $X\text{-}RAY \, DIFFRACTION$

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$\begin{array}{c} \textbf{Whole archive} \\ (\#\textbf{Entries}) \end{array}$	${f Similar\ resolution}\ (\#{ m Entries,\ resolution\ range}({ m \AA}))$
R_{free}	130704	8085 (2.00-2.00)
Clashscore	141614	9178 (2.00-2.00)
Ramachandran outliers	138981	9054 (2.00-2.00)
Sidechain outliers	138945	9053 (2.00-2.00)
RSRZ outliers	127900	7900 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain	
1	А	334	86%	11% ••
1	В	334	88%	9% ••
1	С	334	84%	9% • 6%
1	D	334	85%	8% 6%
1	Е	334	86%	7% • 6%



Mol	Chain	Length	Quality of chain	
1	F	334	84%	10% • 6%
1	G	334	87%	10% ••
1	Н	334	86%	10% ••



2 Entry composition (i)

There are 5 unique types of molecules in this entry. The entry contains 21351 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

Mol	Chain	Residues		Ate	oms			ZeroOcc	AltConf	Trace			
1	1 1	326	Total	С	Ν	0	S	0	0	0			
	A	320	2429	1516	414	491	8	0	0	0			
1	В	326	Total	С	Ν	0	S	0	1	0			
1	D	520	2434	1519	415	492	8	0	L	0			
1	С	314	Total	С	Ν	Ο	S	0	0	0			
1	U	514	2346	1467	400	471	8	0	0	U			
1	D	314	Total	С	Ν	Ο	\mathbf{S}	0	2	0			
1	D	514	2354	1473	400	472	9			0			
1	Е	313	Total	С	Ν	Ο	\mathbf{S}	0	0	0			
L	Ľ	515	2342	1465	399	470	8		0	0			
1	F	313	Total	\mathbf{C}	Ν	Ο	\mathbf{S}	0	1	0			
	Ľ	T,	T,	T,		2347	1468	399	472	8	0	1	U
1	G	326	Total	\mathbf{C}	Ν	Ο	\mathbf{S}	0	1	0			
	I G	320	2432	1518	414	492	8	0	1	0			
1	1 H	326	Total	\mathbf{C}	Ν	Ο	\mathbf{S}	0	0	0			
		520	2429	1516	414	491	8			0			

• Molecule 1 is a protein called L-asparaginase 2.

There are 80 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Actual Comment	
А	-7	MET	-	initiating methionine	UNP P00805
А	-6	ASP	-	expression tag	UNP P00805
А	-5	HIS	-	expression tag	UNP P00805
А	-4	HIS	-	expression tag	UNP P00805
А	-3	HIS	-	expression tag	UNP P00805
А	-2	HIS	-	expression tag	UNP P00805
А	-1	HIS	-	expression tag	UNP P00805
А	0	HIS	-	expression tag	UNP P00805
А	89	VAL	THR	engineered mutation	UNP P00805
А	162	THR	LYS	engineered mutation	UNP P00805
В	-7	MET	-	initiating methionine	UNP P00805
В	-6	ASP	-	expression tag	UNP P00805
В	-5	HIS	-	expression tag	UNP P00805



F

F

F

-5

-4

-3

HIS

HIS

HIS

-

-

-

Chain	Residue	Modelled	Actual	Comment	Reference
В	-4	HIS	-	expression tag	UNP P00805
В	-3	HIS	-	expression tag	UNP P00805
В	-2	HIS	-	expression tag	UNP P00805
В	-1	HIS	-	expression tag	UNP P00805
В	0	HIS	-	expression tag	UNP P00805
В	89	VAL	THR	engineered mutation	UNP P00805
В	162	THR	LYS	engineered mutation	UNP P00805
С	-7	MET	-	initiating methionine	UNP P00805
С	-6	ASP	-	expression tag	UNP P00805
С	-5	HIS	_	expression tag	UNP P00805
С	-4	HIS	-	expression tag	UNP P00805
С	-3	HIS	-	expression tag	UNP P00805
С	-2	HIS	-	expression tag	UNP P00805
С	-1	HIS	-	expression tag	UNP P00805
С	0	HIS	-	expression tag	UNP P00805
С	89	VAL	THR	engineered mutation	UNP P00805
С	162	THR	LYS	engineered mutation	UNP P00805
D	-7	MET	-	initiating methionine	UNP P00805
D	-6	ASP	-	expression tag	UNP P00805
D	-5	HIS	-	expression tag	UNP P00805
D	-4	HIS	-	expression tag	UNP P00805
D	-3	HIS	-	expression tag	UNP P00805
D	-2	HIS	-	expression tag	UNP P00805
D	-1	HIS	-	expression tag	UNP P00805
D	0	HIS	-	expression tag	UNP P00805
D	89	VAL	THR	engineered mutation	UNP P00805
D	162	THR	LYS	engineered mutation	UNP P00805
Е	-7	MET	-	initiating methionine	UNP P00805
Е	-6	ASP	-	expression tag	UNP P00805
Е	-5	HIS	-	expression tag	UNP P00805
Е	-4	HIS	-	expression tag	UNP P00805
Е	-3	HIS	-	expression tag	UNP P00805
Е	-2	HIS	-	expression tag	UNP P00805
Е	-1	HIS	-	expression tag	UNP P00805
Е	0	HIS	-	expression tag	UNP P00805
Е	89	VAL	THR	engineered mutation	UNP P00805
Е	162	THR	LYS	engineered mutation	UNP P00805
F	-7	MET	_	initiating methionine	UNP P00805
F	-6	ASP	-	expression tag	UNP P00805
·		-			

Continued from previous page...

Continued on next page...

UNP P00805

UNP P00805

UNP P00805



expression tag


expression tag

expression tag

Chain	Residue	Modelled	Actual	Comment	Reference
F	-2	HIS	-	expression tag	UNP P00805
F	-1	HIS	-	expression tag	UNP P00805
F	0	HIS	-	expression tag	UNP P00805
F	89	VAL	THR	engineered mutation	UNP P00805
F	162	THR	LYS	engineered mutation	UNP P00805
G	-7	MET	-	initiating methionine	UNP P00805
G	-6	ASP	-	expression tag	UNP P00805
G	-5	HIS	-	expression tag	UNP P00805
G	-4	HIS	-	expression tag	UNP P00805
G	-3	HIS	-	expression tag	UNP P00805
G	-2	HIS	-	expression tag	UNP P00805
G	-1	HIS	-	expression tag	UNP P00805
G	0	HIS	-	expression tag	UNP P00805
G	89	VAL	THR	engineered mutation	UNP P00805
G	162	THR	LYS	engineered mutation	UNP P00805
Н	-7	MET	-	initiating methionine	UNP P00805
Н	-6	ASP	-	expression tag	UNP P00805
Н	-5	HIS	-	expression tag	UNP P00805
Н	-4	HIS	-	expression tag	UNP P00805
Н	-3	HIS	-	expression tag	UNP P00805
Н	-2	HIS	-	expression tag	UNP P00805
Н	-1	HIS	-	expression tag	UNP P00805
Н	0	HIS	-	expression tag	UNP P00805
Н	89	VAL	THR	engineered mutation	UNP P00805
Н	162	THR	LYS	engineered mutation	UNP P00805

• Molecule 2 is ASPARAGINE (three-letter code: ASN) (formula: $C_4H_8N_2O_3$) (labeled as "Ligand of Interest" by depositor).






Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	А	1	$\begin{array}{ccccc} \text{Total} & \text{C} & \text{N} & \text{O} \\ 9 & 4 & 2 & 3 \end{array}$	0	0
2	В	1	$\begin{array}{ccccc} \text{Total} & \text{C} & \text{N} & \text{O} \\ 9 & 4 & 2 & 3 \end{array}$	0	0
2	С	1	$\begin{array}{cccc} \text{Total} & \text{C} & \text{N} & \text{O} \\ 9 & 4 & 2 & 3 \end{array}$	0	0
2	D	1	$\begin{array}{ccccc} \text{Total} & \text{C} & \text{N} & \text{O} \\ 9 & 4 & 2 & 3 \end{array}$	0	0
2	Ε	1	$\begin{array}{ccccc} \text{Total} & \text{C} & \text{N} & \text{O} \\ 9 & 4 & 2 & 3 \end{array}$	0	0
2	F	1	$\begin{array}{cccc} \text{Total} & \text{C} & \text{N} & \text{O} \\ 9 & 4 & 2 & 3 \end{array}$	0	0
2	G	1	$\begin{array}{ccccc} \text{Total} & \text{C} & \text{N} & \text{O} \\ 9 & 4 & 2 & 3 \end{array}$	0	0
2	Н	1	$\begin{array}{ccccc} \text{Total} & \text{C} & \text{N} & \text{O} \\ 9 & 4 & 2 & 3 \end{array}$	0	0

 $\bullet\,$ Molecule 3 is IMIDAZOLE (three-letter code: IMD) (formula: C_3H_5N_2).






Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	А	1	$\begin{array}{ccc} \text{Total} & \text{C} & \text{N} \\ 5 & 3 & 2 \end{array}$	0	0
3	В	1	$\begin{array}{ccc} \text{Total} \text{C} \text{N} \\ 5 3 2 \end{array}$	0	0
3	С	1	$\begin{array}{ccc} \text{Total} \text{C} \text{N} \\ 5 3 2 \end{array}$	0	0
3	D	1	$\begin{array}{ccc} \text{Total} \text{C} \text{N} \\ 5 3 2 \end{array}$	0	0
3	Е	1	$\begin{array}{ccc} \text{Total} & \text{C} & \text{N} \\ 5 & 3 & 2 \end{array}$	0	0
3	F	1	$\begin{array}{ccc} \text{Total} & \text{C} & \text{N} \\ 5 & 3 & 2 \end{array}$	0	0
3	G	1	$\begin{array}{ccc} \text{Total} & \text{C} & \text{N} \\ 5 & 3 & 2 \end{array}$	0	0

• Molecule 4 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).





Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	А	1	$\begin{array}{ccc} \text{Total} \text{C} \text{O} \\ 6 3 3 \end{array}$	0	0
4	В	1	$\begin{array}{ccc} \text{Total} \text{C} \text{O} \\ 6 3 3 \end{array}$	0	0
4	D	1	$\begin{array}{ccc} \text{Total} \text{C} \text{O} \\ 6 3 3 \end{array}$	0	0


• Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	А	306	Total O 307 307	0	1
5	В	307	Total O 307 307	0	1
5	С	301	Total O 303 303	0	2
5	D	291	Total O 293 293	0	3
5	Е	209	Total O 210 210	0	1
5	F	211	Total O 211 211	0	0
5	G	240	Total O 240 240	0	0
5	Н	242	Total O 242 242	0	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.




• Molecule 1: L-asparaginase 2





• Molecule 1: L-asparaginase 2





4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants	140.53Å 62.34Å 151.05Å	Deperitor
a, b, c, α , β , γ	90.00° 117.68° 90.00°	Depositor
Resolution (Å)	26.21 - 2.00	Depositor
Resolution (A)	$26.21 \ - \ 2.00$	EDS
% Data completeness	95.5 (26.21-2.00)	Depositor
(in resolution range)	99.6 (26.21-2.00)	EDS
R _{merge}	0.07	Depositor
R _{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	2.62 (at 1.99 Å)	Xtriage
Refinement program	REFMAC 5.8.0158	Depositor
D D	0.170 , 0.214	Depositor
R, R_{free}	0.205 , 0.244	DCC
R_{free} test set	4189 reflections (2.67%)	wwPDB-VP
Wilson B-factor $(Å^2)$	26.2	Xtriage
Anisotropy	0.050	Xtriage
Bulk solvent $k_{sol}(e/Å^3), B_{sol}(Å^2)$	0.32, 46.9	EDS
L-test for twinning ²	$< L > = 0.40, < L^2 > = 0.24$	Xtriage
Estimated twinning fraction	0.000 for h,-k,-h-l	Xtriage
Dependent of twinning fraction	0.736 for H, K, L	Denesiten
Reported twinning fraction	0.264 for -H, -K, H+L	Depositor
Outliers	8 of 156927 reflections (0.005%)	Xtriage
F_o, F_c correlation	0.98	EDS
Total number of atoms	21351	wwPDB-VP
Average B, all atoms $(Å^2)$	32.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The analyses of the Patterson function reveals a significant off-origin peak that is 99.10 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 3.6338e-12. The detected translational NCS is most likely also responsible for the elevated intensity ratio.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for a centric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: IMD, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bo	nd lengths	В	ond angles
	Unam	RMSZ	# Z > 5	RMSZ	# Z > 5
1	А	1.05	2/2466~(0.1%)	1.07	14/3360~(0.4%)
1	В	1.07	1/2474~(0.0%)	1.07	8/3371~(0.2%)
1	С	1.01	0/2381	1.06	14/3243~(0.4%)
1	D	0.98	0/2395	1.03	12/3261~(0.4%)
1	Ε	0.93	0/2377	1.00	7/3238~(0.2%)
1	F	0.94	0/2385	1.00	9/3249~(0.3%)
1	G	0.96	0/2472	1.01	9/3368~(0.3%)
1	Н	0.96	0/2466	1.00	6/3360~(0.2%)
All	All	0.99	3/19416~(0.0%)	1.03	79/26450~(0.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	В	0	1

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Ζ	$\operatorname{Observed}(\operatorname{\AA})$	Ideal(Å)
1	В	252	SER	CB-OG	-8.41	1.31	1.42
1	А	252	SER	CB-OG	-7.95	1.31	1.42
1	А	1	LEU	N-CA	5.33	1.57	1.46

The worst 5 of 79 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
1	Е	144	ARG	NE-CZ-NH1	10.46	125.53	120.30
1	F	144	ARG	NE-CZ-NH1	9.43	125.01	120.30



Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
1	В	204	ASP	CB-CG-OD2	-9.08	110.12	118.30
1	Н	167	ASP	CB-CG-OD1	8.21	125.69	118.30
1	F	144	ARG	NE-CZ-NH2	-7.67	116.46	120.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	В	26	THR	Peptide

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	А	2429	0	2420	16	0
1	В	2434	0	2426	11	0
1	С	2346	0	2343	14	0
1	D	2354	0	2357	11	0
1	Е	2342	0	2340	7	0
1	F	2347	0	2344	11	0
1	G	2432	0	2425	13	0
1	Н	2429	0	2420	14	0
2	А	9	0	5	0	0
2	В	9	0	5	0	0
2	С	9	0	5	0	0
2	D	9	0	5	0	0
2	Е	9	0	5	0	0
2	F	9	0	5	0	0
2	G	9	0	5	0	0
2	Н	9	0	5	0	0
3	А	5	0	5	0	0
3	В	5	0	5	0	0
3	С	5	0	5	0	0
3	D	5	0	5	0	0
3	Е	5	0	5	0	0
3	F	5	0	5	0	0
3	G	5	0	5	0	0



Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	А	6	0	8	0	0
4	В	6	0	8	0	0
4	D	6	0	8	0	0
5	А	307	0	0	2	1
5	В	307	0	0	0	0
5	С	303	0	0	2	2
5	D	293	0	0	4	0
5	Ε	210	0	0	1	2
5	F	211	0	0	2	0
5	G	240	0	0	3	1
5	Н	242	0	0	0	0
All	All	21351	0	19174	91	3

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

The worst 5 of 91 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:G:188:ASP:O	5:G:501:HOH:O	1.59	1.20
1:D:41:GLN:HE21	1:D:41:GLN:H	1.18	0.91
5:E:509:HOH:O	1:H:21:THR:HG21	1.73	0.89
1:C:41:GLN:HE21	1:C:41:GLN:H	1.18	0.87
5:F:503:HOH:O	1:G:21:THR:HG21	1.73	0.86

All (3) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:A:745:HOH:O	5:C:758:HOH:O[1_545]	1.78	0.42
5:C:788:HOH:O	5:E:697:HOH:O[2_756]	1.82	0.38
5:E:689:HOH:O	5:G:619:HOH:O[1_545]	2.14	0.06

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries



of similar resolution.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	А	324/334~(97%)	319~(98%)	5 (2%)	0	100 100
1	В	325/334~(97%)	319~(98%)	6 (2%)	0	100 100
1	С	310/334~(93%)	305~(98%)	5 (2%)	0	100 100
1	D	312/334~(93%)	306~(98%)	6 (2%)	0	100 100
1	Ε	309/334~(92%)	302~(98%)	6 (2%)	1 (0%)	41 37
1	F	310/334~(93%)	301~(97%)	8 (3%)	1 (0%)	41 37
1	G	325/334~(97%)	319~(98%)	6 (2%)	0	100 100
1	Н	324/334~(97%)	318~(98%)	6 (2%)	0	100 100
All	All	2539/2672~(95%)	2489 (98%)	48 (2%)	2(0%)	51 49

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	Е	198	THR
1	F	198	THR

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent side chain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Perce	ntiles
1	А	266/274~(97%)	258~(97%)	8~(3%)	41	41
1	В	267/274~(97%)	260 (97%)	7 (3%)	46	48
1	С	257/274~(94%)	246 (96%)	11 (4%)	29	26
1	D	259/274~(94%)	247~(95%)	12~(5%)	27	23
1	Ε	257/274~(94%)	248~(96%)	9~(4%)	36	35
1	F	258/274~(94%)	247~(96%)	11 (4%)	29	26
1	G	267/274~(97%)	255~(96%)	12~(4%)	27	24



Continued from previous page...

Mol	Chain	Analysed	Analysed Rotameric Outlie		Percentiles	
1	Н	266/274~(97%)	252~(95%)	14~(5%)	22 18	
All	All	2097/2192~(96%)	2013 (96%)	84 (4%)	31 29	

 $5~{\rm of}~84$ residues with a non-rotameric side chain are listed below:

Mol	Chain	Res	Type
1	F	314	LYS
1	Н	29	LYS
1	G	29	LYS
1	G	211	LEU
1	Н	103	VAL

Sometimes side chains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 22 such side chains are listed below:

Mol	Chain	Res	Type
1	F	197	HIS
1	G	318	GLN
1	G	197	HIS
1	Н	64	ASN
1	С	41	GLN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

18 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and



the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Туре	Chain	Res	Link	В	ond leng	gths	B	ond ang	gles
WIOI	Type	Ullalli	nes	LIIIK	Counts	RMSZ	# Z >2	Counts	RMSZ	# Z >2
2	ASN	А	401	-	$6,\!8,\!8$	1.31	1 (16%)	8,10,10	0.47	0
2	ASN	D	401	-	$6,\!8,\!8$	1.12	1 (16%)	$8,\!10,\!10$	1.20	2 (25%)
3	IMD	D	402	-	$3,\!5,\!5$	0.68	0	$4,\!5,\!5$	0.78	0
2	ASN	В	401	-	6,8,8	0.74	0	8,10,10	1.53	2 (25%)
3	IMD	В	402	-	$3,\!5,\!5$	0.26	0	$4,\!5,\!5$	0.86	0
4	GOL	А	403	-	$5,\!5,\!5$	0.66	0	$5,\!5,\!5$	0.72	0
4	GOL	D	403	-	$5,\!5,\!5$	1.01	0	$5,\!5,\!5$	1.17	0
3	IMD	А	402	-	$3,\!5,\!5$	0.24	0	$4,\!5,\!5$	1.28	0
2	ASN	Е	401	-	$6,\!8,\!8$	0.91	0	8,10,10	0.80	0
2	ASN	С	401	-	6,8,8	0.82	0	8,10,10	1.14	1 (12%)
2	ASN	Н	401	-	6,8,8	0.79	0	8,10,10	1.02	1 (12%)
3	IMD	F	402	-	$3,\!5,\!5$	0.27	0	$4,\!5,\!5$	0.71	0
3	IMD	Е	402	-	$3,\!5,\!5$	0.27	0	4,5,5	0.63	0
3	IMD	G	402	-	$3,\!5,\!5$	0.34	0	4,5,5	0.42	0
4	GOL	В	403	-	$5,\!5,\!5$	0.51	0	$5,\!5,\!5$	0.62	0
3	IMD	С	402	-	$3,\!5,\!5$	0.23	0	4,5,5	0.78	0
2	ASN	F	401	-	6,8,8	0.85	0	8,10,10	1.19	1 (12%)
2	ASN	G	401	-	6,8,8	1.01	1 (16%)	8,10,10	0.76	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	ASN	А	401	-	-	4/8/8/8	-
2	ASN	D	401	-	-	2/8/8/8	-
4	GOL	D	403	-	-	2/4/4/4	-
2	ASN	В	401	-	-	2/8/8/8	-
4	GOL	А	403	-	-	4/4/4/4	-
3	IMD	В	402	-	-	-	0/1/1/1
3	IMD	D	402	-	-	-	0/1/1/1
3	IMD	А	402	-	-	-	0/1/1/1
2	ASN	Е	401	-	-	4/8/8/8	-



Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	ASN	С	401	-	-	4/8/8/8	-
2	ASN	Н	401	-	-	2/8/8/8	-
3	IMD	F	402	-	-	-	0/1/1/1
3	IMD	Е	402	-	-	-	0/1/1/1
4	GOL	В	403	-	-	4/4/4/4	-
3	IMD	G	402	-	-	-	0/1/1/1
3	IMD	С	402	-	-	-	0/1/1/1
2	ASN	F	401	-	-	2/8/8/8	-
2	ASN	G	401	-	-	4/8/8/8	-

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Ζ	Observed(Å)	Ideal(Å)
2	D	401	ASN	OXT-C	-2.33	1.22	1.30
2	G	401	ASN	OXT-C	-2.16	1.23	1.30
2	А	401	ASN	OXT-C	-2.02	1.23	1.30

The worst 5 of 7 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$Observed(^{o})$	$Ideal(^{o})$
2	В	401	ASN	OXT-C-CA	3.25	124.45	113.38
2	F	401	ASN	OXT-C-CA	2.52	121.95	113.38
2	С	401	ASN	OXT-C-O	-2.48	118.45	124.09
2	D	401	ASN	OXT-C-CA	2.42	121.64	113.38
2	Н	401	ASN	OXT-C-CA	2.21	120.89	113.38

There are no chirality outliers.

5 of 34 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	А	401	ASN	O-C-CA-N
2	В	401	ASN	O-C-CA-N
2	С	401	ASN	O-C-CA-N
2	D	401	ASN	O-C-CA-N
2	Е	401	ASN	O-C-CA-N


There are no ring outliers.

No monomer is involved in short contacts.


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In



addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.











6PA5














5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ $>$	#RSRZ>2	$OWAB(Å^2)$	$\mathbf{Q}{<}0.9$
1	А	326/334~(97%)	-0.63	1 (0%) 94 93	19, 26, 44, 72	0
1	В	326/334~(97%)	-0.63	1 (0%) 94 93	20, 26, 43, 70	0
1	С	314/334~(94%)	-0.65	0 100 100	19, 25, 42, 75	0
1	D	314/334~(94%)	-0.65	0 100 100	20, 25, 42, 74	0
1	Е	313/334~(93%)	-0.36	2 (0%) 89 88	24, 34, 53, 78	0
1	F	313/334~(93%)	-0.37	1 (0%) 94 93	24, 34, 52, 78	0
1	G	326/334~(97%)	-0.56	0 100 100	24, 31, 49, 77	0
1	Н	326/334~(97%)	-0.56	0 100 100	23, 32, 47, 76	0
All	All	2558/2672~(95%)	-0.55	5 (0%) 95 94	19, 29, 49, 78	0

All (5) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	А	28	GLY	3.3
1	В	28	GLY	3.2
1	F	78	ASP	2.4
1	Е	78	ASP	2.2
1	Е	106	ASP	2.1

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.




6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	GOL	D	403	6/6	0.32	0.37	63,89,94,101	0
3	IMD	G	402	5/5	0.73	0.23	67,72,74,82	0
4	GOL	А	403	6/6	0.74	0.21	$54,\!55,\!59,\!67$	0
4	GOL	В	403	6/6	0.84	0.19	$54,\!57,\!61,\!69$	0
3	IMD	А	402	5/5	0.84	0.18	45,48,50,52	0
3	IMD	В	402	5/5	0.87	0.19	43,46,47,48	0
3	IMD	Е	402	5/5	0.91	0.22	59,59,61,65	0
3	IMD	F	402	5/5	0.91	0.22	58,60,61,64	0
3	IMD	С	402	5/5	0.94	0.23	52,53,53,56	0
2	ASN	F	401	9/9	0.95	0.07	26,29,31,31	0
2	ASN	Е	401	9/9	0.95	0.08	26,28,30,31	0
3	IMD	D	402	5/5	0.95	0.22	22,23,23,24	5
2	ASN	С	401	9/9	0.97	0.06	20,22,24,25	0
2	ASN	D	401	9/9	0.97	0.06	20,21,24,24	0
2	ASN	G	401	9/9	0.98	0.06	26,27,30,30	0
2	ASN	Н	401	9/9	0.98	0.06	24,27,30,30	0
2	ASN	А	401	9/9	0.98	0.07	21,24,27,28	0
2	ASN	В	401	9/9	0.98	0.06	20,24,28,28	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.







































6.5 Other polymers (i)

There are no such residues in this entry.

