



# wwPDB EM Validation Summary Report ⓘ

May 13, 2024 – 07:20 pm BST

PDB ID : 6TUT  
EMDB ID : EMD-10595  
Title : Cryo-EM structure of the RNA Polymerase III-Maf1 complex  
Authors : Vorlaender, M.K.; Hagen, W.J.H.; Mueller, C.W.  
Deposited on : 2020-01-08  
Resolution : 3.25 Å (reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev92  
MolProbity : 4.02b-467  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
MapQ : 1.9.13  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36.2


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*ELECTRON MICROSCOPY*

The reported resolution of this entry is 3.25 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)
Clashscore	158937	4297
Ramachandran outliers	154571	4023
Sidechain outliers	154315	3826

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion  $< 40\%$ ). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	1460	
2	B	1149	
3	C	335	
4	D	161	
5	E	215	
6	F	155	
7	G	212	
8	H	146	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
9	I	110	
10	J	70	
11	K	142	
12	L	70	
13	M	282	
14	N	422	
15	O	654	
16	P	317	
17	Q	251	
18	R	267	

## 2 Entry composition [i](#)

There are 19 unique types of molecules in this entry. The entry contains 40324 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DNA-directed RNA polymerase III subunit RPC1.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
1	A	1390	10908	6880	1926	2046	56	0	0

- Molecule 2 is a protein called DNA-directed RNA polymerase III subunit RPC2.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
2	B	1097	8657	5481	1493	1623	60	0	0

- Molecule 3 is a protein called DNA-directed RNA polymerases I and III subunit RPAC1.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
3	C	334	2647	1676	453	510	8	0	0

- Molecule 4 is a protein called DNA-directed RNA polymerase III subunit RPC9.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
4	D	104	857	553	138	161	5	0	0

- Molecule 5 is a protein called DNA-directed RNA polymerases I, II, and III subunit RPABC1.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
5	E	215	1759	1116	310	321	12	0	0

- Molecule 6 is a protein called DNA-directed RNA polymerases I, II, and III subunit RPABC2.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
6	F	83	671	429	114	125	3	0	0

- Molecule 7 is a protein called DNA-directed RNA polymerase III subunit RPC8.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
7	G	199	Total	C	N	O	S	0	0
			1594	1038	258	291	7		

- Molecule 8 is a protein called DNA-directed RNA polymerases I, II, and III subunit RPABC3.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
8	H	146	Total	C	N	O	S	0	0
			1161	726	195	235	5		

- Molecule 9 is a protein called DNA-directed RNA polymerase III subunit RPC10.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
9	I	43	Total	C	N	O	S	0	0
			332	210	51	65	6		

- Molecule 10 is a protein called DNA-directed RNA polymerases I, II, and III subunit RPABC5.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
10	J	67	Total	C	N	O	S	0	0
			549	350	95	98	6		

- Molecule 11 is a protein called DNA-directed RNA polymerases I and III subunit RPAC2.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
11	K	102	Total	C	N	O	S	0	0
			801	501	131	164	5		

- Molecule 12 is a protein called DNA-directed RNA polymerases I, II, and III subunit RPABC4.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
12	L	45	Total	C	N	O	S	0	0
			358	221	71	62	4		

- Molecule 13 is a protein called DNA-directed RNA polymerase III subunit RPC5.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
13	M	166	Total	C	N	O	S	0	0
			1345	862	225	257	1		

- Molecule 14 is a protein called DNA-directed RNA polymerase III subunit RPC4.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
14	N	125	968	610	173	181	4	0	0

- Molecule 15 is a protein called DNA-directed RNA polymerase III subunit RPC3.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
15	O	551	4433	2819	763	833	18	0	0

- Molecule 16 is a protein called DNA-directed RNA polymerase III subunit RPC6.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
16	P	196	1599	1036	258	299	6	0	0

- Molecule 17 is a protein called DNA-directed RNA polymerase III subunit RPC7.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
17	Q	79	503	319	89	94	1	0	0

- Molecule 18 is a protein called Repressor of RNA polymerase III transcription MAF1, Repressor of RNA polymerase III transcription MAF1.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
18	R	142	1175	760	189	221	5	0	0

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
R	-2	GLY	-	expression tag	UNP P41910
R	-1	ALA	-	expression tag	UNP P41910
R	0	MET	-	expression tag	UNP P41910
R	1	GLY	-	expression tag	UNP P41910

- Molecule 19 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		AltConf
			Total	Zn	
19	A	2	2	2	0

*Continued on next page...*


*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Residues</b>	<b>Atoms</b>		<b>AltConf</b>
19	B	1	Total 1	Zn 1	0
19	I	1	Total 1	Zn 1	0
19	J	2	Total 2	Zn 2	0
19	L	1	Total 1	Zn 1	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



- Molecule 1: DNA-directed RNA polymerase III subunit RPC1










• Molecule 2: DNA-directed RNA polymerase III subunit RPC2




• Molecule 3: DNA-directed RNA polymerases I and III subunit RPAC1




• Molecule 4: DNA-directed RNA polymerase III subunit RPC9




• Molecule 5: DNA-directed RNA polymerases I, II, and III subunit RPABC1




• Molecule 6: DNA-directed RNA polymerases I, II, and III subunit RPABC2





• Molecule 7: DNA-directed RNA polymerase III subunit RPC8







- Molecule 8: DNA-directed RNA polymerases I, II, and III subunit RPABC3




- Molecule 9: DNA-directed RNA polymerase III subunit RPC10




- Molecule 10: DNA-directed RNA polymerases I, II, and III subunit RPABC5



- Molecule 11: DNA-directed RNA polymerases I and III subunit RPAC2




- Molecule 12: DNA-directed RNA polymerases I, II, and III subunit RPABC4




- Molecule 13: DNA-directed RNA polymerase III subunit RPC5








• Molecule 16: DNA-directed RNA polymerase III subunit RPC6



• Molecule 17: DNA-directed RNA polymerase III subunit RPC7





## 4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, C1	Depositor
Number of particles used	117442	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ( $e^-/\text{\AA}^2$ )	60.5	Depositor
Minimum defocus (nm)	750	Depositor
Maximum defocus (nm)	3500	Depositor
Magnification	130000	Depositor
Image detector	GATAN K2 SUMMIT (4k x 4k)	Depositor
Maximum map value	0.419	Depositor
Minimum map value	-0.239	Depositor
Average map value	0.000	Depositor
Map value standard deviation	0.010	Depositor
Recommended contour level	0.075	Depositor
Map size (Å)	374.76, 374.76, 374.76	wwPDB
Map dimensions	360, 360, 360	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	1.041, 1.041, 1.041	Depositor

## 5 Model quality i

### 5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section:  
ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.48	0/11101	0.64	5/14993 (0.0%)
2	B	0.53	1/8808 (0.0%)	0.67	5/11879 (0.0%)
3	C	0.53	0/2703	0.68	4/3666 (0.1%)
4	D	0.28	0/870	0.50	0/1167
5	E	0.38	0/1795	0.60	1/2416 (0.0%)
6	F	0.55	1/683 (0.1%)	0.62	0/923
7	G	0.35	0/1634	0.54	0/2217
8	H	0.56	0/1181	0.70	2/1602 (0.1%)
9	I	0.40	0/339	0.81	2/459 (0.4%)
10	J	0.62	0/558	0.70	1/750 (0.1%)
11	K	0.54	0/812	0.66	1/1096 (0.1%)
12	L	0.47	0/360	0.80	2/478 (0.4%)
13	M	0.35	0/1377	0.57	0/1863
14	N	0.32	0/978	0.62	1/1310 (0.1%)
15	O	0.33	0/4501	0.60	1/6072 (0.0%)
16	P	0.34	0/1632	0.56	0/2206
17	Q	0.35	0/510	0.60	1/695 (0.1%)
18	R	0.34	0/1204	0.57	0/1633
All	All	0.46	2/41046 (0.0%)	0.64	26/55425 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	4
2	B	0	3
3	C	0	1
13	M	0	1
All	All	0	9



All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	489	LEU	C-N	-6.44	1.19	1.34
6	F	137	TYR	C-N	-5.06	1.22	1.34

The worst 5 of 26 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	638	ASP	CB-CG-OD1	8.38	125.84	118.30
12	L	50	ASP	CB-CG-OD1	7.67	125.20	118.30
1	A	576	LEU	CA-CB-CG	7.22	131.90	115.30
3	C	287	ASP	CB-CG-OD1	7.03	124.63	118.30
5	E	181	ALA	C-N-CA	6.73	138.52	121.70

There are no chirality outliers.

5 of 9 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	1043	ASP	Peptide
1	A	1178	LYS	Peptide
1	A	477	GLN	Peptide
1	A	932	PRO	Peptide
2	B	223	SER	Peptide

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	10908	0	11035	124	0
2	B	8657	0	8786	102	0
3	C	2647	0	2616	28	0
4	D	857	0	868	12	0
5	E	1759	0	1788	25	0
6	F	671	0	691	10	0
7	G	1594	0	1580	20	0
8	H	1161	0	1124	13	0
9	I	332	0	316	3	0
10	J	549	0	559	6	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
11	K	801	0	795	11	0
12	L	358	0	383	5	0
13	M	1345	0	1307	16	0
14	N	968	0	1014	14	0
15	O	4433	0	4600	51	0
16	P	1599	0	1579	15	0
17	Q	503	0	395	7	0
18	R	1175	0	1143	16	0
19	A	2	0	0	0	0
19	B	1	0	0	0	0
19	I	1	0	0	0	0
19	J	2	0	0	0	0
19	L	1	0	0	0	0
All	All	40324	0	40579	414	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 414 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
15:O:530:GLU:HG3	15:O:534:ARG:HE	1.56	0.71
1:A:1185:GLN:HE21	1:A:1229:ARG:HD3	1.55	0.69
1:A:1316:THR:HG22	1:A:1318:HIS:H	1.59	0.68
1:A:483:LEU:HB3	1:A:507:PRO:HB3	1.77	0.66
1:A:497:THR:HG23	1:A:499:ARG:HD2	1.77	0.66

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	1378/1460 (94%)	1269 (92%)	107 (8%)	2 (0%)	51	82
2	B	1093/1149 (95%)	1007 (92%)	86 (8%)	0	100	100
3	C	332/335 (99%)	306 (92%)	26 (8%)	0	100	100
4	D	98/161 (61%)	96 (98%)	2 (2%)	0	100	100
5	E	213/215 (99%)	197 (92%)	16 (8%)	0	100	100
6	F	81/155 (52%)	77 (95%)	4 (5%)	0	100	100
7	G	195/212 (92%)	181 (93%)	14 (7%)	0	100	100
8	H	144/146 (99%)	125 (87%)	19 (13%)	0	100	100
9	I	41/110 (37%)	31 (76%)	10 (24%)	0	100	100
10	J	65/70 (93%)	60 (92%)	5 (8%)	0	100	100
11	K	100/142 (70%)	98 (98%)	2 (2%)	0	100	100
12	L	43/70 (61%)	35 (81%)	8 (19%)	0	100	100
13	M	162/282 (57%)	151 (93%)	11 (7%)	0	100	100
14	N	121/422 (29%)	106 (88%)	15 (12%)	0	100	100
15	O	547/654 (84%)	513 (94%)	34 (6%)	0	100	100
16	P	188/317 (59%)	164 (87%)	24 (13%)	0	100	100
17	Q	73/251 (29%)	60 (82%)	10 (14%)	3 (4%)	3	17
18	R	138/267 (52%)	126 (91%)	12 (9%)	0	100	100
All	All	5012/6418 (78%)	4602 (92%)	405 (8%)	5 (0%)	54	82

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
17	Q	1080	ASP
1	A	642	PRO
17	Q	1084	ASN
17	Q	49	ASN
1	A	439	ASP

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	1206/1257 (96%)	1188 (98%)	18 (2%)	65	80
2	B	959/1006 (95%)	952 (99%)	7 (1%)	84	90
3	C	295/296 (100%)	290 (98%)	5 (2%)	60	78
4	D	97/145 (67%)	95 (98%)	2 (2%)	53	75
5	E	197/197 (100%)	195 (99%)	2 (1%)	76	85
6	F	73/137 (53%)	72 (99%)	1 (1%)	67	81
7	G	173/190 (91%)	173 (100%)	0	100	100
8	H	128/128 (100%)	128 (100%)	0	100	100
9	I	39/98 (40%)	39 (100%)	0	100	100
10	J	62/65 (95%)	61 (98%)	1 (2%)	62	79
11	K	92/130 (71%)	91 (99%)	1 (1%)	73	84
12	L	40/57 (70%)	40 (100%)	0	100	100
13	M	142/249 (57%)	142 (100%)	0	100	100
14	N	105/360 (29%)	102 (97%)	3 (3%)	42	68
15	O	506/593 (85%)	498 (98%)	8 (2%)	62	79
16	P	179/285 (63%)	176 (98%)	3 (2%)	60	78
17	Q	35/212 (16%)	34 (97%)	1 (3%)	42	68
18	R	134/243 (55%)	131 (98%)	3 (2%)	52	74
All	All	4462/5648 (79%)	4407 (99%)	55 (1%)	72	83

5 of 55 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
3	C	165	ARG
10	J	42	LYS
18	R	342	ARG
16	P	189	ASN
3	C	277	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 60 such sidechains are listed below:

Mol	Chain	Res	Type
2	B	1036	HIS
15	O	544	ASN
5	E	61	GLN
15	O	514	ASN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
18	R	17	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 7 ligands modelled in this entry, 7 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

The following chains have linkage breaks:

Mol	Chain	Number of breaks
17	Q	2
2	B	1

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	Q	75:MET	C	1076:SER	N	22.36
1	Q	1096:LEU	C	1104:ASP	N	12.78
1	B	489:LEU	C	490:GLN	N	1.19


## 6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-10595. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

### 6.1 Orthogonal projections [i](#)


#### 6.1.1 Primary map



X




Y




Z


#### 6.1.2 Raw map



X



Y




Z


The images above show the map projected in three orthogonal directions.

## 6.2 Central slices [i](#)


### 6.2.1 Primary map



X Index: 180




Y Index: 180




Z Index: 180


### 6.2.2 Raw map



X Index: 180



Y Index: 180




Z Index: 180

The images above show central slices of the map in three orthogonal directions.




## 6.3 Largest variance slices [i](#)


### 6.3.1 Primary map



X Index: 183




Y Index: 170




Z Index: 135


### 6.3.2 Raw map



X Index: 174



Y Index: 170




Z Index: 195


The images above show the largest variance slices of the map in three orthogonal directions.

## 6.4 Orthogonal standard-deviation projections (False-color) [i](#)


### 6.4.1 Primary map



X




Y




Z


### 6.4.2 Raw map



X



Y



Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.


## 6.5 Orthogonal surface views [i](#)

### 6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.075. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

### 6.5.2 Raw map



These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.


## 6.6 Mask visualisation [i](#)

This section was not generated. No masks/segmentation were deposited.

## 7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

### 7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


## 7.2 Volume estimate [i](#)



The volume at the recommended contour level is 202 nm<sup>3</sup>; this corresponds to an approximate mass of 182 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

### 7.3 Rotationally averaged power spectrum [i](#)




\*Reported resolution corresponds to spatial frequency of 0.308 Å<sup>-1</sup>

## 8 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

### 8.1 FSC [i](#)



\*Reported resolution corresponds to spatial frequency of 0.308 Å<sup>-1</sup>

## 8.2 Resolution estimates [i](#)

Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	3.25	-	-
Author-provided FSC curve	3.25	3.79	3.32
Unmasked-calculated*	3.85	6.91	3.95


\*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 3.85 differs from the reported value 3.25 by more than 10 %



## 9 Map-model fit [i](#)


This section contains information regarding the fit between EMDB map EMD-10595 and PDB model 6TUT. Per-residue inclusion information can be found in section 3 on page 8.

### 9.1 Map-model overlay [i](#)




The images above show the 3D surface view of the map at the recommended contour level 0.075 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

## 9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

## 9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.075).







































## 9.4 Atom inclusion [i](#)



At the recommended contour level, 81% of all backbone atoms, 69% of all non-hydrogen atoms, are inside the map.

## 9.5 Map-model fit summary

The table lists the average atom inclusion at the recommended contour level (0.075) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	 0.6860	 0.4900
A	 0.7660	 0.5280
B	 0.7990	 0.5480
C	 0.8440	 0.5590
D	 0.2940	 0.3260
E	 0.6280	 0.4320
F	 0.8530	 0.5610
G	 0.4770	 0.3950
H	 0.7740	 0.5360
I	 0.5610	 0.4210
J	 0.8800	 0.5730
K	 0.8710	 0.5860
L	 0.7510	 0.5170
M	 0.5030	 0.4300
N	 0.5010	 0.4390
O	 0.5510	 0.4070
P	 0.3280	 0.3190
Q	 0.5770	 0.4160
R	 0.4480	 0.3990

