



# Full wwPDB X-ray Structure Validation Report ⓘ

Jun 24, 2024 – 07:50 AM EDT

PDB ID : 7ATI  
Title : Crystal structure of dimeric chlorite dismutase variant Q74V (CCld Q74V)  
from *Cyanothece* sp. PCC7425  
Authors : Schmidt, D.; Mlynek, G.; Djinovic-Carugo, K.; Obinger, C.  
Deposited on : 2020-10-30  
Resolution : 1.51 Å (reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 2022.3.0, CSD as543be (2022)  
Xtriage (Phenix) : 1.20.1  
EDS : 2.37.1  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.37.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.51 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	4009 (1.54-1.50)
Clashscore	141614	4249 (1.54-1.50)
Ramachandran outliers	138981	4148 (1.54-1.50)
Sidechain outliers	138945	4146 (1.54-1.50)
RSRZ outliers	127900	3943 (1.54-1.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	188	
1	B	188	

## 2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 6751 atoms, of which 3144 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Chlorite dismutase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
1	A	182	3058	980	1524	275	275	4	0	10	0
1	B	182	3053	979	1522	274	273	5	0	8	0

There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-5	GLY	-	expression tag	UNP B8HNS6
A	-4	PRO	-	expression tag	UNP B8HNS6
A	-3	GLY	-	expression tag	UNP B8HNS6
A	-2	TYR	-	expression tag	UNP B8HNS6
A	-1	GLN	-	expression tag	UNP B8HNS6
A	0	ASP	-	expression tag	UNP B8HNS6
A	1	PRO	-	expression tag	UNP B8HNS6
A	74	VAL	GLN	engineered mutation	UNP B8HNS6
B	-5	GLY	-	expression tag	UNP B8HNS6
B	-4	PRO	-	expression tag	UNP B8HNS6
B	-3	GLY	-	expression tag	UNP B8HNS6
B	-2	TYR	-	expression tag	UNP B8HNS6
B	-1	GLN	-	expression tag	UNP B8HNS6
B	0	ASP	-	expression tag	UNP B8HNS6
B	1	PRO	-	expression tag	UNP B8HNS6
B	74	VAL	GLN	engineered mutation	UNP B8HNS6

- Molecule 2 is GLYCEROL (three-letter code: GOL) (formula: C<sub>3</sub>H<sub>8</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	H	O		
2	A	1	9	3	3	3	0	0
2	A	1	9	3	3	3	0	0
2	A	1	9	3	3	3	0	0
2	B	1	9	3	3	3	0	0
2	B	1	9	3	3	3	0	0
2	B	1	9	3	3	3	0	0

- Molecule 3 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (formula:  $C_{34}H_{32}FeN_4O_4$ ) (labeled as "Ligand of Interest" by depositor).




- Molecule 5 is water.

<b>Mol</b>	<b>Chain</b>	<b>Residues</b>	<b>Atoms</b>		<b>ZeroOcc</b>	<b>AltConf</b>
5	A	220	Total 220	O 220	0	0
5	B	186	Total 186	O 186	0	0



### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Chlorite dismutase



- Molecule 1: Chlorite dismutase



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	55.01Å 72.68Å 60.17Å 90.00° 112.73° 90.00°	Depositor
Resolution (Å)	36.34 – 1.51 36.34 – 1.51	Depositor EDS
% Data completeness (in resolution range)	98.3 (36.34-1.51) 98.3 (36.34-1.51)	Depositor EDS
$R_{merge}$	0.09	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.01 (at 1.51Å)	Xtrriage
Refinement program	PHENIX dev_3885	Depositor
R, $R_{free}$	0.175 , 0.193 0.177 , 0.195	Depositor DCC
$R_{free}$ test set	1575 reflections (2.34%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	25.3	Xtrriage
Anisotropy	0.275	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.41 , 46.5	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.98	EDS
Total number of atoms	6751	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	36.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.60% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, HEM, PEG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	1.22	6/1604 (0.4%)	1.46	18/2171 (0.8%)
1	B	0.74	1/1608 (0.1%)	0.85	8/2176 (0.4%)
All	All	1.01	7/3212 (0.2%)	1.20	26/4347 (0.6%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2
1	B	0	1
All	All	0	3

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	168	TRP	CG-CD1	-27.43	0.98	1.36
1	A	168	TRP	CG-CD2	12.73	1.65	1.43
1	A	150	GLU	CD-OE1	12.50	1.39	1.25
1	A	150	GLU	CG-CD	11.91	1.69	1.51
1	B	168	TRP	CG-CD1	6.95	1.46	1.36
1	A	168	TRP	CD2-CE2	-6.77	1.33	1.41
1	A	150	GLU	CA-CB	5.06	1.65	1.53

All (26) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	168	TRP	CD1-CG-CD2	-33.82	79.24	106.30
1	A	168	TRP	N-CA-CB	18.78	144.40	110.60

*Continued on next page...*

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	168	TRP	CD1-NE1-CE2	-15.40	95.14	109.00
1	A	21	ARG	NE-CZ-NH1	-14.78	112.91	120.30
1	A	168	TRP	CB-CG-CD2	-14.48	107.78	126.60
1	A	150	GLU	OE1-CD-OE2	-13.70	106.87	123.30
1	A	168	TRP	CA-CB-CG	13.36	139.07	113.70
1	B	168	TRP	CB-CG-CD1	-12.32	110.98	127.00
1	A	168	TRP	CB-CA-C	-10.83	88.75	110.40
1	B	168	TRP	CD1-CG-CD2	-9.67	98.56	106.30
1	A	168	TRP	NE1-CE2-CD2	-8.69	98.61	107.30
1	A	168	TRP	CB-CG-CD1	8.57	138.14	127.00
1	A	150	GLU	CA-CB-CG	-8.27	95.20	113.40
1	A	21	ARG	CD-NE-CZ	7.96	134.74	123.60
1	B	168	TRP	CA-CB-CG	7.82	128.56	113.70
1	A	51	ARG	NE-CZ-NH1	-7.42	116.59	120.30
1	B	21	ARG	CG-CD-NE	7.25	127.03	111.80
1	A	21	ARG	CG-CD-NE	-7.19	96.71	111.80
1	A	167	GLU	C-N-CA	-7.17	103.78	121.70
1	A	150	GLU	N-CA-CB	6.35	122.02	110.60
1	A	51	ARG	NE-CZ-NH2	6.33	123.46	120.30
1	B	21	ARG	NE-CZ-NH2	-5.72	117.44	120.30
1	B	166	LYS	CD-CE-NZ	-5.62	98.77	111.70
1	B	168	TRP	N-CA-C	5.55	126.00	111.00
1	B	21	ARG	CD-NE-CZ	5.17	130.83	123.60
1	A	51	ARG	CD-NE-CZ	5.06	130.68	123.60

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	150	GLU	Sidechain
1	A	21	ARG	Sidechain
1	B	21	ARG	Sidechain

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1534	1524	1484	13	0
1	B	1531	1522	1481	12	0
2	A	18	9	24	0	0
2	B	18	9	23	1	0
3	A	43	30	30	0	0
3	B	43	30	30	1	0
4	A	7	10	10	1	0
4	B	7	10	10	1	0
5	A	220	0	0	0	2
5	B	186	0	0	1	1
All	All	3607	3144	3092	25	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (25) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:163:ARG:HA	1:A:168:TRP:HD1	1.53	0.73
1:A:41:VAL:O	1:A:51:ARG:NH2	2.28	0.66
1:A:70:LEU:HD23	1:A:119:LEU:CD2	2.29	0.63
1:A:70:LEU:HD23	1:A:119:LEU:HD21	1.84	0.59
1:A:137:GLU:OE1	4:A:204:PEG:O4	2.22	0.55
1:B:44:ILE:HG21	1:B:51:ARG:HB2	1.89	0.53
1:A:163:ARG:HA	1:A:168:TRP:CD1	2.41	0.52
1:A:168:TRP:HA	1:A:168:TRP:CE3	2.45	0.51
1:B:37:LEU:HD12	1:B:41:VAL:HG21	1.94	0.50
1:A:168:TRP:HA	1:A:168:TRP:HE3	1.77	0.48
1:B:22:ASN:ND2	1:B:27:GLY:HA3	2.29	0.48
1:B:138:GLU:HG3	4:B:204:PEG:H12	1.96	0.47
1:B:158:LEU:O	1:B:162[A]:MET:HG3	2.14	0.47
1:A:148:ALA:CB	1:A:150:GLU:OE1	2.62	0.47
1:A:122:LEU:N	1:A:123:PRO:CD	2.79	0.46
1:B:44:ILE:HD12	1:B:48:SER:HB2	1.99	0.45
1:B:86:VAL:HG22	1:B:178:TRP:CE3	2.52	0.45
1:B:70:LEU:HD23	1:B:119:LEU:CD2	2.47	0.44
1:A:148:ALA:HB1	1:A:150:GLU:OE1	2.18	0.44
1:A:20:ILE:HD12	1:A:30:LEU:HA	2.01	0.42
1:B:19:LYS:HE2	2:B:205:GOL:O2	2.20	0.41
1:A:11:ARG:HD2	1:A:32:GLU:OE1	2.20	0.41
1:B:145:PHE:CZ	3:B:202:HEM:C2C	3.09	0.40
1:B:151:HIS:HE1	5:B:315:HOH:O	2.03	0.40

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:A:306:HOH:O	5:A:470:HOH:O[2_545]	2.00	0.20
5:A:482:HOH:O	5:B:446:HOH:O[1_454]	2.19	0.01

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	190/188 (101%)	186 (98%)	4 (2%)	0	100	100
1	B	188/188 (100%)	181 (96%)	7 (4%)	0	100	100
All	All	378/376 (100%)	367 (97%)	11 (3%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	162/159 (102%)	162 (100%)	0	100	100
1	B	163/159 (102%)	163 (100%)	0	100	100
All	All	325/318 (102%)	325 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such

sidechains are listed below:

Mol	Chain	Res	Type
1	B	16	GLN
1	B	22	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

10 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	GOL	B	201	-	5,5,5	0.49	0	5,5,5	0.92	0
4	PEG	B	204	-	6,6,6	0.56	0	5,5,5	0.33	0
3	HEM	B	202	1,5	42,50,50	1.79	14 (33%)	46,82,82	2.05	14 (30%)
2	GOL	B	203	-	5,5,5	1.36	1 (20%)	5,5,5	1.07	0
2	GOL	A	203	-	5,5,5	0.99	0	5,5,5	1.14	0
4	PEG	A	204	-	6,6,6	0.43	0	5,5,5	0.46	0
3	HEM	A	202	1,5	42,50,50	1.51	7 (16%)	46,82,82	1.93	15 (32%)
2	GOL	B	205	-	5,5,5	0.89	0	5,5,5	0.96	0
2	GOL	A	201	-	5,5,5	0.53	0	5,5,5	1.25	1 (20%)
2	GOL	A	205	-	5,5,5	0.65	0	5,5,5	1.20	1 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GOL	B	201	-	-	0/4/4/4	-
4	PEG	B	204	-	-	4/4/4/4	-
3	HEM	B	202	1,5	-	4/12/54/54	-
2	GOL	B	203	-	-	4/4/4/4	-
2	GOL	A	203	-	-	2/4/4/4	-
4	PEG	A	204	-	-	1/4/4/4	-
3	HEM	A	202	1,5	-	4/12/54/54	-
2	GOL	B	205	-	-	2/4/4/4	-
2	GOL	A	201	-	-	2/4/4/4	-
2	GOL	A	205	-	-	2/4/4/4	-

All (22) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	202	HEM	C3C-C2C	-4.10	1.34	1.40
3	B	202	HEM	C1A-NA	3.72	1.43	1.36
3	A	202	HEM	C3C-C4C	3.54	1.46	1.41
3	A	202	HEM	C1A-NA	3.49	1.43	1.36
3	B	202	HEM	C3C-C4C	3.19	1.46	1.41
3	B	202	HEM	C4D-ND	-2.80	1.35	1.40
3	A	202	HEM	CAB-C3B	2.60	1.54	1.47
3	B	202	HEM	O1D-CGD	2.53	1.30	1.22
3	B	202	HEM	CAA-C2A	2.50	1.58	1.52
3	A	202	HEM	O2D-CGD	-2.41	1.22	1.30
2	B	203	GOL	O2-C2	-2.39	1.36	1.43
3	B	202	HEM	C1A-CHA	-2.35	1.34	1.41
3	A	202	HEM	CMD-C2D	2.34	1.55	1.50
3	B	202	HEM	FE-ND	2.29	2.10	1.98
3	A	202	HEM	CBA-CGA	2.28	1.55	1.50
3	B	202	HEM	CMD-C2D	2.24	1.55	1.50
3	B	202	HEM	CBA-CGA	2.18	1.55	1.50
3	B	202	HEM	C3C-CAC	2.18	1.52	1.47
3	A	202	HEM	CAA-C2A	2.16	1.57	1.52
3	B	202	HEM	C3B-C4B	-2.13	1.40	1.44
3	B	202	HEM	CAB-C3B	2.09	1.53	1.47
3	B	202	HEM	C2C-C1C	2.01	1.47	1.42

All (31) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	202	HEM	C4D-ND-C1D	4.90	111.01	105.21
3	B	202	HEM	C4D-ND-C1D	4.30	110.31	105.21
3	B	202	HEM	CAA-CBA-CGA	-4.04	102.96	113.83
3	A	202	HEM	CAA-CBA-CGA	-3.93	103.23	113.83
3	A	202	HEM	C3B-C4B-NB	-3.73	106.79	109.47
3	B	202	HEM	C4B-C3B-C2B	3.71	110.69	107.28
3	B	202	HEM	CHD-C1D-ND	3.67	128.39	124.44
3	B	202	HEM	CMA-C3A-C4A	-3.56	123.24	128.46
3	A	202	HEM	C1B-NB-C4B	3.42	109.25	105.21
3	A	202	HEM	CHD-C1D-ND	3.35	128.04	124.44
3	B	202	HEM	CHB-C1B-NB	3.29	128.45	124.37
3	B	202	HEM	C3B-C4B-NB	-3.29	107.10	109.47
3	A	202	HEM	C2D-C1D-ND	-3.11	106.31	109.90
3	A	202	HEM	CHB-C1B-NB	3.06	128.17	124.37
3	B	202	HEM	O1A-CGA-CBA	-3.05	113.42	123.09
3	A	202	HEM	CMA-C3A-C4A	-3.00	124.07	128.46
3	B	202	HEM	C2D-C1D-ND	-2.87	106.59	109.90
3	B	202	HEM	CBA-CAA-C2A	-2.66	108.07	112.54
3	B	202	HEM	C3D-C4D-ND	-2.54	107.38	110.17
3	A	202	HEM	C4C-CHD-C1D	2.53	125.90	122.56
3	B	202	HEM	C4C-CHD-C1D	2.49	125.85	122.56
3	A	202	HEM	O1A-CGA-CBA	-2.42	115.41	123.09
3	A	202	HEM	C3D-C4D-ND	-2.37	107.58	110.17
3	B	202	HEM	CHC-C4B-C3B	2.34	128.16	124.57
2	A	201	GOL	C3-C2-C1	-2.33	103.25	111.80
3	A	202	HEM	C4B-C3B-C2B	2.21	109.31	107.28
3	A	202	HEM	CHC-C4B-NB	2.11	126.71	124.44
3	A	202	HEM	CBB-CAB-C3B	-2.09	117.07	127.53
3	A	202	HEM	O1D-CGD-CBD	-2.09	116.47	123.09
2	A	205	GOL	C3-C2-C1	-2.09	104.14	111.80
3	B	202	HEM	O2A-CGA-CBA	2.02	120.38	114.00

There are no chirality outliers.

All (25) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	203	GOL	O1-C1-C2-C3
2	B	203	GOL	O1-C1-C2-C3
2	B	203	GOL	C1-C2-C3-O3
2	B	205	GOL	O1-C1-C2-O2
2	B	205	GOL	O1-C1-C2-C3

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms
2	A	203	GOL	O1-C1-C2-O2
4	A	204	PEG	O1-C1-C2-O2
2	A	201	GOL	O1-C1-C2-C3
2	A	205	GOL	C1-C2-C3-O3
2	A	201	GOL	O1-C1-C2-O2
2	A	205	GOL	O2-C2-C3-O3
2	B	203	GOL	O1-C1-C2-O2
2	B	203	GOL	O2-C2-C3-O3
4	B	204	PEG	O1-C1-C2-O2
4	B	204	PEG	C1-C2-O2-C3
4	B	204	PEG	O2-C3-C4-O4
4	B	204	PEG	C4-C3-O2-C2
3	A	202	HEM	CAA-CBA-CGA-O1A
3	B	202	HEM	CAD-CBD-CGD-O1D
3	B	202	HEM	CAA-CBA-CGA-O1A
3	A	202	HEM	CAD-CBD-CGD-O1D
3	A	202	HEM	CAA-CBA-CGA-O2A
3	B	202	HEM	CAD-CBD-CGD-O2D
3	B	202	HEM	CAA-CBA-CGA-O2A
3	A	202	HEM	CAD-CBD-CGD-O2D


There are no ring outliers.


4 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	B	204	PEG	1	0
3	B	202	HEM	1	0
4	A	204	PEG	1	0
2	B	205	GOL	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	182/188 (96%)	0.05	4 (2%) 62 67	17, 28, 43, 72	0
1	B	182/188 (96%)	0.59	16 (8%) 10 10	21, 33, 59, 93	0
All	All	364/376 (96%)	0.32	20 (5%) 25 27	17, 31, 53, 93	0

All (20) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	42	ALA	11.9
1	B	44	ILE	10.8
1	B	46	LEU	9.0
1	B	41	VAL	6.2
1	B	43	GLU	5.6
1	A	1	PRO	5.6
1	A	46	LEU	5.6
1	B	1	PRO	5.5
1	B	11	ARG	4.2
1	B	45	PRO	4.1
1	B	47	ASP	4.1
1	A	168	TRP	3.9
1	B	30	LEU	3.4
1	B	48	SER	3.0
1	A	47	ASP	3.0
1	B	72	ALA	2.9
1	B	95[A]	GLN	2.7
1	B	18	VAL	2.6
1	B	88	ILE	2.2
1	B	94	ALA	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)


There are no monosaccharides in this entry.


## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	GOL	A	203	6/6	0.69	0.38	45,52,66,70	0
4	PEG	B	204	7/7	0.70	0.23	50,60,71,71	0
2	GOL	A	205	6/6	0.76	0.25	43,47,59,63	0
4	PEG	A	204	7/7	0.81	0.38	40,54,67,67	0
2	GOL	B	205	6/6	0.84	0.14	47,53,63,69	0
2	GOL	B	203	6/6	0.88	0.11	33,45,55,66	0
2	GOL	B	201	6/6	0.90	0.17	36,44,52,55	0
2	GOL	A	201	6/6	0.92	0.10	32,43,46,46	0
3	HEM	B	202	43/43	0.98	0.11	20,24,30,34	0
3	HEM	A	202	43/43	0.99	0.10	18,23,29,33	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





## 6.5 Other polymers [i](#)

There are no such residues in this entry.