



Full wwPDB X-ray Structure Validation Report ⓘ

Aug 23, 2022 – 10:17 pm BST

PDB ID : 7BFZ
Title : X-ray structure of human prostate-specific membrane antigen(PSMA) in complex with a inhibitor Glu-490
Authors : Rakhimbekova, A.; Motlova, L.; Barinka, C.
Deposited on : 2021-01-05
Resolution : 1.73 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at <http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.4, CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.30
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0267
CCP4 : 7.1.010 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.30


1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.73 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3764 (1.76-1.72)
Clashscore	141614	3923 (1.76-1.72)
Ramachandran outliers	138981	3878 (1.76-1.72)
Sidechain outliers	138945	3878 (1.76-1.72)
RSRZ outliers	127900	3705 (1.76-1.72)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	707	 7% 91% 7%
2	B	3	 100%
2	F	3	 67% 33%
3	C	5	 100%
4	D	2	 100%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
5	E	4	 25% 75%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	BMA	B	3	-	-	-	X
4	NAG	D	2	-	-	-	X
5	BMA	E	3	-	-	-	X
5	FUC	E	4	-	-	-	X

2 Entry composition i

There are 14 unique types of molecules in this entry. The entry contains 6622 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Glutamate carboxypeptidase 2.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	694	5836	3753	968	1093	22	0	63	0

- Molecule 2 is an oligosaccharide called beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
2	B	3	39	22	2	15	0	0	0
2	F	3	39	22	2	15	0	0	0

- Molecule 3 is an oligosaccharide called alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
3	C	5	61	34	2	25	0	0	0

- Molecule 4 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
4	D	2	28	16	2	10	0	0	0

- Molecule 5 is an oligosaccharide called beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
5	E	4	49	28	2	19	0	0	0

- Molecule 6 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Zn		
6	A	2	2	2	0	0

- Molecule 7 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Ca		
7	A	1	1	1	0	0

- Molecule 8 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Cl		
8	A	1	1	1	0	0

- Molecule 9 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: C₈H₁₅NO₆).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	
9	A	1	Total	C	N	O	0	0
			14	8	1	5		
9	A	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 10 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: $C_2H_6O_2$).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
10	A	1	Total	C	O	0	0
			4	2	2		
10	A	1	Total	C	O	0	0
			4	2	2		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
10	A	1	Total	C	O	0	0
			4	2	2		

- Molecule 11 is DI(HYDROXYETHYL)ETHER (three-letter code: PEG) (formula: $C_4H_{10}O_3$).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
11	A	1	Total	C	O	0	0
			7	4	3		

- Molecule 12 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
12	A	2	Total	Na	0	0
			2	2		

- Molecule 13 is (((S)-1-carboxy-5-((E)-2-cyano-3-(5-(1-(3-methoxy-3-oxopropyl)-1,2,3,4-tetrahydroquinolin-6-yl)thiophen-2-yl)acrylamido)pentyl)carbamoyl)-L-glutamic acid (three-letter code: TKZ) (formula: $C_{33}H_{39}N_5O_{10}S$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	S		
13	A	1	49	33	5	10	1	0	0


- Molecule 14 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
14	A	467	Total	O	0	1
			468	468		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Glutamate carboxypeptidase 2




- Molecule 2: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 2: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 3: alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose




- Molecule 4: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain D:  100%

MAG1
MAG2

- Molecule 5: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose

Chain E:  25% 75%

MAG1
MAG2
BMA3
FUC4

4 Data and refinement statistics

Property	Value	Source
Space group	I 2 2 2	Depositor
Cell constants a, b, c, α , β , γ	101.54Å 130.10Å 158.91Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	46.96 – 1.73 46.96 – 1.73	Depositor EDS
% Data completeness (in resolution range)	99.5 (46.96-1.73) 99.5 (46.96-1.73)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.75 (at 1.73Å)	Xtriage
Refinement program	REFMAC 5.8.0135	Depositor
R, R_{free}	0.163 , 0.183 0.171 , 0.187	Depositor DCC
R_{free} test set	2101 reflections (1.93%)	wwPDB-VP
Wilson B-factor (Å ²)	23.9	Xtriage
Anisotropy	0.576	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	(Not available) , (Not available)	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	6622	wwPDB-VP
Average B, all atoms (Å ²)	36.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.19% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CL, BMA, CA, FUC, NAG, ZN, PEG, EDO, MAN, NA, TKZ

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.91	2/6177 (0.0%)	0.85	4/8361 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	434	TRP	CE3-CZ3	5.43	1.47	1.38
1	A	460	TYR	CG-CD1	5.11	1.45	1.39

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	465	ASP	CB-CG-OD1	6.33	123.99	118.30
1	A	673	ARG	NE-CZ-NH1	5.33	122.97	120.30
1	A	668[A]	LEU	CB-CG-CD1	5.16	119.77	111.00
1	A	668[B]	LEU	CB-CG-CD1	5.16	119.77	111.00

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	546	PHE	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5836	0	5757	28	0
2	B	39	0	34	0	0
2	F	39	0	34	0	0
3	C	61	0	52	0	0
4	D	28	0	25	0	0
5	E	49	0	43	0	0
6	A	2	0	0	0	0
7	A	1	0	0	0	0
8	A	1	0	0	0	0
9	A	28	0	26	1	0
10	A	12	0	18	0	0
11	A	7	0	10	0	0
12	A	2	0	0	0	0
13	A	49	0	0	1	0
14	A	468	0	0	7	0
All	All	6622	0	5999	28	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (28) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:437[B]:GLU:OE1	14:A:902:HOH:O	1.76	1.01
1:A:620[B]:GLN:NE2	14:A:901:HOH:O	1.58	0.82
1:A:620[A]:GLN:NE2	14:A:901:HOH:O	2.21	0.72
1:A:58:MET:CE	1:A:586:GLU:HG2	2.32	0.59
1:A:400:ARG:O	1:A:404[B]:THR:HG23	2.02	0.58
1:A:162:SER:HB3	1:A:258[B]:ILE:HD13	1.86	0.58
1:A:612:TYR:CZ	1:A:616:MET:HG3	2.39	0.57
1:A:688[B]:ARG:NH1	14:A:903:HOH:O	1.86	0.55
1:A:131[B]:ILE:O	1:A:131[B]:ILE:HG23	2.07	0.55
1:A:58:MET:HE1	1:A:586:GLU:HG2	1.92	0.50
1:A:90:PHE:CE2	1:A:94:LYS:HE2	2.46	0.50
1:A:96[B]:ILE:HD13	14:A:942:HOH:O	2.12	0.48

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:367:GLU:OE1	1:A:662[A]:ARG:NH1	2.42	0.48
1:A:131[B]:ILE:O	1:A:131[B]:ILE:CG2	2.63	0.47
1:A:626[B]:SER:HA	14:A:1082:HOH:O	2.15	0.46
1:A:626[A]:SER:HA	14:A:1082:HOH:O	2.16	0.45
1:A:258[A]:ILE:HD13	1:A:294:VAL:HB	1.99	0.45
1:A:246:TRP:CD1	9:A:806:NAG:H83	2.52	0.44
1:A:131[A]:ILE:HG22	1:A:137:GLU:HG2	1.99	0.44
1:A:659[B]:ILE:HD13	1:A:659[B]:ILE:HA	1.67	0.44
1:A:506:PHE:HB2	1:A:509[A]:MET:HG3	2.00	0.43
1:A:198:GLY:O	1:A:223:LYS:HE2	2.18	0.43
1:A:551:LEU:HD22	1:A:556:TYR:HB2	2.00	0.42
1:A:58:MET:HE2	1:A:586:GLU:HG2	1.99	0.42
1:A:514[A]:LYS:HG3	13:A:813:TKZ:C1	2.49	0.42
1:A:387:ASP:HA	1:A:388:PRO:HA	1.89	0.41
1:A:517[A]:SER:HB2	1:A:694:PRO:HG3	2.02	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	754/707 (107%)	735 (98%)	16 (2%)	3 (0%)	34 17

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	544	ASN
1	A	382	VAL
1	A	548	GLY

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	656/603 (109%)	647 (99%)	9 (1%)	67 50

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	174	LEU
1	A	187	LYS
1	A	519	ASN
1	A	537	TYR
1	A	542	GLU
1	A	600	TYR
1	A	668[A]	LEU
1	A	668[B]	LEU
1	A	673	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	136	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

17 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	NAG	B	1	1,2	14,14,15	0.35	0	17,19,21	1.13	2 (11%)
2	NAG	B	2	2	14,14,15	0.57	0	17,19,21	1.48	3 (17%)
2	BMA	B	3	2	11,11,12	0.69	0	15,15,17	1.02	1 (6%)
3	NAG	C	1	3,1	14,14,15	0.65	0	17,19,21	1.31	2 (11%)
3	NAG	C	2	3	14,14,15	0.48	0	17,19,21	0.90	1 (5%)
3	BMA	C	3	3	11,11,12	0.51	0	15,15,17	0.97	1 (6%)
3	MAN	C	4	3	11,11,12	0.54	0	15,15,17	1.03	1 (6%)
3	MAN	C	5	3	11,11,12	0.73	0	15,15,17	0.90	1 (6%)
4	NAG	D	1	4,1	14,14,15	0.70	0	17,19,21	1.46	3 (17%)
4	NAG	D	2	4	14,14,15	0.71	0	17,19,21	1.37	3 (17%)
5	NAG	E	1	1,5	14,14,15	0.70	0	17,19,21	1.38	2 (11%)
5	NAG	E	2	5	14,14,15	0.58	0	17,19,21	1.12	3 (17%)
5	BMA	E	3	5	11,11,12	0.58	0	15,15,17	1.33	3 (20%)
5	FUC	E	4	5	10,10,11	0.80	0	14,14,16	0.74	0
2	NAG	F	1	1,2	14,14,15	0.52	0	17,19,21	0.95	0
2	NAG	F	2	2	14,14,15	0.46	0	17,19,21	0.70	0
2	BMA	F	3	2	11,11,12	0.60	0	15,15,17	1.22	2 (13%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	B	1	1,2	-	0/6/23/26	0/1/1/1
2	NAG	B	2	2	-	2/6/23/26	0/1/1/1
2	BMA	B	3	2	-	0/2/19/22	0/1/1/1
3	NAG	C	1	3,1	-	1/6/23/26	0/1/1/1
3	NAG	C	2	3	-	2/6/23/26	0/1/1/1
3	BMA	C	3	3	-	0/2/19/22	0/1/1/1
3	MAN	C	4	3	-	0/2/19/22	0/1/1/1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	MAN	C	5	3	-	2/2/19/22	0/1/1/1
4	NAG	D	1	4,1	-	0/6/23/26	0/1/1/1
4	NAG	D	2	4	-	0/6/23/26	0/1/1/1
5	NAG	E	1	1,5	-	2/6/23/26	0/1/1/1
5	NAG	E	2	5	-	2/6/23/26	0/1/1/1
5	BMA	E	3	5	-	2/2/19/22	0/1/1/1
5	FUC	E	4	5	-	-	0/1/1/1
2	NAG	F	1	1,2	-	0/6/23/26	0/1/1/1
2	NAG	F	2	2	-	0/6/23/26	0/1/1/1
2	BMA	F	3	2	-	2/2/19/22	0/1/1/1

There are no bond length outliers.

All (28) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	2	NAG	C2-N2-C7	3.92	128.49	122.90
4	D	1	NAG	C1-O5-C5	3.61	117.08	112.19
3	C	1	NAG	C1-O5-C5	3.52	116.96	112.19
2	F	3	BMA	C1-O5-C5	3.48	116.90	112.19
5	E	1	NAG	C2-N2-C7	3.41	127.76	122.90
5	E	3	BMA	C3-C4-C5	2.91	115.43	110.24
5	E	3	BMA	C1-O5-C5	2.86	116.07	112.19
4	D	1	NAG	C4-C3-C2	2.77	115.07	111.02
4	D	2	NAG	C2-N2-C7	2.76	126.83	122.90
4	D	2	NAG	O5-C5-C6	2.70	111.44	107.20
3	C	5	MAN	O5-C5-C6	2.70	111.44	107.20
5	E	2	NAG	C2-N2-C7	2.67	126.70	122.90
3	C	1	NAG	O5-C1-C2	-2.58	107.22	111.29
4	D	2	NAG	C4-C3-C2	2.56	114.77	111.02
5	E	1	NAG	C8-C7-N2	2.48	120.30	116.10
3	C	3	BMA	C1-O5-C5	2.45	115.51	112.19
2	B	2	NAG	C4-C3-C2	2.39	114.51	111.02
3	C	2	NAG	C1-O5-C5	2.34	115.36	112.19
2	B	2	NAG	C8-C7-N2	2.28	119.97	116.10
2	F	3	BMA	C3-C4-C5	2.20	114.17	110.24
5	E	2	NAG	O5-C5-C6	2.16	110.59	107.20
5	E	2	NAG	C4-C3-C2	2.14	114.16	111.02
3	C	4	MAN	C1-O5-C5	2.09	115.03	112.19
5	E	3	BMA	C2-C3-C4	2.08	114.49	110.89
2	B	1	NAG	C4-C3-C2	2.08	114.06	111.02
2	B	3	BMA	C3-C4-C5	2.06	113.91	110.24

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	D	1	NAG	O5-C5-C6	2.03	110.39	107.20
2	B	1	NAG	O5-C5-C6	2.03	110.38	107.20

There are no chirality outliers.



All (15) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	C	5	MAN	O5-C5-C6-O6
5	E	3	BMA	O5-C5-C6-O6
3	C	5	MAN	C4-C5-C6-O6
2	B	2	NAG	C8-C7-N2-C2
2	B	2	NAG	O7-C7-N2-C2
5	E	1	NAG	C8-C7-N2-C2
5	E	1	NAG	O7-C7-N2-C2
5	E	3	BMA	C4-C5-C6-O6
2	F	3	BMA	C4-C5-C6-O6
3	C	2	NAG	C4-C5-C6-O6
5	E	2	NAG	C4-C5-C6-O6
3	C	2	NAG	O5-C5-C6-O6
5	E	2	NAG	O5-C5-C6-O6
3	C	1	NAG	C4-C5-C6-O6
2	F	3	BMA	O5-C5-C6-O6


There are no ring outliers.


No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.









5.6 Ligand geometry [i](#)

Of 13 ligands modelled in this entry, 6 are monoatomic - leaving 7 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
9	NAG	A	806	1	14,14,15	0.88	0	17,19,21	1.43	4 (23%)
9	NAG	A	805	1	14,14,15	0.67	0	17,19,21	1.48	2 (11%)
13	TKZ	A	813	6	48,51,51	1.00	2 (4%)	57,68,68	1.19	3 (5%)
10	EDO	A	807	-	3,3,3	0.44	0	2,2,2	0.46	0
10	EDO	A	810	-	3,3,3	0.47	0	2,2,2	0.39	0
10	EDO	A	808	-	3,3,3	0.42	0	2,2,2	0.60	0
11	PEG	A	809	-	6,6,6	0.45	0	5,5,5	0.30	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
9	NAG	A	806	1	-	0/6/23/26	0/1/1/1
9	NAG	A	805	1	-	0/6/23/26	0/1/1/1
13	TKZ	A	813	6	-	9/47/60/60	0/3/3/3
10	EDO	A	807	-	-	1/1/1/1	-
10	EDO	A	810	-	-	0/1/1/1	-
10	EDO	A	808	-	-	0/1/1/1	-
11	PEG	A	809	-	-	1/4/4/4	-

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
13	A	813	TKZ	C24-C26	-4.65	1.41	1.50
13	A	813	TKZ	C13-N3	-2.01	1.38	1.43

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
13	A	813	TKZ	C12-C13-N3	-5.07	117.73	121.62
9	A	806	NAG	C1-O5-C5	3.93	117.52	112.19
9	A	805	NAG	C1-O5-C5	3.84	117.40	112.19
9	A	805	NAG	O5-C5-C6	3.40	112.53	107.20
13	A	813	TKZ	C24-C26-N4	3.04	119.89	116.86
13	A	813	TKZ	C18-C13-N3	2.72	122.50	118.87
9	A	806	NAG	O3-C3-C4	-2.47	104.64	110.35
9	A	806	NAG	O5-C5-C6	2.03	110.39	107.20
9	A	806	NAG	O5-C1-C2	-2.03	108.09	111.29

There are no chirality outliers.

All (11) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
13	A	813	TKZ	C22-C23-C24-C26
13	A	813	TKZ	C23-C24-C26-O10
13	A	813	TKZ	C32-C33-O1-C1
13	A	813	TKZ	C22-C23-C24-C25
13	A	813	TKZ	O2-C33-O1-C1
10	A	807	EDO	O1-C1-C2-O2

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
13	A	813	TKZ	C25-C24-C26-O10
13	A	813	TKZ	C27-C28-C29-C30
13	A	813	TKZ	C32-C31-N3-C13
13	A	813	TKZ	C4-C3-C30-C29
11	A	809	PEG	O2-C3-C4-O4

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
9	A	806	NAG	1	0
13	A	813	TKZ	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	694/707 (98%)	0.51	51 (7%) 15 19	22, 32, 53, 97	0

All (51) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	546	PHE	10.8
1	A	155	SER	5.9
1	A	543	THR	5.7
1	A	55	LYS	5.3
1	A	507	SER	5.1
1	A	542	GLU	4.5
1	A	656[A]	SER	4.4
1	A	154	VAL	4.1
1	A	153	ASN	4.0
1	A	506	PHE	3.9
1	A	541	TRP	3.9
1	A	136	ASN	3.6
1	A	135	GLY	3.5
1	A	487	GLY	3.4
1	A	337	PHE	3.4
1	A	719	VAL	3.4
1	A	134	ASP	3.4
1	A	131[A]	ILE	3.3
1	A	56	HIS	3.2
1	A	544	ASN	3.2
1	A	659[A]	ILE	2.9
1	A	620[A]	GLN	2.9
1	A	505	GLU	2.9
1	A	124	HIS	2.8
1	A	104	GLY	2.7
1	A	336	ASN	2.6
1	A	545	LYS	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	428	LEU	2.6
1	A	123	THR	2.5
1	A	490	GLY	2.5
1	A	386	ILE	2.5
1	A	700	TYR	2.4
1	A	133	GLU	2.4
1	A	338	SER	2.4
1	A	547	SER	2.4
1	A	456	ILE	2.3
1	A	515	LEU	2.3
1	A	521	PHE	2.3
1	A	125	PRO	2.3
1	A	284	ALA	2.3
1	A	340	GLN	2.2
1	A	651	GLN	2.2
1	A	209	PHE	2.2
1	A	248	LEU	2.2
1	A	407	LYS	2.1
1	A	189	GLU	2.1
1	A	408	GLU	2.1
1	A	132	ASN	2.1
1	A	191	ASP	2.1
1	A	450	ILE	2.1
1	A	335	GLY	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [\(i\)](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	BMA	B	3	11/12	0.52	0.59	93,100,105,105	0
5	BMA	E	3	11/12	0.54	0.53	109,116,121,124	0
2	NAG	B	2	14/15	0.60	0.35	46,61,72,83	0
5	FUC	E	4	10/11	0.66	0.48	74,81,84,86	0


Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	BMA	F	3	11/12	0.68	0.30	69,74,82,89	0
4	NAG	D	2	14/15	0.69	0.49	88,92,96,104	0
4	NAG	D	1	14/15	0.72	0.27	57,65,82,89	0
5	NAG	E	1	14/15	0.79	0.23	46,51,57,62	0
3	MAN	C	5	11/12	0.81	0.27	67,74,84,91	0
5	NAG	E	2	14/15	0.83	0.35	59,69,81,98	0
3	NAG	C	1	14/15	0.85	0.13	27,35,44,57	0
3	NAG	C	2	14/15	0.89	0.20	44,47,54,64	0
3	MAN	C	4	11/12	0.90	0.11	49,53,56,56	0
3	BMA	C	3	11/12	0.91	0.11	45,47,53,54	0
2	NAG	F	2	14/15	0.91	0.14	42,51,59,64	0
2	NAG	F	1	14/15	0.91	0.12	34,37,40,48	0
2	NAG	B	1	14/15	0.92	0.17	38,45,51,56	0




The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.








Electron density around Chain C:


$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around Chain D:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
12	NA	A	811	1/1	0.70	0.21	55,55,55,55	0
9	NAG	A	805	14/15	0.76	0.31	67,77,84,84	0
9	NAG	A	806	14/15	0.82	0.14	37,52,57,60	0
10	EDO	A	807	4/4	0.84	0.28	49,57,60,62	0
10	EDO	A	808	4/4	0.87	0.17	50,57,59,62	0
11	PEG	A	809	7/7	0.89	0.13	61,62,68,69	0
10	EDO	A	810	4/4	0.89	0.22	44,55,58,64	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
12	NA	A	812	1/1	0.89	0.38	57,57,57,57	0
13	TKZ	A	813	49/49	0.91	0.17	23,45,69,77	0
8	CL	A	804	1/1	0.99	0.20	26,26,26,26	0
6	ZN	A	802	1/1	1.00	0.11	25,25,25,25	0
7	CA	A	803	1/1	1.00	0.10	24,24,24,24	0
6	ZN	A	801	1/1	1.00	0.09	25,25,25,25	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers [i](#)

There are no such residues in this entry.