



wwPDB EM Validation Summary Report ⓘ

May 27, 2024 – 01:52 PM JST

PDB ID : 7EKO
EMDB ID : EMD-31171
Title : CrClpP-S1
Authors : Wang, N.; Wang, Y.F.; Cong, Y.; Liu, C.M.
Deposited on : 2021-04-06
Resolution : 3.30 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:


EMDB validation analysis : 0.0.1.dev92
MolProbity : 4.02b-467
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36.2

1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:
ELECTRON MICROSCOPY

The reported resolution of this entry is 3.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)
Clashscore	158937	4297
Ramachandran outliers	154571	4023
Sidechain outliers	154315	3826

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	238	 68% 24% 7%
2	B	238	 64% 16% 19%
2	D	238	 59% 21% 19%
2	F	238	 7% 55% 17% 28%
3	C	296	 42% 17% 41%
3	E	296	 36% 22% 41%
3	G	296	 5% 45% 15% 40%
4	H	208	 49% 21% 29%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
4	J	208	<p>12% 52% 26% 21%</p>
4	M	208	<p>44% 19% 36%</p>
5	I	246	<p>61% 27% 12%</p>
6	K	251	<p>19% 64% 27% 7%</p>
7	L	250	<p>54% 20% 25%</p>
8	N	383	<p>6% 65% 23% 12%</p>
9	O	180	<p>10% 56% 32% 10%</p>

2 Entry composition [i](#)

There are 9 unique types of molecules in this entry. The entry contains 22470 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ATP-dependent Clp protease proteolytic subunit.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
1	A	222	1728	1093	288	333	14	0	0

- Molecule 2 is a protein called ATP-dependent Clp protease proteolytic subunit.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
2	B	193	1477	925	258	283	11	0	0
2	D	193	1477	925	258	283	11	0	0
2	F	172	1307	817	227	253	10	0	0

- Molecule 3 is a protein called ATP-dependent Clp protease proteolytic subunit.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
3	C	174	1329	836	229	252	12	0	0
3	E	174	1329	836	229	252	12	0	0
3	G	179	1369	862	235	260	12	0	0

- Molecule 4 is a protein called ATP-dependent Clp protease proteolytic subunit.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
4	H	148	1131	713	192	219	7	0	0
4	J	165	1279	805	220	247	7	0	0
4	M	133	1014	644	172	193	5	0	0

- Molecule 5 is a protein called ATP-dependent Clp protease proteolytic subunit.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
5	I	217	1745	1123	300	315	7	0	0

- Molecule 6 is a protein called ATP-dependent Clp protease proteolytic subunit.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
6	K	233	1841	1179	308	342	12	0	0

- Molecule 7 is a protein called ATP-dependent Clp protease proteolytic subunit.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
7	L	187	1494	950	256	275	13	0	0

- Molecule 8 is a protein called ATP-dependent Clp protease proteolytic subunit.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
8	N	338	2691	1699	474	504	14	0	0


- Molecule 9 is a protein called ATP-dependent Clp protease ATP-binding subunit CLPT4, chloroplastic.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
9	O	162	1259	807	207	239	6	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: ATP-dependent Clp protease proteolytic subunit




- Molecule 2: ATP-dependent Clp protease proteolytic subunit





- Molecule 2: ATP-dependent Clp protease proteolytic subunit










• Molecule 2: ATP-dependent Clp protease proteolytic subunit




• Molecule 3: ATP-dependent Clp protease proteolytic subunit




• Molecule 3: ATP-dependent Clp protease proteolytic subunit




• Molecule 3: ATP-dependent Clp protease proteolytic subunit




• Molecule 4: ATP-dependent Clp protease proteolytic subunit




• Molecule 4: ATP-dependent Clp protease proteolytic subunit




• Molecule 4: ATP-dependent Clp protease proteolytic subunit







• Molecule 5: ATP-dependent Clp protease proteolytic subunit




• Molecule 6: ATP-dependent Clp protease proteolytic subunit




• Molecule 7: ATP-dependent Clp protease proteolytic subunit



• Molecule 8: ATP-dependent Clp protease proteolytic subunit



• Molecule 9: ATP-dependent Clp protease ATP-binding subunit CLPT4, chloroplatic



4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	131245	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	38	Depositor
Minimum defocus (nm)	Not provided	
Maximum defocus (nm)	Not provided	
Magnification	Not provided	
Image detector	GATAN K2 SUMMIT (4k x 4k)	Depositor
Maximum map value	0.258	Depositor
Minimum map value	-0.142	Depositor
Average map value	0.000	Depositor
Map value standard deviation	0.007	Depositor
Recommended contour level	0.026	Depositor
Map size (Å)	284.688, 284.688, 284.688	wwPDB
Map dimensions	216, 216, 216	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	1.318, 1.318, 1.318	Depositor

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.47	0/1758	0.56	0/2376
2	B	0.50	0/1500	0.55	0/2028
2	D	0.43	0/1500	0.55	0/2028
2	F	0.39	0/1326	0.54	0/1792
3	C	0.48	0/1350	0.58	0/1814
3	E	0.39	0/1350	0.57	1/1814 (0.1%)
3	G	0.44	0/1391	0.58	0/1870
4	H	0.49	0/1149	0.62	2/1553 (0.1%)
4	J	0.39	0/1301	0.52	0/1758
4	M	0.45	0/1031	0.58	0/1393
5	I	0.47	0/1790	0.54	0/2428
6	K	0.42	0/1881	0.57	0/2542
7	L	0.42	0/1529	0.57	0/2072
8	N	0.45	0/2746	0.53	1/3718 (0.0%)
9	O	0.45	0/1285	0.53	0/1743
All	All	0.44	0/22887	0.56	4/30929 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
6	K	0	1
All	All	0	2

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
8	N	170	GLY	C-N-CA	6.84	138.81	121.70

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	H	483	MET	CA-CB-CG	6.10	123.67	113.30
4	H	483	MET	CB-CG-SD	-5.54	95.79	112.40
3	E	29	LEU	CA-CB-CG	5.37	127.64	115.30

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	110	SER	Mainchain
6	K	134	PRO	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1728	0	1718	48	0
2	B	1477	0	1477	38	0
2	D	1477	0	1477	44	0
2	F	1307	0	1301	28	0
3	C	1329	0	1330	38	0
3	E	1329	0	1330	51	0
3	G	1369	0	1365	30	0
4	H	1131	0	1123	41	0
4	J	1279	0	1263	56	0
4	M	1014	0	1013	29	0
5	I	1745	0	1731	57	0
6	K	1841	0	1867	58	0
7	L	1494	0	1481	41	0
8	N	2691	0	2708	63	0
9	O	1259	0	1246	35	0
All	All	22470	0	22430	549	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 12.

The worst 5 of 549 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:141:SER:OG	5:I:156:ALA:CB	2.13	0.96
1:A:141:SER:OG	5:I:156:ALA:HB2	1.67	0.93
1:A:183:ARG:HH22	2:B:165:GLN:NE2	1.71	0.87
1:A:183:ARG:NH2	2:B:165:GLN:HE21	1.74	0.85
2:D:46:VAL:HA	2:D:49:ARG:HD2	1.59	0.84

There are no symmetry-related clashes.

5.3 Torsion angles [\(i\)](#)

5.3.1 Protein backbone [\(i\)](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	220/238 (92%)	200 (91%)	20 (9%)	0	100	100
2	B	191/238 (80%)	182 (95%)	9 (5%)	0	100	100
2	D	191/238 (80%)	181 (95%)	10 (5%)	0	100	100
2	F	170/238 (71%)	162 (95%)	8 (5%)	0	100	100
3	C	172/296 (58%)	164 (95%)	8 (5%)	0	100	100
3	E	172/296 (58%)	161 (94%)	11 (6%)	0	100	100
3	G	177/296 (60%)	168 (95%)	9 (5%)	0	100	100
4	H	146/208 (70%)	140 (96%)	6 (4%)	0	100	100
4	J	163/208 (78%)	155 (95%)	8 (5%)	0	100	100
4	M	131/208 (63%)	127 (97%)	4 (3%)	0	100	100
5	I	215/246 (87%)	201 (94%)	14 (6%)	0	100	100
6	K	231/251 (92%)	206 (89%)	24 (10%)	1 (0%)	34	66
7	L	185/250 (74%)	170 (92%)	15 (8%)	0	100	100
8	N	336/383 (88%)	314 (94%)	22 (6%)	0	100	100
9	O	160/180 (89%)	153 (96%)	6 (4%)	1 (1%)	25	57
All	All	2860/3774 (76%)	2684 (94%)	174 (6%)	2 (0%)	54	81

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
9	O	88	SER
6	K	35	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	186/200 (93%)	181 (97%)	5 (3%)	44	71
2	B	161/199 (81%)	160 (99%)	1 (1%)	86	91
2	D	161/199 (81%)	160 (99%)	1 (1%)	86	91
2	F	141/199 (71%)	138 (98%)	3 (2%)	53	75
3	C	140/250 (56%)	140 (100%)	0	100	100
3	E	140/250 (56%)	140 (100%)	0	100	100
3	G	144/250 (58%)	136 (94%)	8 (6%)	21	52
4	H	120/175 (69%)	117 (98%)	3 (2%)	47	72
4	J	136/175 (78%)	133 (98%)	3 (2%)	52	74
4	M	105/175 (60%)	100 (95%)	5 (5%)	25	56
5	I	183/207 (88%)	180 (98%)	3 (2%)	62	79
6	K	197/213 (92%)	188 (95%)	9 (5%)	27	58
7	L	161/217 (74%)	158 (98%)	3 (2%)	57	77
8	N	290/327 (89%)	287 (99%)	3 (1%)	76	86
9	O	130/146 (89%)	115 (88%)	15 (12%)	5	22
All	All	2395/3182 (75%)	2333 (97%)	62 (3%)	49	71

5 of 62 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
6	K	61	PHE
9	O	84	LEU
6	K	210	LYS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
9	O	82	ASP
9	O	94	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 15 such sidechains are listed below:

Mol	Chain	Res	Type
4	J	370	GLN
4	M	451	HIS
4	J	412	GLN
8	N	229	GLN
7	L	171	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-31171. These allow visual inspection of the internal detail of the map and identification of artifacts.

No raw map or half-maps were deposited for this entry and therefore no images, graphs, etc. pertaining to the raw map can be shown.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)

6.2.1 Primary map




The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 95



Y Index: 126




Z Index: 99


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [i](#)


6.4.1 Primary map



X



Y




Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.

6.5 Orthogonal surface views [i](#)


6.5.1 Primary map



X



Y



Z

The images above show the 3D surface view of the map at the recommended contour level 0.026. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.


6.6 Mask visualisation [i](#)

This section was not generated. No masks/segmentation were deposited.

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [\(i\)](#)



The volume at the recommended contour level is 193 nm³; this corresponds to an approximate mass of 175 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum i



*Reported resolution corresponds to spatial frequency of 0.303 Å⁻¹


8 Fourier-Shell correlation

This section was not generated. No FSC curve or half-maps provided.

9 Map-model fit [i](#)


This section contains information regarding the fit between EMDB map EMD-31171 and PDB model 7EKO. Per-residue inclusion information can be found in section [3](#) on page [6](#).

9.1 Map-model overlay [i](#)




The images above show the 3D surface view of the map at the recommended contour level 0.026 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.026).

































9.4 Atom inclusion [i](#)



At the recommended contour level, 83% of all backbone atoms, 78% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary

The table lists the average atom inclusion at the recommended contour level (0.026) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	 0.7840	 0.5110
A	 0.8320	 0.5430
B	 0.8730	 0.5600
C	 0.8700	 0.5580
D	 0.8220	 0.5350
E	 0.8030	 0.5170
F	 0.7390	 0.4780
G	 0.7680	 0.4990
H	 0.8400	 0.5490
I	 0.8300	 0.5270
J	 0.6620	 0.4490
K	 0.6290	 0.4270
L	 0.8080	 0.5280
M	 0.8280	 0.5370
N	 0.7930	 0.5190
O	 0.6710	 0.4470

