



wwPDB NMR Structure Validation Summary Report ⓘ

Jun 6, 2023 – 05:56 pm BST

PDB ID : 7QYI
BMRB ID : 34704
Title : Solution structure of the DNA-binding minor pilin FimT from Legionella pneumophila
Authors : Braus, S.A.G.; Hospenthal, M.K.; Gossert, A.D.
Deposited on : 2022-01-28

This is a wwPDB NMR Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
wwPDB-RCI : v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV : Wang et al. (2010)
wwPDB-ShiftChecker : v1.2
BMRB Restraints Analysis : v1.2
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.33


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

SOLUTION NMR

The overall completeness of chemical shifts assignment is 87%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	158937	12864
Ramachandran outliers	154571	11451
Sidechain outliers	154315	11428

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	125	73% 23%

2 Ensemble composition and analysis i

This entry contains 20 models. Model 7 is the overall representative, medoid model (most similar to other models). The authors have identified model 1 as representative, based on the following criterion: *target function*.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model
1	A:31-A:63, A:68-A:104, A:110-A:117, A:122-A:139 (96)	0.58	7

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 3 clusters and 1 single-model cluster was found.

Cluster number	Models
1	1, 2, 3, 4, 6, 7, 8, 9, 12, 13, 14, 17, 18
2	5, 15, 16, 19
3	10, 20
Single-model clusters	11

3 Entry composition

There is only 1 type of molecule in this entry. The entry contains 2041 atoms, of which 1027 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called Pilus assembly protein.


Mol	Chain	Residues	Atoms					Trace	
			Total	C	H	N	O		S
1	A	125	2041	635	1027	195	182	2	0

4 Residue-property plots [i](#)

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.


- Molecule 1: Pilus assembly protein



4.2 Residue scores for the representative (medoid) model from the NMR ensemble

The representative model is number 7. Colouring as in section 4.1 above.

- Molecule 1: Pilus assembly protein



5 Refinement protocol and experimental data overview

The models were refined using the following method: *simulated annealing, molecular dynamics*.

Of the 100 calculated structures, 20 were deposited, based on the following criterion: *structures with the least restraint violations*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
CYANA	structure calculation	3.98.2
Amber	geometry optimization	20

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	working_cs.cif
Number of chemical shift lists	1
Total number of shifts	1511
Number of shifts mapped to atoms	1510
Number of unparsed shifts	0
Number of shifts with mapping errors	1
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	87%

6 Model quality [i](#)

6.1 Standard geometry [i](#)

There are no covalent bond-length or bond-angle outliers.

There are no bond-length outliers.

There are no bond-angle outliers.

There are no chirality outliers.

There are no planarity outliers.

6.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	792	809	809	4±2
All	All	15840	16180	16180	83

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

5 of 19 unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:75:LEU:CD2	1:A:88:ILE:HD12	0.54	2.33	15	1
1:A:75:LEU:HD22	1:A:88:ILE:HD12	0.53	1.80	3	10
1:A:35:THR:O	1:A:35:THR:HG22	0.52	2.04	8	6
1:A:58:ILE:HD12	1:A:75:LEU:HD21	0.51	1.83	6	8
1:A:46:TYR:CD2	1:A:46:TYR:O	0.49	2.65	6	12

6.3 Torsion angles [i](#)

6.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	96/125 (77%)	81±2 (84±2%)	13±2 (14±2%)	2±1 (2±1%)	11	53
All	All	1920/2500 (77%)	1621 (84%)	262 (14%)	37 (2%)	11	53

5 of 6 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	83	ASN	20
1	A	134	ASN	10
1	A	69	TRP	3
1	A	32	GLU	2
1	A	80	GLN	1

6.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	89/113 (79%)	87±1 (97±1%)	2±1 (3±1%)	49	91
All	All	1780/2260 (79%)	1734 (97%)	46 (3%)	49	91

5 of 15 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	87	LEU	8
1	A	93	TRP	8
1	A	48	LYS	6
1	A	131	LYS	6
1	A	68	ASN	3

6.3.3 RNA [i](#)

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.6 Ligand geometry [i](#)

There are no ligands in this entry.

6.7 Other polymers [i](#)

There are no such molecules in this entry.

6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

7 Chemical shift validation [i](#)

The completeness of assignment taking into account all chemical shift lists is 87% for the well-defined parts and 84% for the entire structure.

7.1 Chemical shift list 1

File name: working_cs.cif

Chemical shift list name: *FimT_after_corrections.str*

7.1.1 Bookkeeping [i](#)

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	1511
Number of shifts mapped to atoms	1510
Number of unparsed shifts	0
Number of shifts with mapping errors	1
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	11

The following assigned chemical shifts were not mapped to the molecules present in the coordinate file.

- No matching atom found in the structure. All 1 occurrences are reported below.

List ID	Chain	Res	Type	Atom	Shift Data		
					Value	Uncertainty	Ambiguity
1	A	53	HIS	HD1	8.604	0.020	1

7.1.2 Chemical shift referencing [i](#)

The following table shows the suggested chemical shift referencing corrections.

Nucleus	# values	Correction \pm precision, ppm	Suggested action
$^{13}\text{C}_\alpha$	125	-0.30 ± 0.16	None needed (< 0.5 ppm)
$^{13}\text{C}_\beta$	117	0.12 ± 0.10	None needed (< 0.5 ppm)
$^{13}\text{C}'$	0	—	None (insufficient data)
^{15}N	109	0.29 ± 0.44	None needed (< 0.5 ppm)

7.1.3 Completeness of resonance assignments [i](#)

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 87%, i.e. 1219 atoms were assigned a chemical shift out of a possible 1397. 0 out of 13 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	¹ H	¹³ C	¹⁵ N
Backbone	376/478 (79%)	190/193 (98%)	96/192 (50%)	90/93 (97%)
Sidechain	735/800 (92%)	500/519 (96%)	217/243 (89%)	18/38 (47%)
Aromatic	108/119 (91%)	54/60 (90%)	49/51 (96%)	5/8 (62%)
Overall	1219/1397 (87%)	744/772 (96%)	362/486 (74%)	113/139 (81%)

7.1.4 Statistically unusual chemical shifts [i](#)


The following table lists the statistically unusual chemical shifts. These are statistical measures, and large deviations from the mean do not necessarily imply incorrect assignments. Molecules containing paramagnetic centres or hemes are expected to give rise to anomalous chemical shifts.

List Id	Chain	Res	Type	Atom	Shift, ppm	Expected range, ppm	Z-score
1	A	133	THR	HG1	6.28	0.08 – 2.19	24.4
1	A	35	THR	HG1	5.10	0.08 – 2.19	18.8
1	A	130	ASN	HB3	0.39	1.12 – 4.38	-7.2
1	A	72	GLY	HA3	1.62	2.08 – 5.71	-6.2
1	A	102	TRP	HE3	4.97	5.27 – 9.37	-5.7
1	A	111	ILE	HD11	-0.86	-0.72 – 2.09	-5.5
1	A	111	ILE	HD12	-0.86	-0.72 – 2.09	-5.5
1	A	111	ILE	HD13	-0.86	-0.72 – 2.09	-5.5
1	A	43	ALA	HB1	0.11	0.14 – 2.58	-5.1
1	A	43	ALA	HB2	0.11	0.14 – 2.58	-5.1
1	A	43	ALA	HB3	0.11	0.14 – 2.58	-5.1

7.1.5 Random Coil Index (RCI) plots [i](#)

The image below reports *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition. If well-defined core and ill-defined regions are not identified then it is shown as gray bars.

Random coil index (RCI) for chain A:



8 NMR restraints analysis

8.1 Conformationally restricting restraints

The following table provides the summary of experimentally observed NMR restraints in different categories. Restraints are classified into different categories based on the sequence separation of the atoms involved.

Description	Value
Total distance restraints	2148
Intra-residue ($ i-j =0$)	572
Sequential ($ i-j =1$)	489
Medium range ($ i-j >1$ and $ i-j <5$)	322
Long range ($ i-j \geq 5$)	765
Inter-chain	0
Hydrogen bond restraints	0
Disulfide bond restraints	0
Total dihedral-angle restraints	0
Number of unmapped restraints	2
Number of restraints per residue	17.2
Number of long range restraints per residue ¹	6.1

¹Long range hydrogen bonds and disulfide bonds are counted as long range restraints while calculating the number of long range restraints per residue

8.2 Residual restraint violations

This section provides the overview of the restraint violations analysis. The violations are binned as small, medium and large violations based on its absolute value. Average number of violations per model is calculated by dividing the total number of violations in each bin by the size of the ensemble.

8.2.1 Average number of distance violations per model

Distance violations less than 0.1 Å are not included in the calculation.

Bins (Å)	Average number of violations per model	Max (Å)
0.1-0.2 (Small)	45.6	0.2
0.2-0.5 (Medium)	33.0	0.5
>0.5 (Large)	7.2	1.95

8.2.2 Average number of dihedral-angle violations per model

Dihedral-angle violations less than 1° are not included in the calculation. There are no dihedral-angle violations

9 Distance violation analysis [i](#)


9.1 Summary of distance violations [i](#)

The following table shows the summary of distance violations in different restraint categories based on the sequence separation of the atoms involved. Each category is further sub-divided into three sub-categories based on the atoms involved. Violations less than 0.1 Å are not included in the statistics.

Restrains type	Count	% ¹	Violated ³			Consistently Violated ⁴		
			Count	% ²	% ¹	Count	% ²	% ¹
Intra-residue (i-j =0)	572	26.6	38	6.6	1.8	3	0.5	0.1
Backbone-Backbone	1	0.0	0	0.0	0.0	0	0.0	0.0
Backbone-Sidechain	427	19.9	32	7.5	1.5	3	0.7	0.1
Sidechain-Sidechain	144	6.7	6	4.2	0.3	0	0.0	0.0
Sequential (i-j =1)	489	22.8	109	22.3	5.1	13	2.7	0.6
Backbone-Backbone	102	4.7	3	2.9	0.1	0	0.0	0.0
Backbone-Sidechain	308	14.3	89	28.9	4.1	13	4.2	0.6
Sidechain-Sidechain	79	3.7	17	21.5	0.8	0	0.0	0.0
Medium range (i-j >1 & i-j <5)	322	15.0	51	15.8	2.4	3	0.9	0.1
Backbone-Backbone	56	2.6	6	10.7	0.3	0	0.0	0.0
Backbone-Sidechain	139	6.5	23	16.5	1.1	2	1.4	0.1
Sidechain-Sidechain	127	5.9	22	17.3	1.0	1	0.8	0.0
Long range (i-j ≥5)	765	35.6	109	14.2	5.1	2	0.3	0.1
Backbone-Backbone	74	3.4	6	8.1	0.3	0	0.0	0.0
Backbone-Sidechain	275	12.8	50	18.2	2.3	1	0.4	0.0
Sidechain-Sidechain	416	19.4	53	12.7	2.5	1	0.2	0.0
Inter-chain	0	0.0	0	0.0	0.0	0	0.0	0.0
Backbone-Backbone	0	0.0	0	0.0	0.0	0	0.0	0.0
Backbone-Sidechain	0	0.0	0	0.0	0.0	0	0.0	0.0
Sidechain-Sidechain	0	0.0	0	0.0	0.0	0	0.0	0.0
Hydrogen bond	0	0.0	0	0.0	0.0	0	0.0	0.0
Disulfide bond	0	0.0	0	0.0	0.0	0	0.0	0.0
Total	2148	100.0	307	14.3	14.3	21	1.0	1.0
Backbone-Backbone	233	10.8	15	6.4	0.7	0	0.0	0.0
Backbone-Sidechain	1149	53.5	194	16.9	9.0	19	1.7	0.9
Sidechain-Sidechain	766	35.7	98	12.8	4.6	2	0.3	0.1

¹ percentage calculated with respect to the total number of distance restraints, ² percentage calculated with respect to the number of restraints in a particular restraint category, ³ violated in at least one model, ⁴ violated in all the models

9.1.1 Bar chart : Distribution of distance restraints and violations [i](#)



Violated and consistently violated restraints are shown using different hatch patterns in their respective categories. The hydrogen bonds and disulfid bonds are counted in their appropriate category on the x-axis

9.2 Distance violation statistics for each model [i](#)

The following table provides the distance violation statistics for each model in the ensemble. Violations less than 0.1 Å are not included in the statistics.

Model ID	Number of violations						Mean (Å)	Max (Å)	SD ⁶ (Å)	Median (Å)
	IR ¹	SQ ²	MR ³	LR ⁴	IC ⁵	Total				
1	7	44	13	17	0	81	0.23	1.41	0.17	0.17
2	6	41	12	19	0	78	0.26	1.34	0.18	0.19
3	11	42	11	15	0	79	0.25	1.46	0.21	0.18
4	10	33	15	16	0	74	0.23	0.84	0.14	0.18
5	6	47	12	25	0	90	0.26	1.35	0.2	0.18
6	9	39	13	23	0	84	0.26	1.28	0.18	0.19
7	7	37	9	19	0	72	0.24	1.3	0.17	0.18
8	9	37	13	17	0	76	0.24	1.16	0.17	0.18
9	7	45	15	22	0	89	0.25	1.27	0.18	0.2
10	9	45	14	22	0	90	0.25	1.29	0.18	0.19
11	20	62	27	36	0	145	0.41	1.95	0.34	0.29


Continued on next page...

Continued from previous page...

Model ID	Number of violations						Mean (Å)	Max (Å)	SD ⁶ (Å)	Median (Å)
	IR ¹	SQ ²	MR ³	LR ⁴	IC ⁵	Total				
12	8	38	12	18	0	76	0.25	1.27	0.17	0.2
13	10	41	14	31	0	96	0.29	1.42	0.24	0.2
14	7	42	10	17	0	76	0.24	1.39	0.18	0.18
15	7	43	12	24	0	86	0.27	1.31	0.22	0.21
16	9	37	12	22	0	80	0.23	1.04	0.16	0.19
17	7	41	18	24	0	90	0.24	1.38	0.2	0.17
18	10	40	16	26	0	92	0.23	1.4	0.17	0.17
19	5	36	11	20	0	72	0.26	1.24	0.18	0.2
20	7	43	18	23	0	91	0.25	1.34	0.18	0.2

¹Intra-residue restraints, ²Sequential restraints, ³Medium range restraints, ⁴Long range restraints, ⁵Inter-chain restraints, ⁶Standard deviation

9.2.1 Bar graph : Distance Violation statistics for each model [\(i\)](#)



The mean(dot),median(x) and the standard deviation are shown in blue with respect to the y axis on the right


9.3 Distance violation statistics for the ensemble

Violation analysis may find that some restraints are violated in few models and some are violated in most of models. The following table provides this information as number of violated restraints for a given fraction of the ensemble. In total, 1841(IR:534, SQ:380, MR:271, LR:656, IC:0) restraints are not violated in the ensemble.

Number of violated restraints						Fraction of the ensemble	
IR ¹	SQ ²	MR ³	LR ⁴	IC ⁵	Total	Count ⁶	%
22	24	22	52	0	120	1	5.0
1	14	6	14	0	35	2	10.0
6	10	2	6	0	24	3	15.0
0	5	2	9	0	16	4	20.0
0	4	2	4	0	10	5	25.0
1	2	1	2	0	6	6	30.0
1	4	3	5	0	13	7	35.0
1	5	0	1	0	7	8	40.0
0	4	3	2	0	9	9	45.0
0	2	1	1	0	4	10	50.0
0	4	0	2	0	6	11	55.0
0	5	1	1	0	7	12	60.0
0	0	0	3	0	3	13	65.0
0	4	0	1	0	5	14	70.0
0	2	2	0	0	4	15	75.0
3	1	0	1	0	5	16	80.0
0	2	2	0	0	4	17	85.0
0	1	0	1	0	2	18	90.0
0	3	1	2	0	6	19	95.0
3	13	3	2	0	21	20	100.0

¹Intra-residue restraints, ²Sequential restraints, ³Medium range restraints, ⁴Long range restraints, ⁵Inter-chain restraints, ⁶ Number of models with violations


9.3.1 Bar graph : Distance violation statistics for the ensemble [i](#)



9.4 Most violated distance restraints in the ensemble [i](#)

9.4.1 Histogram : Distribution of mean distance violations [i](#)

The following histogram shows the distribution of the average value of the violation. The average is calculated for each restraint that is violated in more than one model over all the violated models in the ensemble



9.4.2 Table: Most violated distance restraints [i](#)

The following table provides the mean and the standard deviation of the violations for the 10 worst performing restraints, sorted by number of violated models and the mean violation value. The Key (restraint list ID, restraint ID) is the unique identifier for a given restraint. Rows with same key represent combinatorial or ambiguous restraints and are counted as a single restraint.

Key	Atom-1	Atom-2	Models ¹	Mean (Å)	SD ¹ (Å)	Median (Å)
(1,1584)	1:A:36:LEU:HD11	1:A:40:ILE:H	20	0.63	0.34	0.5
(1,1584)	1:A:36:LEU:HD12	1:A:40:ILE:H	20	0.63	0.34	0.5
(1,1584)	1:A:36:LEU:HD13	1:A:40:ILE:H	20	0.63	0.34	0.5
(1,1584)	1:A:36:LEU:HD21	1:A:40:ILE:H	20	0.63	0.34	0.5
(1,1584)	1:A:36:LEU:HD22	1:A:40:ILE:H	20	0.63	0.34	0.5
(1,1584)	1:A:36:LEU:HD23	1:A:40:ILE:H	20	0.63	0.34	0.5
(1,176)	1:A:50:GLN:HB3	1:A:51:ALA:H	20	0.6	0.05	0.61
(1,1148)	1:A:115:ASN:HA	1:A:115:ASN:HD21	20	0.59	0.03	0.59
(1,1533)	1:A:38:ASN:HB3	1:A:39:SER:H	20	0.52	0.09	0.53
(1,81)	1:A:50:GLN:H	1:A:50:GLN:HE21	20	0.41	0.19	0.4
(1,1230)	1:A:52:ILE:HA	1:A:115:ASN:HD21	20	0.37	0.06	0.37
(1,2062)	1:A:128:LEU:HD11	1:A:136:LYS:HG2	20	0.36	0.21	0.32
(1,2062)	1:A:128:LEU:HD12	1:A:136:LYS:HG2	20	0.36	0.21	0.32
(1,2062)	1:A:128:LEU:HD13	1:A:136:LYS:HG2	20	0.36	0.21	0.32
(1,2062)	1:A:128:LEU:HD21	1:A:136:LYS:HG2	20	0.36	0.21	0.32
(1,2062)	1:A:128:LEU:HD22	1:A:136:LYS:HG2	20	0.36	0.21	0.32

Continued on next page...

Continued from previous page...


Key	Atom-1	Atom-2	Models ¹	Mean (Å)	SD ¹ (Å)	Median (Å)
(1,2062)	1:A:128:LEU:HD23	1:A:136:LYS:HG2	20	0.36	0.21	0.32
(1,1410)	1:A:46:TYR:HB2	1:A:47:SER:H	20	0.32	0.05	0.34
(1,1298)	1:A:42:THR:HG21	1:A:43:ALA:H	20	0.32	0.02	0.32
(1,1298)	1:A:42:THR:HG22	1:A:43:ALA:H	20	0.32	0.02	0.32
(1,1298)	1:A:42:THR:HG23	1:A:43:ALA:H	20	0.32	0.02	0.32
(1,19)	1:A:45:GLN:HB2	1:A:46:TYR:H	20	0.31	0.05	0.33

¹Number of violated models, ²Standard deviation

9.5 All violated distance restraints [i](#)

9.5.1 Histogram : Distribution of distance violations [i](#)

The following histogram shows the distribution of the absolute value of the violation for all violated restraints in the ensemble.



9.5.2 Table : All distance violations [i](#)

The following table provides the 10 worst performing restraints, sorted by the violation value. The Key (restraint list ID, restraint ID) is the unique identifier for a given restraint. Rows with same key represent combinatorial or ambiguous restraints and are counted as a single restraint.

Key	Atom-1	Atom-2	Model ID	Violation (Å)
(1,1236)	1:A:50:GLN:HE22	1:A:54:LEU:HD21	11	1.95
(1,1236)	1:A:50:GLN:HE22	1:A:54:LEU:HD22	11	1.95
(1,1236)	1:A:50:GLN:HE22	1:A:54:LEU:HD23	11	1.95
(1,1201)	1:A:38:ASN:HA	1:A:41:LYS:HB2	11	1.92
(1,1673)	1:A:48:LYS:HG2	1:A:49:ILE:HA	11	1.59
(1,1673)	1:A:48:LYS:HG3	1:A:49:ILE:HA	11	1.59
(1,1584)	1:A:36:LEU:HD11	1:A:40:ILE:H	3	1.46
(1,1584)	1:A:36:LEU:HD12	1:A:40:ILE:H	3	1.46
(1,1584)	1:A:36:LEU:HD13	1:A:40:ILE:H	3	1.46
(1,1584)	1:A:36:LEU:HD21	1:A:40:ILE:H	3	1.46
(1,1584)	1:A:36:LEU:HD22	1:A:40:ILE:H	3	1.46
(1,1584)	1:A:36:LEU:HD23	1:A:40:ILE:H	3	1.46
(1,1584)	1:A:36:LEU:HD11	1:A:40:ILE:H	13	1.42
(1,1584)	1:A:36:LEU:HD12	1:A:40:ILE:H	13	1.42
(1,1584)	1:A:36:LEU:HD13	1:A:40:ILE:H	13	1.42
(1,1584)	1:A:36:LEU:HD21	1:A:40:ILE:H	13	1.42
(1,1584)	1:A:36:LEU:HD22	1:A:40:ILE:H	13	1.42
(1,1584)	1:A:36:LEU:HD23	1:A:40:ILE:H	13	1.42
(1,1593)	1:A:36:LEU:HD11	1:A:98:TRP:HE3	1	1.41
(1,1593)	1:A:36:LEU:HD12	1:A:98:TRP:HE3	1	1.41
(1,1593)	1:A:36:LEU:HD13	1:A:98:TRP:HE3	1	1.41
(1,1593)	1:A:36:LEU:HD21	1:A:98:TRP:HE3	1	1.41
(1,1593)	1:A:36:LEU:HD22	1:A:98:TRP:HE3	1	1.41
(1,1593)	1:A:36:LEU:HD23	1:A:98:TRP:HE3	1	1.41
(1,1593)	1:A:36:LEU:HD11	1:A:98:TRP:HE3	18	1.4
(1,1593)	1:A:36:LEU:HD12	1:A:98:TRP:HE3	18	1.4
(1,1593)	1:A:36:LEU:HD13	1:A:98:TRP:HE3	18	1.4
(1,1593)	1:A:36:LEU:HD21	1:A:98:TRP:HE3	18	1.4
(1,1593)	1:A:36:LEU:HD22	1:A:98:TRP:HE3	18	1.4
(1,1593)	1:A:36:LEU:HD23	1:A:98:TRP:HE3	18	1.4
(1,1593)	1:A:36:LEU:HD11	1:A:98:TRP:HE3	14	1.39
(1,1593)	1:A:36:LEU:HD12	1:A:98:TRP:HE3	14	1.39
(1,1593)	1:A:36:LEU:HD13	1:A:98:TRP:HE3	14	1.39
(1,1593)	1:A:36:LEU:HD21	1:A:98:TRP:HE3	14	1.39
(1,1593)	1:A:36:LEU:HD22	1:A:98:TRP:HE3	14	1.39
(1,1593)	1:A:36:LEU:HD23	1:A:98:TRP:HE3	14	1.39
(1,1593)	1:A:36:LEU:HD11	1:A:98:TRP:HE3	17	1.38
(1,1593)	1:A:36:LEU:HD12	1:A:98:TRP:HE3	17	1.38
(1,1593)	1:A:36:LEU:HD13	1:A:98:TRP:HE3	17	1.38
(1,1593)	1:A:36:LEU:HD21	1:A:98:TRP:HE3	17	1.38
(1,1593)	1:A:36:LEU:HD22	1:A:98:TRP:HE3	17	1.38
(1,1593)	1:A:36:LEU:HD23	1:A:98:TRP:HE3	17	1.38
(1,1584)	1:A:36:LEU:HD11	1:A:40:ILE:H	5	1.35

10 Dihedral-angle violation analysis

No dihedral-angle restraints found