



# wwPDB EM Validation Summary Report ⓘ

Oct 6, 2024 – 04:25 AM JST

PDB ID : 7V2A  
EMDB ID : EMD-31639  
Title : SARS-CoV-2 Spike trimer in complex with XG014 Fab  
Authors : Wang, K.; Wang, X.; Pan, L.  
Deposited on : 2021-08-07  
Resolution : 3.40 Å (reported)  
Based on initial models : 6VXX, 6Z3P, 6PZE

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:


EMDB validation analysis : 0.0.1.dev113  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
MolProbity : 4.02b-467  
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)  
MapQ : 1.9.13  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.39

# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:  
*ELECTRON MICROSCOPY*

The reported resolution of this entry is 3.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	EM structures (#Entries)
Clashscore	210492	15764
Ramachandran outliers	207382	16835
Sidechain outliers	206894	16415

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion  $< 40\%$ ). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	1208	
1	B	1208	
1	C	1208	
2	D	216	
2	F	216	
2	H	216	
3	E	237	
3	G	237	

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
3	I	237	 <p>A horizontal bar chart representing the quality of chain. The bar is divided into three segments: a small red segment at the beginning, a green segment labeled '25%', a yellow segment labeled '25%', and a grey segment labeled '49%'. A small black dot is located at the end of the grey segment.</p>

## 2 Entry composition i

There are 4 unique types of molecules in this entry. The entry contains 28017 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Spike glycoprotein.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
1	A	988	7464	4774	1240	1416	34	0	0
1	B	988	7464	4774	1240	1416	34	0	0
1	C	988	7464	4774	1240	1416	34	0	0

There are 15 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	682	GLY	ARG	engineered mutation	UNP P0DTC2
A	683	SER	ARG	engineered mutation	UNP P0DTC2
A	685	SER	ARG	engineered mutation	UNP P0DTC2
A	986	PRO	LYS	engineered mutation	UNP P0DTC2
A	987	PRO	VAL	engineered mutation	UNP P0DTC2
B	682	GLY	ARG	engineered mutation	UNP P0DTC2
B	683	SER	ARG	engineered mutation	UNP P0DTC2
B	685	SER	ARG	engineered mutation	UNP P0DTC2
B	986	PRO	LYS	engineered mutation	UNP P0DTC2
B	987	PRO	VAL	engineered mutation	UNP P0DTC2
C	682	GLY	ARG	engineered mutation	UNP P0DTC2
C	683	SER	ARG	engineered mutation	UNP P0DTC2
C	685	SER	ARG	engineered mutation	UNP P0DTC2
C	986	PRO	LYS	engineered mutation	UNP P0DTC2
C	987	PRO	VAL	engineered mutation	UNP P0DTC2

- Molecule 2 is a protein called The light chain of XG014 Fab.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
2	D	111	799	496	139	162	2	0	0
2	F	111	799	496	139	162	2	0	0

*Continued on next page...*


Continued from previous page...

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
2	H	111	Total 799	C 496	N 139	O 162	S 2	0	0

- Molecule 3 is a protein called The heavy chain of XG014.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
3	E	120	Total 964	C 623	N 153	O 183	S 5	0	0
3	G	120	Total 964	C 623	N 153	O 183	S 5	0	0
3	I	120	Total 964	C 623	N 153	O 183	S 5	0	0

- Molecule 4 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula:  $C_8H_{15}NO_6$ ).



Mol	Chain	Residues	Atoms				AltConf
			Total	C	N	O	
4	A	1	Total 14	C 8	N 1	O 5	0
4	A	1	Total 14	C 8	N 1	O 5	0
4	A	1	Total 14	C 8	N 1	O 5	0
4	A	1	Total 14	C 8	N 1	O 5	0
4	A	1	Total 14	C 8	N 1	O 5	0

Continued on next page...


*Continued from previous page...*


Mol	Chain	Residues	Atoms				AltConf
			Total	C	N	O	
4	A	1	Total 14	8	1	5	0
4	A	1	Total 14	8	1	5	0
4	A	1	Total 14	8	1	5	0
4	B	1	Total 14	8	1	5	0
4	B	1	Total 14	8	1	5	0
4	B	1	Total 14	8	1	5	0
4	B	1	Total 14	8	1	5	0
4	B	1	Total 14	8	1	5	0
4	B	1	Total 14	8	1	5	0
4	B	1	Total 14	8	1	5	0
4	B	1	Total 14	8	1	5	0
4	B	1	Total 14	8	1	5	0
4	B	1	Total 14	8	1	5	0
4	C	1	Total 14	8	1	5	0
4	C	1	Total 14	8	1	5	0
4	C	1	Total 14	8	1	5	0
4	C	1	Total 14	8	1	5	0
4	C	1	Total 14	8	1	5	0
4	C	1	Total 14	8	1	5	0
4	C	1	Total 14	8	1	5	0
4	C	1	Total 14	8	1	5	0
4	C	1	Total 14	8	1	5	0

### 3 Residue-property plots


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### ● Molecule 1: Spike glycoprotein





• Molecule 1: Spike glycoprotein











VAL ASP LEU GLY ASP ASP ILE SER SER ILE ILE ALA SER VAL VAL VAL ASN ILE GLN LYS GLU ILE ASP ARG LEU ASN ASN ASN GLU VAL VAL ALA LYS ASN LYS ASN LYS LEU LEU LEU ILE ASP ASP ASP ASP ASP ASP GLN LEU LEU GLY TYR TYR TYR TYR TYR GLU GLN


• Molecule 2: The light chain of XG014 Fab



• Molecule 2: The light chain of XG014 Fab



• Molecule 2: The light chain of XG014 Fab





## 4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	414922	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING ONLY	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ( $e^-/\text{\AA}^2$ )	50	Depositor
Minimum defocus (nm)	1200	Depositor
Maximum defocus (nm)	2500	Depositor
Magnification	Not provided	
Image detector	GATAN K2 SUMMIT (4k x 4k)	Depositor
Maximum map value	0.071	Depositor
Minimum map value	-0.034	Depositor
Average map value	0.000	Depositor
Map value standard deviation	0.002	Depositor
Recommended contour level	0.01	Depositor
Map size ( $\text{\AA}$ )	416.0, 416.0, 416.0	wwPDB
Map dimensions	400, 400, 400	wwPDB
Map angles ( $^\circ$ )	90.0, 90.0, 90.0	wwPDB
Pixel spacing ( $\text{\AA}$ )	1.04, 1.04, 1.04	Depositor

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.58	1/7625 (0.0%)	0.54	0/10400
1	B	0.58	1/7625 (0.0%)	0.53	0/10400
1	C	0.58	1/7625 (0.0%)	0.54	0/10400
2	D	0.41	0/817	0.54	0/1113
2	F	0.41	0/817	0.54	0/1113
2	H	0.40	0/817	0.54	0/1113
3	E	0.50	0/996	0.57	0/1352
3	G	0.53	0/996	0.60	1/1352 (0.1%)
3	I	0.49	0/996	0.61	1/1352 (0.1%)
All	All	0.56	3/28314 (0.0%)	0.54	2/38595 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	B	0	1
1	C	0	1
2	D	0	1
2	F	0	1
2	H	0	1
3	E	0	1
3	G	0	1
All	All	0	8

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	1088	HIS	C-N	-5.44	1.21	1.34

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	1088	HIS	C-N	-5.42	1.21	1.34
1	B	1088	HIS	C-N	-5.41	1.21	1.34

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	G	110	TYR	CB-CA-C	-7.57	95.26	110.40
3	I	110	TYR	CB-CA-C	-7.50	95.39	110.40

There are no chirality outliers.

5 of 8 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	443	SER	Peptide
1	B	443	SER	Peptide
1	C	443	SER	Peptide
2	D	50	TYR	Peptide
3	E	104	ASN	Peptide

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	7464	0	7086	415	0
1	B	7464	0	7086	413	0
1	C	7464	0	7086	422	0
2	D	799	0	782	59	0
2	F	799	0	782	57	0
2	H	799	0	782	49	0
3	E	964	0	904	62	0
3	G	964	0	904	65	0
3	I	964	0	904	59	0
4	A	112	0	104	3	0
4	B	112	0	104	3	0
4	C	112	0	104	4	0
All	All	28017	0	26628	1365	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 25.

The worst 5 of 1365 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:697:MET:CE	1:C:864:LEU:HD11	1.11	1.56
1:A:697:MET:CE	1:B:864:LEU:HD11	1.13	1.54
1:A:864:LEU:HD11	1:C:697:MET:CE	1.11	1.51
1:A:864:LEU:CD1	1:C:697:MET:CE	1.90	1.48
1:B:697:MET:CE	1:C:864:LEU:CD1	1.90	1.46

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	964/1208 (80%)	872 (90%)	90 (9%)	2 (0%)	44	72
1	B	964/1208 (80%)	872 (90%)	90 (9%)	2 (0%)	44	72
1	C	964/1208 (80%)	871 (90%)	91 (9%)	2 (0%)	44	72
2	D	109/216 (50%)	96 (88%)	12 (11%)	1 (1%)	14	41
2	F	109/216 (50%)	96 (88%)	12 (11%)	1 (1%)	14	41
2	H	109/216 (50%)	96 (88%)	12 (11%)	1 (1%)	14	41
3	E	118/237 (50%)	107 (91%)	11 (9%)	0	100	100
3	G	118/237 (50%)	106 (90%)	12 (10%)	0	100	100
3	I	118/237 (50%)	107 (91%)	11 (9%)	0	100	100
All	All	3573/4983 (72%)	3223 (90%)	341 (10%)	9 (0%)	38	66

5 of 9 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	42	VAL
1	B	42	VAL
1	C	42	VAL
1	A	444	LYS
2	D	51	ASP

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	796/1053 (76%)	789 (99%)	7 (1%)	75	86
1	B	796/1053 (76%)	788 (99%)	8 (1%)	73	83
1	C	796/1053 (76%)	790 (99%)	6 (1%)	79	87
2	D	89/181 (49%)	89 (100%)	0	100	100
2	F	89/181 (49%)	89 (100%)	0	100	100
2	H	89/181 (49%)	89 (100%)	0	100	100
3	E	102/205 (50%)	101 (99%)	1 (1%)	73	83
3	G	102/205 (50%)	101 (99%)	1 (1%)	73	83
3	I	102/205 (50%)	101 (99%)	1 (1%)	73	83
All	All	2961/4317 (69%)	2937 (99%)	24 (1%)	77	87

5 of 24 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	577	ARG
1	C	357	ARG
1	C	238	PHE
1	C	489	TYR
1	A	662	CYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 48 such sidechains are listed below:



Mol	Chain	Res	Type
1	B	1005	GLN
1	C	536	ASN
1	B	1101	HIS
1	C	437	ASN
1	C	751	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

24 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
4	NAG	B	1305	1	14,14,15	0.41	0	17,19,21	0.63	1 (5%)
4	NAG	C	1302	1	14,14,15	0.35	0	17,19,21	0.57	0
4	NAG	B	1303	1	14,14,15	0.31	0	17,19,21	0.56	0
4	NAG	C	1306	1	14,14,15	0.19	0	17,19,21	0.57	0
4	NAG	B	1302	1	14,14,15	0.36	0	17,19,21	0.57	0
4	NAG	B	1308	1	14,14,15	0.37	0	17,19,21	1.54	1 (5%)
4	NAG	A	1307	1	14,14,15	0.37	0	17,19,21	0.36	0
4	NAG	A	1303	1	14,14,15	0.32	0	17,19,21	0.57	0
4	NAG	A	1308	1	14,14,15	0.37	0	17,19,21	1.58	1 (5%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	NAG	A	1306	1	14,14,15	0.19	0	17,19,21	0.57	0
4	NAG	C	1308	1	14,14,15	0.38	0	17,19,21	1.92	4 (23%)
4	NAG	C	1305	1	14,14,15	0.40	0	17,19,21	0.64	1 (5%)
4	NAG	C	1303	1	14,14,15	0.32	0	17,19,21	0.56	0
4	NAG	A	1304	1	14,14,15	0.22	0	17,19,21	0.48	0
4	NAG	C	1304	1	14,14,15	0.22	0	17,19,21	0.48	0
4	NAG	C	1301	1	14,14,15	0.38	0	17,19,21	0.39	0
4	NAG	A	1302	1	14,14,15	0.36	0	17,19,21	0.57	0
4	NAG	A	1305	1	14,14,15	0.41	0	17,19,21	0.64	1 (5%)
4	NAG	B	1301	1	14,14,15	0.39	0	17,19,21	0.39	0
4	NAG	B	1307	1	14,14,15	0.37	0	17,19,21	0.35	0
4	NAG	B	1306	1	14,14,15	0.18	0	17,19,21	0.58	0
4	NAG	A	1301	1	14,14,15	0.39	0	17,19,21	0.39	0
4	NAG	B	1304	1	14,14,15	0.22	0	17,19,21	0.49	0
4	NAG	C	1307	1	14,14,15	0.37	0	17,19,21	0.36	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '2' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	B	1305	1	-	0/6/23/26	0/1/1/1
4	NAG	C	1302	1	-	1/6/23/26	0/1/1/1
4	NAG	B	1303	1	-	0/6/23/26	0/1/1/1
4	NAG	C	1306	1	-	0/6/23/26	0/1/1/1
4	NAG	B	1302	1	-	1/6/23/26	0/1/1/1
4	NAG	B	1308	1	-	3/6/23/26	0/1/1/1
4	NAG	A	1307	1	-	0/6/23/26	0/1/1/1
4	NAG	A	1303	1	-	0/6/23/26	0/1/1/1
4	NAG	A	1308	1	-	3/6/23/26	0/1/1/1
4	NAG	A	1306	1	-	0/6/23/26	0/1/1/1
4	NAG	C	1308	1	-	2/6/23/26	0/1/1/1
4	NAG	C	1305	1	-	0/6/23/26	0/1/1/1
4	NAG	C	1303	1	-	0/6/23/26	0/1/1/1
4	NAG	A	1304	1	-	0/6/23/26	0/1/1/1
4	NAG	C	1304	1	-	0/6/23/26	0/1/1/1
4	NAG	C	1301	1	-	2/6/23/26	0/1/1/1
4	NAG	A	1302	1	-	1/6/23/26	0/1/1/1
4	NAG	A	1305	1	-	0/6/23/26	0/1/1/1
4	NAG	B	1301	1	-	2/6/23/26	0/1/1/1

Continued on next page...

*Continued from previous page...*

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	B	1307	1	-	0/6/23/26	0/1/1/1
4	NAG	B	1306	1	-	0/6/23/26	0/1/1/1
4	NAG	A	1301	1	-	2/6/23/26	0/1/1/1
4	NAG	B	1304	1	-	0/6/23/26	0/1/1/1
4	NAG	C	1307	1	-	0/6/23/26	0/1/1/1

There are no bond length outliers.

The worst 5 of 9 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	1308	NAG	C1-O5-C5	5.04	119.02	112.19
4	C	1308	NAG	C1-O5-C5	5.03	119.00	112.19
4	B	1308	NAG	C1-O5-C5	4.99	118.96	112.19
4	C	1308	NAG	O5-C5-C6	3.58	112.82	107.20
4	C	1308	NAG	C2-N2-C7	-3.29	118.22	122.90

There are no chirality outliers.

5 of 17 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	1308	NAG	C3-C2-N2-C7
4	A	1308	NAG	C8-C7-N2-C2
4	A	1308	NAG	O7-C7-N2-C2
4	B	1308	NAG	C3-C2-N2-C7
4	B	1308	NAG	C8-C7-N2-C2

There are no ring outliers.

9 monomers are involved in 10 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	B	1305	NAG	1	0
4	C	1306	NAG	1	0
4	A	1307	NAG	1	0
4	A	1306	NAG	1	0
4	C	1305	NAG	1	0
4	A	1305	NAG	1	0
4	B	1307	NAG	1	0
4	B	1306	NAG	1	0
4	C	1307	NAG	2	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


## 6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-31639. These allow visual inspection of the internal detail of the map and identification of artifacts.

No raw map or half-maps were deposited for this entry and therefore no images, graphs, etc. pertaining to the raw map can be shown.

### 6.1 Orthogonal projections [i](#)


#### 6.1.1 Primary map



The images above show the map projected in three orthogonal directions.

### 6.2 Central slices [i](#)


#### 6.2.1 Primary map




The images above show central slices of the map in three orthogonal directions.

## 6.3 Largest variance slices [i](#)


### 6.3.1 Primary map



X Index: 196



Y Index: 207




Z Index: 200


The images above show the largest variance slices of the map in three orthogonal directions.

## 6.4 Orthogonal standard-deviation projections (False-color) [i](#)


### 6.4.1 Primary map



X



Y




Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.

## 6.5 Orthogonal surface views [i](#)

### 6.5.1 Primary map



X



Y



Z

The images above show the 3D surface view of the map at the recommended contour level 0.01. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.


## 6.6 Mask visualisation [i](#)

This section was not generated. No masks/segmentation were deposited.

## 7 Map analysis [i](#)

This section contains the results of statistical analysis of the map.


### 7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.




## 7.2 Volume estimate [\(i\)](#)



The volume at the recommended contour level is 300 nm<sup>3</sup>; this corresponds to an approximate mass of 271 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

### 7.3 Rotationally averaged power spectrum [\(i\)](#)



\*Reported resolution corresponds to spatial frequency of 0.294 Å<sup>-1</sup>


## 8 Fourier-Shell correlation

This section was not generated. No FSC curve or half-maps provided.

## 9 Map-model fit [i](#)


This section contains information regarding the fit between EMDB map EMD-31639 and PDB model 7V2A. Per-residue inclusion information can be found in section 3 on page 7.

### 9.1 Map-model overlay [i](#)



The images above show the 3D surface view of the map at the recommended contour level 0.01 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

## 9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

## 9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.01).





















## 9.4 Atom inclusion [i](#)



At the recommended contour level, 98% of all backbone atoms, 91% of all non-hydrogen atoms, are inside the map.

## 9.5 Map-model fit summary

The table lists the average atom inclusion at the recommended contour level (0.01) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	 0.9100	 0.4820
A	 0.9180	 0.5010
B	 0.9170	 0.4990
C	 0.9170	 0.4990
D	 0.8770	 0.4070
E	 0.8800	 0.4040
F	 0.8730	 0.4050
G	 0.8810	 0.4100
H	 0.8740	 0.4070
I	 0.8770	 0.4010

