



Full wwPDB EM Validation Report ⓘ

Dec 10, 2023 – 12:47 AM JST

PDB ID : 8JH2
EMDB ID : EMD-36251
Title : RNA polymerase II elongation complex bound with Elf1, Spt4/5 and foreign DNA, stalled at SHL(-1) of the nucleosome
Authors : Akatsu, M.; Fujita, R.; Ogasawara, M.; Ehara, H.; Kujirai, T.; Takizawa, Y.; Sekine, S.; Kurumizaka, H.
Deposited on : 2023-05-22
Resolution : 5.70 Å(reported)

This is a Full wwPDB EM Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:


EMDB validation analysis : 0.0.1.dev70
MolProbity : 4.02b-467
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
MapQ : 1.9.9
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36

1 Overall quality at a glance

The following experimental techniques were used to determine the structure:
ELECTRON MICROSCOPY

The reported resolution of this entry is 5.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)
Ramachandran outliers	154571	4023
Sidechain outliers	154315	3826
RNA backbone	4643	859

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	0	40	
2	1	40	
3	A	1743	
4	B	1227	
5	C	304	
6	D	186	
7	E	214	
8	F	155	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
9	G	171	98% 6% 8%
10	H	145	89% 8%
11	I	115	96% 8%
12	J	72	92% 8%
13	K	118	96% 8%
14	L	72	62% 38%
15	M	110	58% 42%
16	N	228	52% 48%
17	P	16	94% 6%
18	T	228	56% 44%
19	V	114	89% 8% 11%
20	W	908	30% 70%
21	a	136	70% 8% 29%
21	e	136	73% 27%
22	b	103	75% 25%
22	f	103	78% 22%
23	c	130	76% 8% 22%
23	g	130	77% 8% 22%
24	d	126	69% 8% 30%
24	h	126	70% 8% 29%

2 Entry composition [i](#)

There are 26 unique types of molecules in this entry. The entry contains 47629 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a DNA chain called DNA (40-MER).

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	P		
1	0	40	823	389	175	219	40	0	0

- Molecule 2 is a DNA chain called DNA (40-MER).

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	P		
2	1	40	817	394	122	261	40	0	0

- Molecule 3 is a protein called DNA-directed RNA polymerase subunit.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
3	A	1411	11116	7009	1937	2100	70	0	0

- Molecule 4 is a protein called DNA-directed RNA polymerase subunit beta.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
4	B	1157	9228	5816	1630	1724	58	0	0

- Molecule 5 is a protein called RNA polymerase II third largest subunit B44, part of central core.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
5	C	263	2098	1319	354	413	12	0	0

- Molecule 6 is a protein called RNA polymerase II subunit B32.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
6	D	168	1314	812	237	263	2	0	0

- Molecule 7 is a protein called RNA polymerase subunit ABC27.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
7	E	213	Total	C	N	O	S	0	0
			1740	1094	312	324	10		

- Molecule 8 is a protein called RNA polymerase subunit ABC23, common to RNA polymerases I, II, and III.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
8	F	84	Total	C	N	O	S	0	0
			677	429	114	131	3		

- Molecule 9 is a protein called RNA polymerase II subunit.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
9	G	171	Total	C	N	O	S	0	0
			1324	858	214	247	5		

- Molecule 10 is a protein called DNA-directed RNA polymerases I, II, and III subunit RPABC3.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
10	H	133	Total	C	N	O	S	0	0
			1052	671	169	208	4		

- Molecule 11 is a protein called DNA-directed RNA polymerase subunit.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
11	I	111	Total	C	N	O	S	0	0
			917	565	161	180	11		

- Molecule 12 is a protein called RNA polymerase subunit ABC10-beta, common to RNA polymerases I, II, and III.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
12	J	66	Total	C	N	O	S	0	0
			545	349	95	95	6		

- Molecule 13 is a protein called RNA polymerase II subunit B12.5.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
13	K	113	Total	C	N	O	S	0	0
			932	599	160	169	4		

- Molecule 14 is a protein called RNA polymerase subunit ABC10-alpha.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
14	L	45	359	221	72	61	5	0	0

- Molecule 15 is a protein called Transcription elongation factor 1 homolog.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
15	M	64	505	318	82	99	6	0	0

- Molecule 16 is a DNA chain called DNA (218-MER).

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	P		
16	N	118	2426	1148	445	715	118	0	0

- Molecule 17 is a RNA chain called RNA (5'-R(P*CP*CP*UP*GP*GP*UP*GP*UP*CP*UP*UP*GP*GP*GP*UP*G)-3').

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	P		
17	P	16	341	151	56	118	16	0	0

- Molecule 18 is a DNA chain called DNA (218-MER).

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	P		
18	T	127	2589	1227	486	750	126	0	0

- Molecule 19 is a protein called Transcription elongation factor SPT4.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
19	V	102	792	492	143	150	7	0	0

- Molecule 20 is a protein called Transcription elongation factor SPT5.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
20	W	275	2226	1425	397	403	1	0	0

- Molecule 21 is a protein called Histone H3.3.

Mol	Chain	Residues	Atoms					AltConf	Trace
21	a	96	Total	C	N	O	S	0	0
			786	497	151	136	2		
21	e	99	Total	C	N	O	S	0	0
			815	515	159	139	2		

- Molecule 22 is a protein called Histone H4.

Mol	Chain	Residues	Atoms					AltConf	Trace
22	b	77	Total	C	N	O	S	0	0
			614	389	119	105	1		
22	f	80	Total	C	N	O	S	0	0
			638	401	125	111	1		

- Molecule 23 is a protein called Histone H2A type 1-B/E.

Mol	Chain	Residues	Atoms				AltConf	Trace
23	c	101	Total	C	N	O	0	0
			778	492	150	136		
23	g	101	Total	C	N	O	0	0
			778	492	150	136		

- Molecule 24 is a protein called Histone H2B type 1-J.

Mol	Chain	Residues	Atoms					AltConf	Trace
24	d	88	Total	C	N	O	S	0	0
			685	433	121	129	2		
24	h	90	Total	C	N	O	S	0	0
			703	444	124	133	2		

- Molecule 25 is ZINC ION (three-letter code: ZN) (formula: Zn) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms		AltConf
25	A	2	Total	Zn	0
			2	2	
25	B	1	Total	Zn	0
			1	1	
25	C	1	Total	Zn	0
			1	1	
25	I	2	Total	Zn	0
			2	2	

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		AltConf
25	J	1	Total 1	Zn 1	0
25	L	1	Total 1	Zn 1	0
25	M	1	Total 1	Zn 1	0
25	V	1	Total 1	Zn 1	0


- Molecule 26 is MAGNESIUM ION (three-letter code: MG) (formula: Mg) (labeled as "Ligand of Interest" by depositor).


Mol	Chain	Residues	Atoms		AltConf
26	A	1	Total 1	Mg 1	0

Chain B:  93% 6%




- Molecule 5: RNA polymerase II third largest subunit B44, part of central core

Chain C:  86% 13%




- Molecule 6: RNA polymerase II subunit B32

Chain D:  6% 90% 10%




- Molecule 7: RNA polymerase subunit ABC27

Chain E:  96% 1%



- Molecule 8: RNA polymerase subunit ABC23, common to RNA polymerases I, II, and III


Chain F:  54% 46% 1%



- Molecule 9: RNA polymerase II subunit


Chain G:  6% 98% 1%





- Molecule 10: DNA-directed RNA polymerases I, II, and III subunit RPABC3

Chain H: 89% 8%




- Molecule 11: DNA-directed RNA polymerase subunit

Chain I: 96%




- Molecule 12: RNA polymerase subunit ABC10-beta, common to RNA polymerases I, II, and III

Chain J: 92% 8%




- Molecule 13: RNA polymerase II subunit B12.5

Chain K: 96%




- Molecule 14: RNA polymerase subunit ABC10-alpha

Chain L: 62% 38%



- Molecule 15: Transcription elongation factor 1 homolog

Chain M: 58% 42%




- Molecule 16: DNA (218-MER)


Chain N: 52% 48%




- Molecule 23: Histone H2A type 1-B/E



- Molecule 24: Histone H2B type 1-J



- Molecule 24: Histone H2B type 1-J



4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	74079	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	59.84	Depositor
Minimum defocus (nm)	1000	Depositor
Maximum defocus (nm)	2500	Depositor
Magnification	Not provided	
Image detector	GATAN K3 BIOQUANTUM (6k x 4k)	Depositor
Maximum map value	0.025	Depositor
Minimum map value	-0.004	Depositor
Average map value	0.000	Depositor
Map value standard deviation	0.001	Depositor
Recommended contour level	0.00307	Depositor
Map size (\AA)	381.59998, 381.59998, 381.59998	wwPDB
Map dimensions	360, 360, 360	wwPDB
Map angles ($^\circ$)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (\AA)	1.06, 1.06, 1.06	Depositor

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	0	0.49	0/930	0.76	0/1430
2	1	0.61	0/908	0.99	0/1402
3	A	0.23	0/11322	0.44	0/15300
4	B	0.23	0/9407	0.47	0/12685
5	C	0.23	0/2139	0.43	0/2895
6	D	0.23	0/1326	0.45	0/1788
7	E	0.23	0/1772	0.46	0/2385
8	F	0.22	0/687	0.46	0/931
9	G	0.25	0/1353	0.43	0/1837
10	H	0.24	0/1069	0.46	0/1444
11	I	0.24	0/934	0.49	0/1257
12	J	0.23	0/554	0.45	0/742
13	K	0.23	0/953	0.44	0/1291
14	L	0.23	0/365	0.53	0/484
15	M	0.23	0/513	0.38	0/693
16	N	0.51	0/2719	0.87	0/4195
17	P	0.12	0/379	0.68	0/589
18	T	0.48	0/2905	0.82	0/4477
19	V	0.23	0/808	0.47	0/1097
20	W	0.23	0/2267	0.46	0/3048
21	a	0.23	0/798	0.49	0/1071
21	e	0.23	0/827	0.49	0/1108
22	b	0.24	0/621	0.51	0/832
22	f	0.23	0/645	0.52	0/862
23	c	0.23	0/788	0.48	0/1065
23	g	0.23	0/788	0.48	0/1065
24	d	0.23	0/696	0.41	0/938
24	h	0.23	0/714	0.41	0/961
All	All	0.29	0/49187	0.55	0/67872

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

Due to software issues we are unable to calculate clashes - this section is therefore empty.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
3	A	1399/1743 (80%)	1301 (93%)	90 (6%)	8 (1%)	25	65
4	B	1145/1227 (93%)	1064 (93%)	75 (7%)	6 (0%)	29	69
5	C	261/304 (86%)	243 (93%)	18 (7%)	0	100	100
6	D	162/186 (87%)	149 (92%)	12 (7%)	1 (1%)	25	65
7	E	211/214 (99%)	203 (96%)	5 (2%)	3 (1%)	11	46
8	F	82/155 (53%)	78 (95%)	4 (5%)	0	100	100
9	G	169/171 (99%)	158 (94%)	9 (5%)	2 (1%)	13	49
10	H	129/145 (89%)	126 (98%)	2 (2%)	1 (1%)	19	60
11	I	109/115 (95%)	95 (87%)	14 (13%)	0	100	100
12	J	64/72 (89%)	59 (92%)	5 (8%)	0	100	100
13	K	111/118 (94%)	107 (96%)	4 (4%)	0	100	100
14	L	43/72 (60%)	37 (86%)	6 (14%)	0	100	100
15	M	62/110 (56%)	55 (89%)	7 (11%)	0	100	100
19	V	100/114 (88%)	94 (94%)	6 (6%)	0	100	100
20	W	265/908 (29%)	238 (90%)	23 (9%)	4 (2%)	10	45
21	a	94/136 (69%)	91 (97%)	3 (3%)	0	100	100
21	e	97/136 (71%)	95 (98%)	2 (2%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
22	b	75/103 (73%)	70 (93%)	5 (7%)	0	100	100
22	f	78/103 (76%)	73 (94%)	5 (6%)	0	100	100
23	c	99/130 (76%)	97 (98%)	2 (2%)	0	100	100
23	g	99/130 (76%)	96 (97%)	3 (3%)	0	100	100
24	d	86/126 (68%)	82 (95%)	3 (4%)	1 (1%)	13	49
24	h	88/126 (70%)	84 (96%)	3 (3%)	1 (1%)	14	51
All	All	5028/6644 (76%)	4695 (93%)	306 (6%)	27 (0%)	32	69

All (27) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
3	A	948	VAL
4	B	830	TYR
20	W	806	PRO
3	A	253	MET
3	A	255	GLU
4	B	555	GLY
20	W	320	LEU
20	W	776	ASP
3	A	887	ILE
4	B	352	GLU
6	D	10	ALA
9	G	46	VAL
10	H	108	GLU
3	A	47	ARG
3	A	960	VAL
4	B	257	THR
4	B	1039	GLY
20	W	800	SER
3	A	179	GLY
4	B	1017	ILE
7	E	113	ASN
24	h	103	PRO
3	A	1109	VAL
9	G	154	VAL
7	E	123	ILE
7	E	150	PRO
24	d	103	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
3	A	1224/1528 (80%)	1220 (100%)	4 (0%)	92	94
4	B	1012/1077 (94%)	1000 (99%)	12 (1%)	71	83
5	C	236/264 (89%)	235 (100%)	1 (0%)	91	94
6	D	143/160 (89%)	143 (100%)	0	100	100
7	E	196/197 (100%)	192 (98%)	4 (2%)	55	74
8	F	75/137 (55%)	75 (100%)	0	100	100
9	G	148/148 (100%)	147 (99%)	1 (1%)	84	90
10	H	120/130 (92%)	117 (98%)	3 (2%)	47	68
11	I	106/109 (97%)	105 (99%)	1 (1%)	78	87
12	J	60/66 (91%)	60 (100%)	0	100	100
13	K	104/109 (95%)	104 (100%)	0	100	100
14	L	38/56 (68%)	38 (100%)	0	100	100
15	M	61/98 (62%)	61 (100%)	0	100	100
19	V	86/97 (89%)	86 (100%)	0	100	100
20	W	241/795 (30%)	241 (100%)	0	100	100
21	a	82/110 (74%)	81 (99%)	1 (1%)	71	83
21	e	85/110 (77%)	85 (100%)	0	100	100
22	b	63/79 (80%)	63 (100%)	0	100	100
22	f	65/79 (82%)	65 (100%)	0	100	100
23	c	80/100 (80%)	78 (98%)	2 (2%)	47	68
23	g	80/100 (80%)	79 (99%)	1 (1%)	69	82
24	d	75/105 (71%)	75 (100%)	0	100	100
24	h	77/105 (73%)	76 (99%)	1 (1%)	69	82
All	All	4457/5759 (77%)	4426 (99%)	31 (1%)	84	90

All (31) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
3	A	200	ARG
3	A	446	ASN
3	A	454	MET
3	A	661	ASN
4	B	278	PHE
4	B	284	VAL
4	B	394	PHE
4	B	466	MET
4	B	500	LYS
4	B	653	ARG
4	B	764	SER
4	B	904	ARG
4	B	992	VAL
4	B	1094	ARG
4	B	1122	ARG
4	B	1201	LYS
5	C	8	ASN
7	E	4	ASN
7	E	83	ASP
7	E	120	ASN
7	E	166	ARG
9	G	87	VAL
10	H	8	ASP
10	H	85	ASP
10	H	117	PHE
11	I	105	ASN
21	a	40	ARG
23	c	50	TYR
23	c	73	ASN
23	g	50	TYR
24	h	63	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (43) such sidechains are listed below:

Mol	Chain	Res	Type
3	A	172	GLN
3	A	197	GLN
3	A	287	GLN
3	A	291	ASN
3	A	364	GLN
3	A	446	ASN
3	A	649	ASN
3	A	651	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
3	A	661	ASN
3	A	787	HIS
3	A	936	GLN
3	A	961	ASN
3	A	968	ASN
3	A	1147	ASN
3	A	1190	GLN
3	A	1435	GLN
4	B	291	GLN
4	B	438	ASN
4	B	776	GLN
4	B	996	HIS
5	C	8	ASN
5	C	13	GLN
6	D	38	GLN
6	D	46	HIS
7	E	4	ASN
7	E	120	ASN
7	E	145	HIS
7	E	193	GLN
10	H	13	GLN
11	I	12	ASN
11	I	31	ASN
11	I	90	GLN
11	I	105	ASN
15	M	65	GLN
19	V	78	GLN
20	W	434	ASN
20	W	441	ASN
20	W	783	HIS
20	W	784	ASN
21	a	108	ASN
23	c	73	ASN
23	c	104	GLN
21	e	108	ASN

5.3.3 RNA

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
17	P	15/16 (93%)	1 (6%)	0

All (1) RNA backbone outliers are listed below:

Mol	Chain	Res	Type
17	P	-4	C

There are no RNA pucker outliers to report.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 11 ligands modelled in this entry, 11 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-36251. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X




Y




Z


6.1.2 Raw map



X



Y




Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map



X Index: 180




Y Index: 180




Z Index: 180


6.2.2 Raw map



X Index: 180



Y Index: 180




Z Index: 180


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 188




Y Index: 188




Z Index: 195


6.3.2 Raw map



X Index: 186



Y Index: 185




Z Index: 196


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [i](#)


6.4.1 Primary map



X




Y




Z


6.4.2 Raw map



X



Y



Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.


6.5 Orthogonal surface views [i](#)

6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.00307. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

6.5.2 Raw map



These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.


6.6 Mask visualisation [i](#)

This section was not generated. No masks/segmentation were deposited.

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [i](#)



The volume at the recommended contour level is 1603 nm^3 ; this corresponds to an approximate mass of 1448 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum [i](#)




*Reported resolution corresponds to spatial frequency of 0.175 Å⁻¹

8 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC [i](#)



*Reported resolution corresponds to spatial frequency of 0.175 Å⁻¹

8.2 Resolution estimates [i](#)


Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	5.70	-	-
Author-provided FSC curve	-	-	-
Unmasked-calculated*	7.49	9.24	7.62

*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 7.49 differs from the reported value 5.7 by more than 10 %

9 Map-model fit [i](#)


This section contains information regarding the fit between EMDB map EMD-36251 and PDB model 8JH2. Per-residue inclusion information can be found in section 3 on page 9.

9.1 Map-model overlay [i](#)




The images above show the 3D surface view of the map at the recommended contour level 0.00307 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.00307).



























































9.4 Atom inclusion [i](#)



At the recommended contour level, 98% of all backbone atoms, 97% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary

The table lists the average atom inclusion at the recommended contour level (0.00307) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	 0.9740	 0.1450
0	 0.8540	 0.0410
1	 0.8000	 0.0240
A	 0.9910	 0.1950
B	 0.9910	 0.1980
C	 0.9900	 0.2100
D	 0.9030	 0.0840
E	 0.9940	 0.1840
F	 0.9990	 0.1890
G	 0.9320	 0.0990
H	 0.9810	 0.2210
I	 0.9870	 0.1020
J	 1.0000	 0.1980
K	 0.9920	 0.1840
L	 1.0000	 0.2090
M	 0.9740	 0.0510
N	 0.9960	 0.0820
P	 1.0000	 0.1830
T	 0.9780	 0.1150
V	 0.8820	 0.0460
W	 0.8850	 0.0560
a	 0.9920	 0.0700
b	 1.0000	 0.0430
c	 0.9960	 0.0380
d	 1.0000	 0.0320
e	 1.0000	 0.0750
f	 0.9950	 0.0640
g	 0.9800	 0.0430
h	 0.9960	 0.0510

