

# wwPDB X-ray Structure Validation Summary Report (i)

#### Aug 26, 2024 – 06:04 pm BST

PDB ID : 8QBZ

Title: Crystal structure of the outer membrane decaheme cytochrome MtrC

(E344Boc-Lys)

Authors: Edwards, M.J.; Lockwood, C.J.; Whiting, K.; Nash, B.W.; Butt, J.N.; Clarke,

T.A.

Deposited on : 2023-08-25

Resolution : 1.90 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity: 4.02b-467

Mogul : 1.8.4, CSD as541be (2020)

Xtriage (Phenix) : 1.13

EDS : 3.0

buster-report : 1.1.7 (2018)

Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)

CCP4 : 9.0.002 (Gargrove)

 $Density-Fitness \quad : \quad 1.0.11$ 

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.38.2

## 1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:  $X\text{-}RAY\ DIFFRACTION$ 

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive	Similar resolution
Metric	$(\#  ext{Entries})$	$(\#  ext{Entries},  ext{ resolution range}( ext{Å}))$
$R_{free}$	164625	7293 (1.90-1.90)
Clashscore	180529	8090 (1.90-1.90)
Ramachandran outliers	177936	8022 (1.90-1.90)
Sidechain outliers	177891	8022 (1.90-1.90)
RSRZ outliers	164620	7292 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain					
1	A	679	83%	9%	8%			



# 2 Entry composition (i)

There are 6 unique types of molecules in this entry. The entry contains 6178 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Extracellular iron oxide respiratory system surface decaheme cytochrome c component MtrC.

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace		
1	Λ	626	Total	С	N	О	S	0	7	0
1	A	020	4748	2938	819	958	33			

There are 34 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	3	MET	-	initiating methionine	UNP Q8EG34
A	4	LYS	-	expression tag	UNP Q8EG34
A	5	PHE	-	expression tag	UNP Q8EG34
A	6	LYS	-	expression tag	UNP Q8EG34
A	7	LEU	-	expression tag	UNP Q8EG34
A	8	ASN	-	expression tag	UNP Q8EG34
A	9	LEU	-	expression tag	UNP Q8EG34
A	10	ILE	-	expression tag	UNP Q8EG34
A	11	THR	-	expression tag	UNP Q8EG34
A	12	LEU	-	expression tag	UNP Q8EG34
A	13	ALA	-	expression tag	UNP Q8EG34
A	14	LEU	-	expression tag	UNP Q8EG34
A	15	LEU	-	expression tag	UNP Q8EG34
A	16	ALA	-	expression tag	UNP Q8EG34
A	17	ASN	-	expression tag	UNP Q8EG34
A	18	THR	-	expression tag	UNP Q8EG34
A	19	GLY	-	expression tag	UNP Q8EG34
A	20	LEU	-	expression tag	UNP Q8EG34
A	21	ALA	-	expression tag	UNP Q8EG34
A	22	VAL	-	expression tag	UNP Q8EG34
A	23	ALA		expression tag	UNP Q8EG34
A	24	ALA	=	expression tag	UNP Q8EG34
A	25	ASP	-	expression tag	UNP Q8EG34
A	344	LBY	GLU	engineered mutation	UNP Q8EG34
A	672	SER	-	expression tag	UNP Q8EG34
A	673	ALA	-	expression tag	UNP Q8EG34



Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	674	TRP	-	expression tag	UNP Q8EG34
A	675	SER	-	expression tag	UNP Q8EG34
A	676	HIS	-	expression tag	UNP Q8EG34
A	677	PRO	-	expression tag	UNP Q8EG34
A	678	GLN	-	expression tag	UNP Q8EG34
A	679	PHE	-	expression tag	UNP Q8EG34
A	680	GLU	-	expression tag	UNP Q8EG34
A	681	LYS	-	expression tag	UNP Q8EG34

• Molecule 2 is HEME C (three-letter code: HEC) (formula:  $C_{34}H_{34}FeN_4O_4$ ) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues		Ato	oms			ZeroOcc	AltConf	
2	2 A	1	Total	С	Fe	N	О	0	0	
	A	1	43	34	1	4	4	0	U	
2	A	1	Total	С	Fe	N	О	0	0	
	A	1	43	34	1	4	4		U	
2	A	1	Total	С	Fe	N	О	0	0	
2	Λ	1	43	34	1	4	4			
2	A	1	Total	С	Fe	N	Ο	0	0	
	Λ	1	43	34	1	4	4		U	
2	A	1	Total	С	Fe	N	Ο	0	0	
	Λ	1	43	34	1	4	4	0	U	
2	A	1	Total	С	Fe	N	Ο	0	0	
	Λ	1	43	34	1	4	4	0		
2	A	1	Total	С	Fe	N	О	0	0	
2	A		43	34	1	4	4			




Mol	Chain	Residues		Ato	oms			ZeroOcc	AltConf	
2	Λ	1	Total	С	Fe	N	О	0	0	
2	A	1	43	34	1	4	4	0	0	
9	٨	1	Total	С	Fe	N	О	0	0	
2	A	1	43	34	1	4	4		U	
9	Λ	1	Total	С	Fe	N	О	0	0	
2	A	1	43	34	1	4	4			

• Molecule 3 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	5	Total Ca 5 5	0	0

 $\bullet$  Molecule 4 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula:  $\mathrm{C_2H_6O_2}).$ 




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C O 4 2 2	0	0
4	A	1	Total C O 4 2 2	0	0
4	A	1	Total C O 4 2 2	0	0
4	A	1	Total C O 4 2 2	0	0
4	A	1	Total C O 4 2 2	0	0



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C O 4 2 2	0	0
4	A	1	Total C O 4 2 2	0	0
4	A	1	Total C O 4 2 2	0	0

 $\bullet$  Molecule 5 is ACETATE ION (three-letter code: ACT) (formula:  $\mathrm{C_2H_3O_2}).$ 



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total C O 4 2 2	0	0

• Molecule 6 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	959	Total O 959 959	0	0



# 3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Extracellular iron oxide respiratory system surface decaheme cytochrome c component MtrC





# 4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants	52.97Å 89.66Å 153.54Å	Depositor
a, b, c, $\alpha$ , $\beta$ , $\gamma$	90.00° 90.00° 90.00°	Depositor
Resolution (Å)	50.12 - 1.90	Depositor
recontion (11)	50.12 - 1.90	EDS
% Data completeness	99.6 (50.12-1.90)	Depositor
(in resolution range)	99.9 (50.12-1.90)	EDS
$R_{merge}$	0.45	Depositor
$R_{sym}$	(Not available)	Depositor
$< I/\sigma(I) > 1$	1.55  (at  1.90Å)	Xtriage
Refinement program	REFMAC 5.8.0267	Depositor
$R, R_{free}$	0.174 , $0.216$	Depositor
it, it free	0.181 , 0.221	DCC
$R_{free}$ test set	2954  reflections  (5.05%)	wwPDB-VP
Wilson B-factor $(A^2)$	15.3	Xtriage
Anisotropy	0.107	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$ , $B_{sol}(Å^2)$	0.36, 49.3	EDS
L-test for twinning <sup>2</sup>	$ < L > = 0.48, < L^2> = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	6178	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	20.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 4.79% of the height of the origin peak. No significant pseudotranslation is detected.

<sup>&</sup>lt;sup>2</sup>Theoretical values of <|L|>,  $<L^2>$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



<sup>&</sup>lt;sup>1</sup>Intensities estimated from amplitudes.

# 5 Model quality (i)

## 5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: ACT, EDO, LBY, HEC, CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
MIOI	Chain	RMSZ	# Z  > 5	RMSZ	# Z  > 5
1	A	0.70	1/4820 (0.0%)	0.79	1/6545 (0.0%)

#### All (1) bond length outliers are listed below:

Mol	Chain	$\operatorname{Res}$	Type	Atoms	$\mathbf{Z}$	$\operatorname{Observed}(\text{\AA})$	$\operatorname{Ideal}( ext{\AA})$
1	A	541	CYS	CB-SG	-5.04	1.73	1.81

#### All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^o)$	$\operatorname{Ideal}(^{o})$
1	A	263	GLU	CB-CA-C	6.36	123.13	110.40

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4748	0	4556	43	0
2	A	430	0	300	16	0
3	A	5	0	0	0	0
4	A	32	0	48	3	0
5	A	4	0	3	0	0
6	A	959	0	0	15	2



Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
All	All	6178	0	4907	52	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 52 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	$\begin{array}{c} {\rm Interatomic} \\ {\rm distance} \ ({\rm \AA}) \end{array}$	Clash overlap (Å)
1:A:75[A]:MET:SD	6:A:1753:HOH:O	2.21	0.97
2:A:804:HEC:CGA	6:A:904:HOH:O	2.16	0.94
2:A:804:HEC:HBA1	6:A:904:HOH:O	1.84	0.75
1:A:493:MET:HE2	1:A:535[B]:LEU:HD21	1.70	0.72
1:A:493:MET:CE	1:A:535[B]:LEU:HD21	2.21	0.69

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	$\begin{array}{c} {\rm Interatomic} \\ {\rm distance} \ ({\rm \AA}) \end{array}$	Clash overlap (Å)
6:A:1536:HOH:O	6:A:1561:HOH:O[4_445]	2.18	0.02
6:A:1537:HOH:O	6:A:1539:HOH:O[4_545]	2.18	0.02

## 5.3 Torsion angles (i)

## 5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	630/679 (93%)	614 (98%)	15 (2%)	1 (0%)	44 36

All (1) Ramachandran outliers are listed below:

	Mol	Chain	Res	Type
ſ	1	A	384	ASN



#### 5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	519/549 (94%)	508 (98%)	11 (2%)	48 45

5 of 11 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	449	LYS
1	A	521	LEU
1	A	625	MET
1	A	602	LYS
1	A	198	HIS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	71	ASN
1	A	138	GLN

#### 5.3.3 RNA (i)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains (i)

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).



Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
IVIOI	туре	Chain	rtes	LIIIK	Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
1	LBY	A	344	1	14,15,16	0.33	0	14,19,21	0.42	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	LBY	A	344	1	-	0/14/15/17	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.5 Carbohydrates (i)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry (i)

Of 24 ligands modelled in this entry, 5 are monoatomic - leaving 19 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Trino	Chain	Res	Link	В	ond leng	$\operatorname{gths}$	Bond angles		
MIOI	Type	Chain	nes	Lilik	Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	HEC	A	807	1	32,50,50	2.00	7 (21%)	24,82,82	2.34	7 (29%)
2	HEC	A	810	1	32,50,50	2.06	10 (31%)	24,82,82	2.74	11 (45%)
5	ACT	A	824	-	3,3,3	1.14	0	3,3,3	0.55	0
2	HEC	A	802	1	32,50,50	1.78	8 (25%)	24,82,82	2.55	7 (29%)



Mol	Type	Chain	Res	Link	В	ond leng	$\operatorname{gths}$	В	ond ang	gles
MIOI	туре	Chain	nes	Lilik	Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	EDO	A	822	-	3,3,3	0.14	0	2,2,2	0.22	0
2	HEC	A	801	1	32,50,50	1.83	8 (25%)	24,82,82	2.81	9 (37%)
2	HEC	A	805	1	32,50,50	1.75	6 (18%)	24,82,82	2.32	6 (25%)
4	EDO	A	819	-	3,3,3	0.07	0	2,2,2	0.20	0
4	EDO	A	817	-	3,3,3	0.32	0	2,2,2	0.44	0
4	EDO	A	821	-	3,3,3	0.48	0	2,2,2	0.41	0
4	EDO	A	823	-	3,3,3	0.24	0	2,2,2	0.20	0
4	EDO	A	816	-	3,3,3	0.01	0	2,2,2	0.12	0
2	HEC	A	806	1	32,50,50	1.87	10 (31%)	24,82,82	2.16	7 (29%)
2	HEC	A	803	1	32,50,50	1.85	8 (25%)	24,82,82	2.08	6 (25%)
2	HEC	A	809	1	32,50,50	1.79	9 (28%)	24,82,82	2.37	6 (25%)
4	EDO	A	818	-	3,3,3	0.30	0	2,2,2	0.38	0
2	HEC	A	808	1	32,50,50	1.71	9 (28%)	24,82,82	2.25	7 (29%)
2	HEC	A	804	1	32,50,50	1.90	8 (25%)	24,82,82	2.52	9 (37%)
4	EDO	A	820	-	3,3,3	0.31	0	2,2,2	0.32	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	HEC	A	805	1	-	0/10/54/54	-
4	EDO	A	819	-	-	0/1/1/1	-
4	EDO	A	817	-	-	1/1/1/1	-
4	EDO	A	818	-	-	1/1/1/1	-
2	HEC	A	810	1	-	0/10/54/54	_
2	HEC	A	807	1	-	4/10/54/54	-
4	EDO	A	821	-	-	1/1/1/1	-
2	HEC	A	808	1	-	4/10/54/54	-
4	EDO	A	823	-	-	0/1/1/1	-
4	EDO	A	822	-	-	0/1/1/1	-
4	EDO	A	816	-	-	0/1/1/1	_
2	HEC	A	802	1	-	2/10/54/54	-
2	HEC	A	806	1	-	4/10/54/54	-
2	HEC	A	804	1	-	3/10/54/54	-
2	HEC	A	803	1	-	2/10/54/54	-
2	HEC	A	801	1	-	3/10/54/54	-
2	HEC	A	809	1	-	2/10/54/54	-



$\mathbf{Mol}$	Type	Chain	$\operatorname{Res}$	Link	Chirals	Torsions	Rings
4	EDO	A	820	-	-	1/1/1/1	-

The worst 5 of 83 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(A)	$Ideal(\AA)$
2	A	807	HEC	C2B-C3B	5.84	1.46	1.40
2	A	806	HEC	C3C-C2C	5.52	1.46	1.40
2	A	810	HEC	C3C-C2C	5.46	1.46	1.40
2	A	810	HEC	C2B-C3B	5.33	1.46	1.40
2	A	803	HEC	C3C-C2C	5.02	1.46	1.40

The worst 5 of 75 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	$\mathbf{Z}$	$Observed(^o)$	$Ideal(^{o})$
2	A	810	HEC	C1D-C2D-C3D	-8.23	101.27	107.00
2	A	801	HEC	C1D-C2D-C3D	-7.65	101.68	107.00
2	A	809	HEC	C1D-C2D-C3D	-6.88	102.21	107.00
2	A	808	HEC	C1D-C2D-C3D	-6.82	102.25	107.00
2	A	806	HEC	C1D-C2D-C3D	-6.72	102.32	107.00

There are no chirality outliers.

5 of 28 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	801	HEC	C2A-CAA-CBA-CGA
4	A	817	EDO	O1-C1-C2-O2
4	A	821	EDO	O1-C1-C2-O2
4	A	820	EDO	O1-C1-C2-O2
2	A	804	HEC	C1A-C2A-CAA-CBA

There are no ring outliers.

10 monomers are involved in 17 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	807	HEC	2	0
2	A	802	HEC	1	0
2	A	801	HEC	2	0
2	A	805	HEC	1	0
4	A	821	EDO	3	0
2	A	806	HEC	2	0
2	A	803	HEC	2	0




Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	809	HEC	2	0
2	A	808	HEC	1	0
2	A	804	HEC	5	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.

















































## 5.7 Other polymers (i)

There are no such residues in this entry.

## 5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



## 6 Fit of model and data (i)

## 6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median,  $95^{th}$  percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	$\langle { m RSRZ} \rangle$	# RSRZ > 2		$OWAB(Å^2)$	Q<0.9	
1	A	625/679 (92%)	-0.42	1 (0%)	92	92	5, 16, 32, 60	7 (1%)

#### All (1) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	241	LYS	2.4

## 6.2 Non-standard residues in protein, DNA, RNA chains (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median,  $95^{th}$  percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-}\mathbf{factors}(\mathbf{\mathring{A}}^2)$	Q < 0.9
1	LBY	A	344	16/17	0.90	0.11	21,29,40,40	0

## 6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

## 6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median,  $95^{th}$  percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	$\operatorname{Res}$	Atoms	RSCC	RSR	$\mathbf{B} ext{-}\mathbf{factors}(\mathbf{\mathring{A}}^2)$	Q<0.9
5	ACT	A	824	4/4	0.70	0.22	44,50,51,51	0




Continued from previous page...


Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-}\mathbf{factors}(\mathbf{\mathring{A}}^2)$	Q<0.9
4	EDO	A	821	4/4	0.77	0.16	36,37,37,37	0
4	EDO	A	817	4/4	0.82	0.14	21,23,25,30	0
4	EDO	A	823	4/4	0.83	0.15	37,38,39,39	0
4	EDO	A	820	4/4	0.85	0.13	32,34,34,35	0
4	EDO	A	816	4/4	0.88	0.12	36,37,38,39	0
4	EDO	A	818	4/4	0.93	0.08	19,21,23,23	0
4	EDO	A	819	4/4	0.93	0.09	32,32,33,33	0
4	EDO	A	822	4/4	0.96	0.06	24,24,25,26	0
3	CA	A	814	1/1	0.97	0.12	35,35,35,35	0
2	HEC	A	805	43/43	0.98	0.06	13,15,20,24	0
2	HEC	A	806	43/43	0.98	0.07	13,14,24,41	0
2	HEC	A	809	43/43	0.98	0.06	10,11,21,30	0
2	HEC	A	801	43/43	0.98	0.07	10,12,26,32	0
3	CA	A	815	1/1	0.98	0.12	27,27,27,27	0
2	HEC	A	802	43/43	0.98	0.05	10,11,15,18	0
2	HEC	A	804	43/43	0.98	0.07	11,13,28,35	0
3	CA	A	813	1/1	0.99	0.03	16,16,16,16	0
2	HEC	A	807	43/43	0.99	0.05	12,13,18,24	0
2	HEC	A	808	43/43	0.99	0.05	9,10,12,13	0
2	HEC	A	803	43/43	0.99	0.05	9,10,11,12	0
2	HEC	A	810	43/43	0.99	0.05	10,12,17,20	0
3	CA	A	812	1/1	0.99	0.02	10,10,10,10	1
3	CA	A	811	1/1	1.00	0.03	11,11,11,11	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

















































# 6.5 Other polymers (i)

There are no such residues in this entry.

